

Review

Jonathan Zhang

1 Differential Calculus in Banach Spaces and the Calculus of Variations

1.1 Differentiation

Our goal will be to generalize the notion of the 1D derivative, in the context of results like the Mean Value Theorem and Taylor's Theorem. Let us start with what we know: the standard 1D definition of the derivative is

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}, \quad (1)$$

but it is not clear how $1/h$ will generalize in higher dimensions. Indeed, it makes no sense. We can, however, define a similar concept.

Definition 1.1 (Little o). Let X, Y be NLS's, $f : X \rightarrow Y$. If

$$\lim_{h \rightarrow 0} \frac{\|f(h)\|_Y}{\|h\|_X} = 0,$$

we say that f is “little oh” of h and write

$$\|f(h)\|_Y = o(\|h\|_X).$$

Here, we have generalized the notion of $1/h$ to multiple dimensions by looking only at the norm of the increment and of the output. We can now formulate a notion of the derivative.

Definition 1.2 (Fréchet Derivative). Let X, Y NLS, $U \subset X$ open, $f : U \rightarrow Y$. We say that f is Fréchet differentiable (or strongly differentiable) at a point $x \in U$ if and only if there exists a bounded linear map $A \in B(X, Y)$ such that the remainder

$$R(x, h) := f(x+h) - f(x) - Ah$$

is $o(\|h\|_X)$. That is,

$$\lim_{h \rightarrow 0} \frac{\|R(x, h)\|_Y}{\|h\|_X} = 0.$$

If A exists, we call A the Fréchet derivative of f at x , denoted $Df(x)$. Note that by construction, $A = Df(x)$ is a linear operator on h . The definition of the Fréchet derivative is consistent with the one dimensional analogue. For $f \in C^1(\mathbb{R})$,

$$R(x, h) = f(x+h) - f(x) - f'(x)h = \left[\frac{f(x+h) - f(x)}{h} - f'(x) \right] h, \quad (2)$$

and

$$\lim_{h \rightarrow 0} \frac{|R(x, h)|}{\|h\|} = \lim_{h \rightarrow 0} \left[\frac{f(x+h) - f(x)}{\|h\|} - f'(x) \right] = 0. \quad (3)$$

More generally, the map Df (as contrasted with $Df(x)$) can be regarded as a map $Df : U \times X \ni (x, h) \rightarrow Df(x)h \in Y$.

Proposition 1.1. *If f is Fréchet differentiable, then $Df(x)$ is unique, and f is continuous at x . Moreover, if g is Fréchet differentiable, then for every $\alpha, \beta \in \mathbb{F}$,*

$$D(\alpha f + \beta g)(x) = \alpha Df(x) + \beta Dg(x).$$

(That is to say, the operator D taking functions $f \mapsto Df$ is linear.)

Proof. Suppose that there are two derivatives at x , $A, B \in B(X, Y)$. Define the two remainders

$$\begin{aligned} R_A(x, h) &= f(x+h) - f(x) - Ah \\ R_B(x, h) &= f(x+h) - f(x) - Bh. \end{aligned} \quad (4)$$

By assumption, both $R_A(x, h) = o(\|h\|_X)$ and $R_B(x, h) = o(\|h\|_X)$. Now we show that in fact, $A = B$.

$$\begin{aligned} \|A - B\|_{B(X, Y)} &= \sup_{\|h\|_X = \varepsilon} \frac{\|Ah - Bh\|_Y}{\|h\|_X} \\ &= \sup_{\|h\|_X = \varepsilon} \frac{\|Ah - Bh\|_Y}{\|h\|_X} \\ &= \sup_{\|h\|_X = \varepsilon} \frac{\|R_A(x, h) - R_B(x, h)\|_Y}{\|h\|_X} \\ &\leq \sup_{\|h\|_X = \varepsilon} \frac{\|R_A(x, h)\|_Y}{\|h\|_X} + \sup_{\|h\|_X = \varepsilon} \frac{\|R_B(x, h)\|_Y}{\|h\|_X}. \end{aligned} \quad (5)$$

Is this the right definition of the norm on $B(X, Y)$? The RHS can be made small as $\varepsilon \rightarrow 0$. Indeed, the argument of the supremum goes to zero as $\|h\| \rightarrow 0$ by the $o(\|h\|_X)$ assumption. Thus, $A = B = Df(x)$ is unique. To prove that f is continuous at x , we have

$$\begin{aligned} \|f(x+h) - f(x)\|_Y &= \|Df(x)h + R(x, h)\|_Y \\ &\leq \|Df(x)h\|_Y + \|R(x, h)\|_Y \\ &\leq \|Df(x)\|_{B(X, Y)}\|h\|_X + \|R(x, h)\|_Y. \end{aligned} \quad (6)$$

The RHS tends to 0 as $h \rightarrow 0$ in X . Since $Df(x)$ is bounded, the first term on the right tends to zero trivially as $h \rightarrow 0$. Since $R(x, h)$ is $o(\|h\|)$, it means that $R(x, h)$ goes to zero faster than h goes to zero. This implies, however, that if $h \rightarrow 0$, then also $R(x, h) \rightarrow 0$. A somewhat obvious fact. \square

Having differentiability, namely *strong* differentiability, at a point x is more than enough to conclude that f is continuous at x . A piecewise linear function may be continuous, but not differentiable at the interface points. Strong differentiability allows us to conclude more:

Lemma 1.1 (Local Lipschitz Property). *If $f : U \rightarrow Y$ is differentiable at $x \in U$, then for every $\varepsilon > 0$, there exists a $\delta = \delta(x, \varepsilon) > 0$ such that for every h with $\|h\|_X \leq \delta$,*

$$\|f(x+h) - f(x)\|_Y \leq \left(\|Df(x)\|_{B(X,Y)} + \varepsilon \right) \|h\|_X.$$

Proof. We have

$$f(x+h) - f(x) = Df(x)h + R(x, h). \quad (7)$$

Since the remainder is $o(\|h\|)$, then for every $\varepsilon > 0$, there exists a $\delta > 0$ such that

$$\frac{\|R(x, h)\|_Y}{\|h\|_X} < \varepsilon \quad (8)$$

whenever $\|h\|_X < \delta$. Definition of convergence of $\|R\|/\|h\| \rightarrow 0$. It follows then that

$$\|f(x+h) - f(x)\|_Y = \|Df(x)h + R(x, h)\| \leq \|Df(x)\|_{B(X,Y)}\|h\|_X + \varepsilon\|h\|_X. \quad (9)$$

□

Even if f is not strongly differentiable, it may still admit some kind of weaker derivative. The key here is that in the Fréchet derivative, we require the remainder goes to zero for all paths that take $\|h\|_X \rightarrow 0$. We now consider incrementing only in a specific direction h . Start with $f : U \rightarrow Y$, fix a direction $h \in X$, and consider the auxillary function

$$g : \mathbb{R} \ni t \rightarrow g(t) := f(x + th) \in Y. \quad (10)$$

Proposition 1.2.

$$f \text{ is Fréchet differentiable at } x \implies f \text{ is Gateaux differentiable at } x.$$

Theorem 1.2 (Chain Rule). *Let X, Y, Z be NLS's and $U \subset X, V \subset Y$ open, $f : U \rightarrow Y$ and $g : V \rightarrow Z$. Let $x \in U$ and $y = f(x) \in V$. Suppose that g is Fréchet differentiable at y and f is Gateaux (or Fréchet) differentiable at x . Then the composition $g \circ f$ is Gateaux (or Fréchet) differentiable at x with*

$$D(g \circ f)(x) = Dg(y) \circ Df(x)$$

Proposition 1.3 (Mean Value Theorem for Curves). Let Y be an NLS and $\varphi : [a, b] \rightarrow Y$ be continuous, where $a < b$ are real numbers. Suppose $\varphi'(t)$ exists on (a, b) and $\|\varphi'(t)\|_{B(\mathbb{R}, Y)} \leq M$. Then

$$\|\varphi(b) - \varphi(a)\|_Y \leq M(b - a).$$

Proof. Use a compactness argument. □

Theorem 1.3 (Mean Value Theorem). Let X, Y be NLS's, $U \subset X$ open. Let $f : U \rightarrow Y$ be Fréchet differentiable at all $x \in U$, and suppose that the line segment

$$\ell = \{tx_2 + (1 - t)x_1 : t \in [0, 1]\}$$

is contained in U . Then

$$\|f(x_2) - f(x_1)\|_Y \leq \sup_{x \in \ell} \|Df(x)\|_{B(X, Y)} \|x_2 - x_1\|_X.$$

Proof. Use the MVT for curves applied to the function $\varphi : [0, 1] \rightarrow Y$,

$$\varphi(t) = f((1 - t)x_1 + tx_2) = f(x_1 + t(x_2 - x_1)) = f(\gamma(t)).$$

□

Definition 1.3. Let $X = X_1 \oplus \cdots \oplus X_m$. Let $U \subset X$ open, $F : U \rightarrow Y$, Y NLS. Let $x = (x_1, \dots, x_m) \in U$ and fix an integer $k \in [1, m]$. For z near x_k (k -th component) in space X_k , the point $(x_1, \dots, x_{k-1}, z, x_{k+1}, \dots, x_m) \in U$ since U is open. Define now

$$f_k(z) = F(x_1, \dots, x_{k-1}, z, x_{k+1}, \dots, x_m).$$

Then f_k maps an open set in X_k to Y . If f_k has a Fréchet derivative at $z = x_k$, then we say that the original F has a k -th partial derivative at x and define $D_k F(x) := Df_k(x_k)$. Notice that $D_k F(x) = Df_k(x_k) \in B(X_k, Y)$.

Proposition 1.4. Let $X = X_1 \oplus \cdots \oplus X_m$ be the direct sum of NLS's, $U \subset X$ open, and $F : U \rightarrow Y$, Y an NLS. Suppose that all partial derivatives $D_j F(x)$ exist for $x \in U$ and $j \in [1, m]$, and that these maps are continuous as a function of x at some $x_0 \in U$. Then, F is Fréchet differentiable at x_0 and for an increment $h = (h_1, \dots, h_m) \in X$,

$$DF(x_0)h = \sum_{j=1}^m D_j F(x_0)h_j.$$

1.2 Fixed Points and Contractive Maps

Definition 1.4 (Contraction Map, Fixed Point). Let (X, d) be a metric space, $G : X \rightarrow X$. We say that the mapping G is a contraction if and only if there exists a θ , $0 \leq \theta < 1$ such that

$$d(G(x), G(y)) \leq \theta d(x, y)$$

for every $x, y \in X$. We say that a point $x \in X$ is a fixed point of the mapping G if and only if $x = G(x)$.

The definition of a contraction immediately implies that every contraction is Lipschitz continuous with Lipschitz constant strictly smaller than 1.

Theorem 1.4 (Banach Contraction Mapping Theorem). Let (X, d) be a complete metric space and G a contraction on X . Then there is a unique fixed point of G in X .

Proof. Uniqueness: go by contradiction, then you would get $x = y$. Existence: Fix $x_0 \in X$. Consider the sequence $x_n = G(x_{n-1})$. Claim: x_n is a Cauchy sequence. Observe that

$$d(x_1, x_2) = d(G(x_0), G(x_1)) \leq \theta d(x_0, x_1)$$

and proceed by induction. Then you find out

$$d(x_n, x_{n+1}) \leq \theta^n d(x_0, x_1).$$

Then for $n \geq 0, m > n$,

$$\begin{aligned} d(x_n, x_m) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + \cdots + d(x_{m-1}, x_m) \\ &\leq (\theta^n + \cdots + \theta^{m-1}) d(x_0, x_1) \\ &= \theta^n (1 + \theta + \cdots + \theta^{m-n-1}) d(x_0, x_1) \\ &\leq \frac{\theta^n}{1 - \theta} d(x_0, x_1). \end{aligned}$$

Then since X is complete, the Cauchy sequence x_n converges to the limit, the fixed point. \square

Corollary 1.4.1 (Fixed Point Iteration). Let the assumptions of Theorem 1.4 hold. Suppose that G has a contraction constant θ , and let $x_0 \in X$. Then, if the sequence x_n is defined successively by $x_{n+1} = G(x_n)$, then $x_n \rightarrow x$, where x is the unique fixed point of G in X . Moreover,

$$d(x_n, x) \leq \frac{\theta^n}{1 - \theta} d(x_0, x_1).$$

Usually, the metric space X will be the space of continuous functions. For nonlinear equations, you may have to restrict yourself to a ball, $B_r(0) \subset X = C^0([0, T])$, or something similar. Once you take the closed ball, you have still a metric space, so everything else still works.

Corollary 1.4.2. *Let X be a Banach space and $f : X \rightarrow X$ a differentiable mapping. Suppose that $\|Df(x)\|_{B(X,X)} \leq \kappa < 1$ for $x \in \overline{B_R(0)}$. If there is an $x_0 \in B_R(0)$ such that $\overline{B_r(x_0)} \subset B_R(0)$ for some $r \geq \|f(x_0) - x_0\|/(1 - \kappa)$, then f has a unique fixed point in $B_r(x_0)$.*

That is to say, a map f which is *locally contractive*, and for which we can find a point that is not too far moved by f , has a fixed point which may be obtained by the sequence $x_{n+1} = f(x_n)$.

1.3 Nonlinear Equations

We have seen that the basic tool for proving the existence and uniqueness of nonlinear equations is the contraction mapping theorem. We consider now the more general problem of solving $f(x) = y$, where $f : X \rightarrow Y$ and X, Y are Banach spaces. Solving $f(x) = y$ is equivalent to solving for the fixed point of the operator

$$G(x) = x - T_x(f(x) - y).$$

where $T_x : Y \rightarrow X$ vanishes only at 0. $x = G(x) \implies 0 = T_x(f(x) - y) \implies f(x) = y$. If T is independent of x , then we also have

$$DG(x) = I - T \circ Df(x).$$

If $\|DG(x)\| \leq \kappa < 1$, that is, if we can select such a T to make the operator norm less than one, then we can apply the result of Corollary 1.4.2 in an attempt to find a fixed point. The result is

Theorem 1.5 (Simplified Newton Method). *Let X, Y Banach, $f : X \rightarrow Y$ a differentiable mapping. Suppose that $A = Df(x_0)$ has a bounded inverse, and that*

$$\|I - A^{-1}Df(x)\| \leq \kappa < 1$$

for every $x \in B_r(x_0)$ for some $r > 0$. Let

$$\delta = \frac{(1 - \kappa)r}{\|A^{-1}\|_{B(Y,x)}}.$$

Then the equation $f(x) = y$ has a unique solution $x \in B_r(x_0)$ whenever $y \in B_\delta(f(x_0))$.

Proof. Let $y \in B_\delta(f(x_0))$ be given, define a mapping

$$g_y : X \rightarrow X, \quad x \mapsto x - A^{-1}(f(x) - y).$$

Note that $f(x) = y$ if and only if $g_y(x) = x$. Same argument as previous. If A is invertible, then A^{-1} vanishes only at zero. Notice also that since A^{-1} is independent of x , we have

$$Dg_y(x) = I - A^{-1}Df(x).$$

By assumption, we have that $\|Dg_y(x)\| = \|I - A^{-1}Df(x)\| \leq \kappa$ for $x \in B_r(x_0)$, so g_y is a contraction on $\overline{B_r(x_0)}$. What remains to show is that g_y takes $B_r(x_0)$ into itself. But with our constructions of y, δ , **revisit this...f(x0) not f(x)!**

$$\begin{aligned} \|g_y(x) - x_0\| &\leq \|g_y(x) - g_y(x_0)\| + \|g_y(x_0) - x_0\| \\ &\leq \dots \\ &\leq r. \end{aligned}$$

The contraction map theorem now implies the existence of a unique solution in the ball. \square

Theorem 1.6 (Newton-Kantorovich Method). *Let X, Y be Banach spaces, $f : X \rightarrow Y$ a differentiable mapping. Assume that there is an $x_0 \in X$ and an $r > 0$ such that*

1. $A = Df(x_0)$ has a bounded inverse,

2. $\|Df(x_1) - Df(x_2)\| \leq \kappa\|x_1 - x_2\|$

for every $x_1, x_2 \in B_r(x_0)$. Let $y \in Y$, and set

$$\varepsilon = \|A^{-1}(f(x_0) - y)\|_X.$$

For any y such that

$$\varepsilon \leq r/2, \quad 4\varepsilon\kappa\|A^{-1}\|_{B(Y,X)} \leq 1,$$

the equation $f(x) = y$ has a unique solution in $B_r(x_0)$. Moreover, the solution is the limit of the iterations

$$x_{k+1} = x_k - Df(x_k)^{-1}(f(x_k) - y).$$

The convergence is asymptotically quadratic, that is

$$\|x_{k+1} - x_k\| \leq C\|x_k - x_{k-1}\|^2$$

for large k .

Theorem 1.7 (Inverse Function Theorem I). *Let the hypotheses of the Simplified Newton Method hold. Then the inverse mapping $f^{-1} : B_\delta(f(x_0)) \rightarrow B_r(x_0)$ is Lipschitz.*

Proof. Consider two points y_1, y_2 such that $f(x_i) = y_i$. Then estimate

$$\begin{aligned} \|f^{-1}(y_1) - f^{-1}(y_2)\| &= \|x_1 - x_2\| \\ &= \|g_{y_1}(x_1) - g_{y_1}(x_2) + A^{-1}(f(x_2) - f(x_1))\| \\ &\leq \kappa\|x_1 - x_2\| + \|A^{-1}\|\|y_2 - y_1\|. \end{aligned}$$

So, f^{-1} is Lipschitz with at worst constant $\|A^{-1}\|/(1 - \kappa)$. \square

Definition 1.5 (Diffeomorphism). Let X, Y be Banach spaces, $U \subset X$ open, $V \subset Y$ open. Let $f : U \rightarrow V$ be 1-1 and onto. We say that f is a diffeomorphism on U (equivalently, that U is diffeomorphic to V) if and only if both f and f^{-1} are C^1 . That is, the maps

$$x \mapsto Df(x) \quad y \mapsto Df^{-1}(y)$$

are continuous from U to $B(X, Y)$ and from V into $B(Y, X)$ respectively.

Theorem 1.8 (Inverse Function Theorem II). Let X, Y be Banach. Let $x_0 \in X$ be such that f is C^1 in a neighborhood of x_0 and $Df(x_0)$ is an isomorphism. Then there is an open set $U \subset X$, $x_0 \in U$, and an open set $V \subset Y$, $f(x_0) \in V$ such that $f : U \rightarrow V$ is a diffeomorphism. Moreover, for $y \in V$, $x \in U$, $y = f(x)$, we have

$$D(f^{-1})(y) = (Df(x))^{-1}.$$

Lemma 1.9. Let X, Y Banach. Then $GL(X, Y)$ is an open subset of $B(X, Y)$. If $GL(X, Y)$ is nonempty, then the mapping $J_{X,Y} : GL(X, Y) \ni A \rightarrow A^{-1} \in GL(Y, X)$ is 1-1, onto, and continuous.

Theorem 1.10 (Implicit Function Theorem). Let X, Y, Z be Banach spaces. Suppose that

- $f : Z \times X \rightarrow Y$ is a C^1 mapping defined at least in the neighborhood of a point (z_0, x_0) .
- Let $y_0 = f(z_0, x_0)$.
- $D_x f(z_0, x_0) \in GL(X, Y)$.

Then there exist open sets

1. $z_0 \in W \subset Z$,
2. $x_0 \in U \subset X$,
3. $y_0 \in V \subset Y$

and a unique mapping $g : W \times V \rightarrow U$ such that $f(z, g(z, y)) = y$ for all $(z, y) \in W \times V$. Moreover, g is C^1 and with $x = g(z, y)$,

$$Dg(z, y)(\eta, \zeta) = D_x f(z, x)^{-1} (\zeta - D_z f(z, x)\eta)$$

for all $(z, y) \in W \times V$ and $(\eta, \zeta) \in Z \times Y$.

If $Z = \{0\}$ is the trivial Banach space, then we recover the Inverse Function Theorem.

Corollary 1.10.1. *Let f be as in the Implicit Function Theorem. Then, there is a unique C^1 branch of solutions to the equation $f(z, x) = y_0$ in a neighborhood of (z_0, x_0) .*

1.4 Higher Derivatives

1.5 Extrema

1.6 Euler-Lagrange Equations

1.7 Constrained Extrema and Lagrange Multipliers