

# Some Notes and Theorems, Part 2

Jonathan Zhang

- 1 Preliminaries
- 2 Normed Linear Spaces and Banach Spaces
- 3 Hilbert Spaces
- 4 Spectral Theory and Compact Operators
- 5 Distributions

**Theorem 5.1.** *Here, we use the following notation:  $(M^\alpha f)(x) = x^\alpha f(x)$ .  
If  $T \in S'$  and  $f \in S$ , then  $T * f$  is smooth and polynomially bounded.*

## 6 The Fourier Transform

We motivate our discussion on the Fourier Transform by observing that it converts the task of solving a PDE into the task of solving an ODE, and that we can represent functions in terms of harmonic functions.

The Fourier Transform can be defined naturally for functions in  $L^1(\mathbb{R}^d)$ , where we will start. Then, we can extend it to functions in  $L^2(\mathbb{R}^d)$ , and we shall see that it maps  $L^2$  onto itself. Then, we will discuss the Fourier transform in context of distributions.

### 6.1 $L^1(\mathbb{R}^d)$ Theory

For (a point)  $\xi \in \mathbb{R}^d$ , the function

$$\varphi_\xi : x \mapsto e^{-ix \cdot \xi} = \cos(x \cdot \xi) - i \sin(x \cdot \xi)$$

for  $x \in \mathbb{R}^d$  defines a wave in the direction  $\xi$ . Moreover, the period of  $\varphi_\xi$  in the  $j$ -th direction is  $2\pi/\xi_j$ . Consider the simplest nontrivial example, say when we are in  $\mathbb{R}^2$  and we let  $\xi = (1, 1)$ . Then, the function  $\varphi_\xi(x) = e^{-i(x+y)} = \cos(x+y) - i \sin(x+y)$  defines a wave that propagates in the direction  $(1, 1)$ . Look at the real part by itself, for example. It is clear that the period in the  $x$  direction is  $2\pi$ , indeed as it should be. The same can be easily seen with the imaginary part. These harmonic functions have certain nice properties:

**Proposition 6.1.** 1.  $|\varphi_\xi| = 1$ , and  $\overline{\varphi_\xi} = \varphi_{-\xi}$  for every  $\xi \in \mathbb{R}^d$ ,

2.  $\varphi_\xi(x+y) = \varphi_\xi(x) \varphi_\xi(y)$  for every  $x, y, \xi \in \mathbb{R}^d$ ,

3.  $-\nabla^2 \varphi_\xi = |\xi|^2 \varphi_\xi$  for every  $\xi \in \mathbb{R}^d$ .

*Proof.* 1. We clearly have

$$|\varphi_\xi|^2 = \cos^2(x \cdot \xi) + \sin^2(x \cdot \xi) = 1,$$

and

$$\overline{\varphi_\xi} = \cos(x \cdot \xi) + i \sin(x \cdot \xi) = \cos(-x \cdot \xi) - i \sin(-x \cdot \xi) = \varphi_{-\xi}.$$

2.

$$\varphi_\xi(x+y) = e^{-i(x+y) \cdot \xi} = e^{-ix \cdot \xi - iy \cdot \xi} = e^{-ix \cdot \xi} e^{-iy \cdot \xi} = \varphi_\xi(x) \varphi_\xi(y).$$

3.

$$-\nabla^2 \varphi_\xi = -(-i\xi)^2 e^{-ix \cdot \xi} = |\xi|^2 \varphi_\xi.$$

We remark that this property demonstrates that functions  $\varphi_\xi$  are eigenfunctions of the Laplacian operator with eigenvalues  $-|\xi|^2$ .

□

We now define the Fourier Transform:

**Definition 6.1.** For  $f \in L^1(\mathbb{R}^d)$ , the Fourier Transform of  $f$  is

$$(\mathcal{F}f)(\xi) = \hat{f}(\xi) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} f(x) e^{-ix \cdot \xi} dx.$$

At this moment, it is not immediately clear what function spaces the Fourier Transform connects. However, for  $f \in L^1$ , the Fourier transform is well defined since

$$|f(x)e^{-ix \cdot \xi}| = |f(x)|.$$

What we mean when we say well defined is that the integral defining  $\hat{f}$  converges whenever  $f \in L^1$ . Indeed,  $f \in L^1$  iff  $\int |f(x)| dx < \infty$ , but as we just showed,  $\int |f(x)| dx = \int |f(x)e^{-ix \cdot \xi}| dx$ , so the integral converges.

Additionally, we remark that it is possible (and equivalent) to define the Fourier transform in different ways, but that the end results are very similar and it is merely a matter of preference.

We preface some additional discussion with a motivating example to show that we need to take care when talking about function spaces:

**Example.** On  $\mathbb{R}$ , let  $f : x \mapsto \chi_{[-1/2, 1/2]}(x)$ , the characteristic function on the interval  $[-1/2, 1/2]$ . Obviously,  $f$  is an  $L^1(\mathbb{R})$  function. Now for every  $\xi \neq 0$ ,

$$\begin{aligned} \hat{f}(\xi) &= (2\pi)^{-1/2} \int_{\mathbb{R}} f(x) e^{-ix \cdot \xi} dx \\ &= (2\pi)^{-1/2} \int_{-1/2}^{1/2} e^{-ix \cdot \xi} dx \\ &= (2\pi)^{-1/2} \frac{2}{k} \sin\left(\frac{k}{2}\right). \end{aligned}$$

This is fine, but notice that  $\hat{f}$  is no longer integrable! So, just because  $f \in L^1$ , we have no reason to believe that  $\hat{f}$  is also in  $L^1$ .

We have now the following proposition:

**Proposition 6.2.** The Fourier Transform viewed in the spaces

$$\mathcal{F} : L^1(\mathbb{R}^d) \rightarrow L^\infty(\mathbb{R}^d)$$

is a bounded linear map, and

$$\|\hat{f}\|_{L^\infty(\mathbb{R}^d)} \leq (2\pi)^{-d/2} \|f\|_{L^1(\mathbb{R}^d)}.$$

*Proof.* Clearly,  $\mathcal{F}$  is linear since the integral is a linear operator. Furthermore,

$$\|\hat{f}\|_{L^\infty} = \max_{\xi \in \mathbb{R}^d} (2\pi)^{-d/2} \left| \int_{\mathbb{R}^d} f(x) e^{-ix \cdot \xi} dx \right| \leq (2\pi)^{-d/2} \int_{\mathbb{R}^d} |f(x) e^{-ix \cdot \xi}| dx = (2\pi)^{-d/2} \|f\|_{L^1}$$

□

**Example.** Consider the characteristic function on the interval  $[-1, 1]^d$ ,

$$f : x \mapsto \chi_{[-1, 1]^d}(x)$$

Certainly,  $f$  is an  $L^1$  function. Its Fourier Transform is

$$\begin{aligned} \hat{f}(\xi) &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(x) e^{-ix \cdot \xi} dx \\ &= (2\pi)^{-d/2} \int_{-1}^1 \int_{-1}^1 \dots \int_{-1}^1 e^{-ix \cdot \xi} dx_1 dx_2 \dots dx_d \\ &= (2\pi)^{-d/2} \prod_{j=1}^d \int_{-1}^1 e^{-ix_j \xi_j} dx_j \\ &= (2\pi)^{-d/2} \prod_{j=1}^d \frac{2 \sin \xi_j}{\xi_j} \end{aligned}$$

We can easily verify that  $\hat{f}$  is bounded, since each term is bounded by  $2/\xi_j$  and so the finite product is bounded.

Additional properties include

**Proposition 6.3.** For  $f \in L^1(\mathbb{R}^d)$  and a translation operator  $\mathcal{T}_y$ , i.e.  $\mathcal{T}_y f(x) = f(x - y)$ ,

1.  $\widehat{(\mathcal{T}_y f)}(\xi) = e^{-iy \cdot \xi} \hat{f}(\xi)$  for every  $y \in \mathbb{R}^d$ , that is, the Fourier transform takes translations into multiplication.
2.  $\widehat{(e^{ix \cdot y} f)}(\xi) = \mathcal{T}_y \hat{f}(\xi)$  for every  $y \in \mathbb{R}^d$ , that is, the Fourier transform takes multiplication by a harmonic function to translation.
3. For any  $r > 0$ ,  $\widehat{f(rx)}(\xi) = r^{-d} \hat{f}(r^{-1}\xi)$ , that is, the Fourier transform takes scaling into premultiplying and scaling.
4.  $\widehat{\bar{f}}(\xi) = \overline{\hat{f}(-\xi)}$ , that is, the Fourier transform of complex conjugate equals complex conjugate of the Fourier transform, reflected.

*Proof.* 1.

$$\begin{aligned}
 \mathcal{F}f(x-y)(\xi) &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(x-y) e^{-ix \cdot \xi} dx \\
 &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(u) e^{-i(u+y) \cdot \xi} du \\
 &= e^{-iy \cdot \xi} \mathcal{F}f(x)(\xi)
 \end{aligned}$$

2.

$$\begin{aligned}
 \mathcal{F}(e^{ix \cdot y} f(x))(\xi) &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} e^{ix \cdot y} f(x) e^{-ix \cdot \xi} dx \\
 &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(x) e^{-ix \cdot \xi + ix \cdot y} dx \\
 &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(x) e^{-ix \cdot (\xi - y)} dx \\
 &= \mathcal{F}f(x)(\xi - y) \\
 &= \mathcal{T}_y \hat{f}(\xi)
 \end{aligned}$$

3.

$$\begin{aligned}
 \mathcal{F}f(rx)(\xi) &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(rx) e^{-ix \cdot \xi} dx \\
 &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(u) e^{-i \frac{u}{r} \cdot \xi} \frac{du}{r^d} \\
 &= r^{-d} (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(u) e^{-i \frac{u}{r} \cdot \xi} du \\
 &= r^{-d} (2\pi)^{-d/2} \hat{f}(r^{-1} \xi)
 \end{aligned}$$

4.

$$\begin{aligned}
\mathcal{F}\bar{f}(\xi) &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} \bar{f}(x) e^{-ix \cdot \xi} dx \\
&= (2\pi)^{-d/2} \int_{\mathbb{R}^d} \overline{f(x) e^{-ix \cdot \xi}} dx \\
&= (2\pi)^{-d/2} \int_{\mathbb{R}^d} \overline{f(x) e^{-ix \cdot (-\xi)}} dx \\
&= (2\pi)^{-d/2} \overline{\int_{\mathbb{R}^d} f(x) e^{-ix \cdot (-\xi)} dx} \\
&= \widehat{\hat{f}}(-\xi)
\end{aligned}$$

Convince yourself that we can pull the complex conjugate out, or let  $f(x) = a(x) + ib(x)$ , and

$$\begin{aligned}
\mathcal{F}\bar{f}(\xi) &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} \bar{f}(x) e^{-ix \cdot \xi} dx \\
&= (2\pi)^{-d/2} \left[ \int_{\mathbb{R}^d} a(x) e^{-ix \cdot \xi} dx - i \int_{\mathbb{R}^d} b(x) e^{-ix \cdot \xi} dx \right] \\
&= (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(x) e^{-ix \cdot \xi} dx
\end{aligned}$$

by linearity of the integral.

□

While the Fourier transform maps  $L^1$  into  $L^\infty$ , it most definitely does not map onto the entire space  $L^\infty$ . Let us attempt to better understand the range of  $\mathcal{F}$ . We do know that it is contained fully in a subspace of  $L^\infty$  we denote  $C_v(\mathbb{R}^d)$ .

**Definition 6.2.** The space  $C_v(\mathbb{R}^d)$  is the space of continuous functions that vanish at infinity, i.e.

$$C_v(\mathbb{R}^d) = \{f \in C_0(\mathbb{R}^d) : f \text{ vanishes at } \infty\}.$$

We say that a continuous function  $f$  vanishes at infinity if and only if for every  $\varepsilon > 0$ , there exists a pre-compact  $K \subset \mathbb{R}^d$  such that

$$|f(x)| < \varepsilon$$

whenever  $x \notin K$ . Compare this definition with that of compact support. Notice that this is a slightly weaker definition, and certainly all continuous functions with compact support belong to  $C_v(\mathbb{R}^d)$ .

**Proposition 6.4.**  $C_v(\mathbb{R}^d)$  is a closed linear subspace of  $L^\infty(\mathbb{R}^d)$ .

*Proof.*

□

**Lemma 6.1 (Reimann-Lebesgue Lemma).** *The Fourier Transform maps*

$$\mathcal{F} : L^1(\mathbb{R}^d) \rightarrow C_v(\mathbb{R}^d).$$

*That is to say, the Fourier Transform of an  $L^1$  function is a  $C_v$  function. Thus, for  $f \in L^1(\mathbb{R}^d)$ ,*

$$\hat{f} \in C_0(\mathbb{R}^d) \quad \text{and} \quad \lim_{|\xi| \rightarrow \infty} |\hat{f}(\xi)| = 0.$$

*(Merely the definition of  $\hat{f} \in C_0$ )*

*Proof.* See Hans' homework problem. Suppose  $f \in L^1(\mathbb{R}^d)$ .

**Sub-Lemma 1.**

□

## 6.2 $\mathcal{S}(\mathbb{R}^d)$ Theory

## 6.3 $L^2(\mathbb{R}^d)$ Theory

## 6.4 $\mathcal{S}'$ Theory

## 6.5 Applications

## 7 Sobolev Spaces

### 7.1 Definitions, Basic Properties

### 7.2 Extensions from $\Omega$ to $\mathbb{R}^d$

### 7.3 Sobolev Imbedding Theorem

### 7.4 Compactness

### 7.5 The Spaces $H^s$

### 7.6 Trace Theorem

### 7.7 The Spaces $W^{s,p}(\Omega)$



## **8 Boundary Value Problems**

### **8.1 Second Order Elliptic PDEs**

### **8.2 A Variational Problem and Minimization of Energy**

### **8.3 Closed Range Theorem, Operators Bounded Below**

### **8.4 The Lax-Milgram Theorem**

### **8.5 Applications to Second Order Elliptic Equations**

### **8.6 Galerkin Approximations**

### **8.7 Green's Functions**

## 9 Differential Calculus in Banach Spaces

## 10 Calculus of Variations