

Homework set 4

shsp
Nov. 2023

Solution suggestion

a) V is a real vector space $\mathbb{R}^{3 \times 3}$. A 6-dimensional subspace of V can be spanned by 6 basis vectors, so 6 linearly independent elements from V . We note that $\dim(V) = 9 < 6$, so a 6-dimensional subspace of V exists according to Lemma 9.31.

We can choose the 6 linearly independent vectors from V :

$$W = \text{Span}\left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}\right)$$

which span a subset from V . According to the remark after Definition 9.36, a spanning is a subspace, hence W is a 6-dimensional subspace within V .

b) In \mathbb{R}^3 we are given a vector space W as:

$$W = \text{Span}\left(\underbrace{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}_{v_1}, \underbrace{\begin{bmatrix} 5 \\ 4 \\ 3 \end{bmatrix}}_{v_2}, \underbrace{\begin{bmatrix} 7 \\ 11 \\ 6 \end{bmatrix}}_{v_3}\right).$$

A basis for W must according to Definition 9.14 span W and its vectors must be linearly independent.

The spanning is given, and thus only lin. independence must be investigated. For this, the vectors are merged as columns in a matrix:

$$\begin{bmatrix} v_1 & v_2 & v_3 \end{bmatrix} = \begin{bmatrix} 1 & 5 & 7 \\ -1 & 4 & 11 \\ 0 & 3 & 6 \end{bmatrix} \xrightarrow{R_2: R_2 + R_1} \begin{bmatrix} 1 & 5 & 7 \\ 0 & 9 & 18 \\ 0 & 3 & 6 \end{bmatrix} \xrightarrow{R_2: \frac{1}{9}R_2} \begin{bmatrix} 1 & 5 & 7 \\ 0 & 1 & 2 \\ 0 & 3 & 6 \end{bmatrix} \xrightarrow{R_3: R_3 - 3R_2} \begin{bmatrix} 1 & 5 & 7 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\xrightarrow{R_1: R_1 - 5R_2} \begin{bmatrix} 1 & 0 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} = \text{rref}\left(\begin{bmatrix} v_1 & v_2 & v_3 \end{bmatrix}\right)$$

The three vectors are according to Theorem 7.8 not linearly independent, since the rank is not 3. Without v_3 a matrix $[v_1 \ v_2]$ would have rank 2, and v_1 and v_2

①

↓ continues...

... continued...

are thus linearly independent. A basis for W is hence $(v_1, v_2) = \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 5 \\ 3 \end{bmatrix} \right)$.

c)

A map $L: C_0 \rightarrow C_0$ between real vector spaces is given by:

$$L(f) = f' + f + 1.$$

L is linear, according to Definition 10.1, if $L(u + c \cdot v) = L(u) + c \cdot L(v)$ for any two functions $u, v \in C_0$ and any scalar $c \in \mathbb{R}$.

We check if that's the case:

$$\begin{aligned} L(u + c \cdot v) &= (u + c \cdot v)' + (u + c \cdot v) + 1 = u' + c \cdot v' + u + c \cdot v + 1 \\ &= u' + u + c(v' + v) + 1 \end{aligned}$$

$$L(u) + c \cdot L(v) = u' + u + 1 + c(v' + v + 1) + 1 = u' + u + c(v' + v) + c + 2$$

We see that $L(u + c \cdot v) \neq L(u) + c \cdot L(v)$ and L is thus not linear.

d) A map $L: \mathbb{C}^3 \rightarrow \mathbb{C}^2$ is given as:

$$L\left(\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix}\right) = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -4 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix}, \quad v_1, v_2, v_3 \in \mathbb{C}$$

Two ordered bases are given:

$$\beta = \left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right) \text{ for } \mathbb{C}^2 \text{ and } \gamma = \left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right) \text{ for } \mathbb{C}^3.$$

The shown matrix $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -4 & 5 \end{bmatrix}$ can be considered a mapping matrix of $L = L_A$ with respect to the standard basis e . Hence $A = {}_e[L]_e$. We now wish to change bases as follows, according to Theorem 10.33:

$${}_B[id]_e {}_e[L]_e [id]_\gamma = {}_B[L]_\gamma \quad \text{to a mapping matrix with respect to the } \gamma\text{- and } B\text{-bases, respectively.}$$

The change-of-basis matrices from the standard basis to B or γ basis, respectively, are needed for this.

according to Definition 10.30

②

↓ continues...

...continued...

We can from the given basis vectors create:

$${}_e[id]_\beta = \begin{bmatrix} 1 & -2 \\ 2 & 1 \end{bmatrix} \quad \text{and} \quad {}_e[id]_\gamma = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \end{bmatrix},$$

and ${}_\beta[id]_e = ({}_e[id]_\beta)^{-1}$ according to Lemma 10.37, bullet 3. Finding this inverse:

$$\begin{aligned} [{}_e[id]_\beta \mid \mathbb{I}_2] &= \begin{bmatrix} 1 & -2 & 1 & 0 \\ 2 & 1 & 0 & 1 \end{bmatrix} \xrightarrow{R_2: R_2 - 2R_1} \begin{bmatrix} 1 & -2 & 1 & 0 \\ 0 & 5 & -2 & 1 \end{bmatrix} \\ &\xrightarrow{R_2: \frac{1}{5}R_2} \begin{bmatrix} 1 & -2 & 1 & 0 \\ 0 & 1 & -\frac{2}{5} & \frac{1}{5} \end{bmatrix} \xrightarrow{R_1: R_1 + 2R_2} \begin{bmatrix} 1 & 0 & -\frac{4}{5} & \frac{2}{5} \\ 0 & 1 & -\frac{2}{5} & \frac{1}{5} \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 & -\frac{4}{5} & \frac{2}{5} \\ 0 & 1 & -\frac{2}{5} & \frac{1}{5} \end{bmatrix} = \begin{bmatrix} 1 & 0 & \underbrace{-\frac{4}{5} \quad \frac{2}{5}}_{{}_\beta[id]_e} \\ 0 & 1 & -\frac{2}{5} \quad \frac{1}{5} \end{bmatrix} \end{aligned}$$

so, ${}_\beta[id]_e = \begin{bmatrix} \frac{1}{5} & \frac{2}{5} \\ -\frac{2}{5} & \frac{1}{5} \end{bmatrix}.$

Computing the new mapping matrix:

$$\begin{aligned} {}_\beta[L]_\gamma &= {}_\beta[id]_e {}_e[L]_e {}_e[id]_\gamma = \begin{bmatrix} \frac{1}{5} & \frac{2}{5} \\ -\frac{2}{5} & \frac{1}{5} \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 & 1 \\ 2 & -4 & 5 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & -\frac{7}{5} & \frac{11}{5} \\ 0 & -\frac{6}{5} & \frac{3}{5} \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} \frac{9}{5} & \frac{4}{5} & \frac{11}{5} \\ -\frac{3}{5} & -\frac{3}{5} & \frac{3}{5} \end{bmatrix} \\ &= \frac{1}{5} \begin{bmatrix} 9 & 4 & 11 \\ -3 & -3 & 3 \end{bmatrix} \end{aligned}$$

c) For a real $\mathbb{R}^{2 \times 2}$ vector space ^{of 2×2 matrices} with real coefficients V , a basis is chosen as: $\beta = ([\begin{smallmatrix} 0 & 0 \\ 0 & 0 \end{smallmatrix}], [\begin{smallmatrix} 0 & 0 \\ 0 & 1 \end{smallmatrix}], [\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}], [\begin{smallmatrix} 1 & 0 \\ 0 & 0 \end{smallmatrix}])$
A linear map $M: V \rightarrow V$ is given by

$$M(\underline{A}) = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \cdot \underline{A}, \quad \underline{A} \in V.$$

A mapping matrix ${}_\beta[M]_\beta$ will, according to Theorem 10.31, consist of images of the β -basis vectors as columns.

3

↓ continues...

We find the image by M of each β -basis vector:

$$M\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

$$M\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = 2 \cdot \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

$$M\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix} = 0 \cdot \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

$$M\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix} = 0 \cdot \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

Expressed in β -basis we have:

$${}_{\beta}[M]_{\beta} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 2 \end{bmatrix}$$

f) Given: $A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 0 & -1 \\ 2 & -1 & 0 \end{bmatrix} \in \mathbb{R}^{3 \times 3}$

Eigenvalues are found by solving the characteristic equation, according to Theorem 11.9:

$$P_A(\lambda) = \det([A - \lambda I_3]) = \det\left(\begin{bmatrix} 1-\lambda & 0 & 0 \\ 2 & -\lambda & -1 \\ 2 & -1 & -\lambda \end{bmatrix}\right) = 0$$

Finding the characteristic polynomial can be done using the expansion method for the determinant by choosing to expand along row $i=1$:

according to Theorem 8.20

$$P_A(\lambda) = \det\left(\begin{bmatrix} 1-\lambda & 0 & 0 \\ 2 & -\lambda & -1 \\ 2 & -1 & -\lambda \end{bmatrix}\right) = \sum_{j=1}^3 (-1)^{1+j} a_{1j} \det(A(1;j))$$

$$= (-1)^2 (1-\lambda) \det\left(\begin{bmatrix} -\lambda & -1 \\ -1 & -\lambda \end{bmatrix}\right) + (-1)^3 \cdot 0 \cdot \det(A(1;2)) + (-1)^4 \cdot 0 \cdot \det(A(1;3))$$

$$= (1-\lambda)((-\lambda)^2 - (-1)^2) = (1-\lambda)(\lambda^2 - 1) = \lambda^2 - 1 - \lambda^3 + \lambda = -\lambda^3 + \lambda^2 + \lambda - 1$$

Solving the characteristic equation:

$-\lambda^3 + \lambda^2 + \lambda - 1 = 0$ can be done by first guessing a root $\lambda_1 = 1$, which holds since $-1^3 + 1^2 + 1 - 1 = 0$

$$\begin{aligned} &\uparrow \\ &-1 + 1 + 1 - 1 = 0 \\ &\uparrow \\ &0 = 0 \end{aligned}$$

Then we perform polynomial division following Example 4.18:

$$\begin{array}{r|l} \lambda - 1 & -\lambda^3 + \lambda^2 + \lambda - 1 \\ & \underline{-\lambda^3 + \lambda^2} \\ & \lambda - 1 \\ & \underline{\lambda - 1} \\ & 0 \end{array} \quad Q(\lambda)$$

In factorized form we now have $P_A(\lambda) = (\lambda - 1)Q(\lambda) = (\lambda - 1)(-\lambda^2 + 1)$

↓ continues...

(4)

Finding roots of $Q(\lambda)$: $Q(\lambda) = 0 \Leftrightarrow -\lambda^2 + 1 = 0$

$$\Leftrightarrow \lambda^2 = 1 \Leftrightarrow \lambda = \pm 1$$

All roots, that is all eigenvalues of A , are thus: $\lambda_1 = 1$ and $\lambda_2 = -1$, with $gm(1) = 2$ and $gm(-1) = 1$.

Finding corresponding eigenspaces by solving $(A - \lambda_i I_3) \underline{v}_i = \underline{0}$ for each eigenvalue λ_i :

$$\text{For } \lambda_1 = 1: \left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 2 & -1 & -1 & 0 \\ 2 & -1 & -1 & 0 \end{array} \right] \xrightarrow{R_3 \leftrightarrow R_1} \left[\begin{array}{ccc|c} 2 & -1 & -1 & 0 \\ 2 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \xrightarrow{R_2: R_2 - R_1} \left[\begin{array}{ccc|c} 2 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\xrightarrow{R_1: \frac{1}{2} R_1} \left[\begin{array}{ccc|c} 1 & -\frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \Leftrightarrow \begin{cases} v_1 - \frac{1}{2} v_2 - \frac{1}{2} v_3 = 0 \\ v_1 = \frac{1}{2} v_2 + \frac{1}{2} v_3 = \frac{1}{2} t + \frac{1}{2} s \end{cases}$$

$$\Leftrightarrow \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ 1 \\ 0 \end{bmatrix} t + \begin{bmatrix} \frac{1}{2} \\ 0 \\ 1 \end{bmatrix} s, \quad t, s \in \mathbb{R}$$

$$\text{For } \lambda_2 = -1: \left[\begin{array}{ccc|c} 2 & 0 & 0 & 0 \\ 2 & 1 & -1 & 0 \\ 2 & -1 & 1 & 0 \end{array} \right] \xrightarrow{R_3: R_3 - R_1} \left[\begin{array}{ccc|c} 2 & 0 & 0 & 0 \\ 2 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \end{array} \right] \xrightarrow{R_2: R_2 - R_1} \left[\begin{array}{ccc|c} 2 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \end{array} \right]$$

$$\xrightarrow{R_1: \frac{1}{2} R_1} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \end{array} \right] \xrightarrow{R_3: R_3 + R_2} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \Leftrightarrow \begin{cases} v_1 = 0 \\ v_2 - v_3 = 0 \end{cases}$$

$$\Leftrightarrow \begin{cases} v_1 = 0 \\ v_2 = v_3 = t \end{cases}$$

$$\Leftrightarrow \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} t, \quad t \in \mathbb{R}$$

We have denoted the coordinates of eigenvectors by $\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix}$.
We see eigenspace E_1 having ^{ordered} basis $\left(\begin{bmatrix} \frac{1}{2} \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} \frac{1}{2} \\ 0 \\ 1 \end{bmatrix} \right)$ and $gm(1) = 2$
and eigenspace E_{-1} having (ordered) basis $\left(\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \right)$ and $gm(-1) = 1$

According to Corollary 11.28, A is diagonalizable since $am(1) = gm(1) = 2$ and $am(-1) = gm(-1) = 1$.

⑤ \downarrow continued...

∴ Hence a matrix V must exist such that

$$\underline{V}^{-1} \underline{A} \underline{V} = \underline{\Lambda} \quad \text{where } \underline{\Lambda} \text{ is a diagonal matrix.}$$

Following Example 11.31, such a diagonal matrix will consist of eigenvalues in the diagonal, and \underline{V} of corresponding linearly independent eigenvectors is corresponding columns, $\underline{V} = [\underline{v}_1 \ \underline{v}_2 \ \underline{v}_3]$, $\underline{\Lambda} = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix}$

Hence $\underline{\Lambda} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$. For $\lambda=1$ where $\dim(1)=2$ we choose two linearly independent eigenvectors, $\begin{bmatrix} 1/2 \\ 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 1/2 \\ 0 \\ 1 \end{bmatrix}$, and have $\underline{V} = \begin{bmatrix} 1/2 & 1/2 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}$.

The inverse \underline{V}^{-1} of \underline{V} is found:

$$\left[\underline{V} \mid \underline{I}_3 \right] = \left[\begin{array}{ccc|ccc} 1/2 & 1/2 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 1/2 & 1/2 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_1: R_1 - \frac{1}{2}R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1/2 & -1/2 & 1 & -1/2 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2: 2 \cdot R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 2 & -1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R_3: R_3 - R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 2 & -1 & 0 \\ 0 & 0 & 2 & -2 & 1 & 1 \end{array} \right] \xrightarrow{R_3: \frac{1}{2}R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 2 & -1 & 0 \\ 0 & 0 & 1 & -1 & 1/2 & 1/2 \end{array} \right]$$

$$\xrightarrow{R_2: R_2 + R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & -1/2 & 1/2 \\ 0 & 0 & 1 & -1 & 1/2 & 1/2 \end{array} \right] \xrightarrow{R_1: R_1 - R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1/2 & -1/2 \\ 0 & 1 & 0 & 1 & -1/2 & 1/2 \\ 0 & 0 & 1 & -1 & 1/2 & 1/2 \end{array} \right]$$

$$\text{So, } \underline{V}^{-1} = \begin{bmatrix} 1 & 1/2 & -1/2 \\ -1 & -1/2 & 1/2 \\ -1 & 1/2 & 1/2 \end{bmatrix}$$

$$\underline{V}^{-1}$$

In the equation $\underline{Q} \underline{A} \underline{Q}^{-1} = \underline{\Lambda}$ we see $\underline{Q} = \underline{V}^{-1}$.

(note, there are several correct \underline{Q} 's)