

應數一線性代數 2020 春, 期末考SOLUTION

學號: _____, 姓名: _____

本次考試共有 12 頁 (包含封面), 有 14 題。如有缺頁或漏題, 請立刻告知監考人員。

考試須知:

- 請在第一頁以及最後一頁填上姓名學號, 並在每一頁的最上方屬名, 避免釘書針斷裂後考卷遺失。
- 不可翻閱課本或筆記。
- 計算題請寫出計算過程, 閱卷人員會視情況給予部份分數。沒有計算過程, 就算回答正確答案也不會得到滿分。答卷請清楚乾淨, 儘可能標記或是框出最終答案。

高師大校訓: 誠敬弘遠

誠, 一生動念都是誠實端正的。敬, 就是對知識的認真尊重。宏, 開拓視界, 恢宏心胸。遠, 任重致遠, 不畏艱難。

請簽名保證以下答題都是由你自己作答的, 並沒有得到任何的外部幫助。

簽名: _____

1. (10 points) Find the projection matrix P for the plane $W : 3x + 2y + z = 0$ in \mathbb{R}^3 and the find the projection \vec{b}_w of $\vec{b} = [4, 2, -1]$ on it.

Answer: $\vec{b}_w = \frac{1}{14} \begin{bmatrix} 11 \\ -2 \\ -29 \end{bmatrix}$, $P = \frac{1}{14} \begin{bmatrix} 5 & -6 & -3 \\ -6 & 10 & -2 \\ -3 & -2 & 13 \end{bmatrix}$.

From 6-4

2. (10 points) Find the lease squares straight line fit to the four points (0,1) (1,3) (2,4) (3,4) and use it to approximate the fifth points (4, a).

Answer: the line equation = $1.5 + x$, a = 5.5 .

From 6-5

3. (5 points) Find the coordinate vector of $\begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix}$ in M_2 relative to $\left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & 3 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} \right)$

Answer: [3, 5, 1, 1]

From 7-1

4. (10 points) Find the five fifth roots of $-243i$. (need not simplify)

$$3 \left[\cos \left(\frac{3\pi}{10} + \frac{2k\pi}{5} \right) + i \sin \left(\frac{3\pi}{10} + \frac{2k\pi}{5} \right) \right], \quad k = 0, 1, 2, 3, 4$$

From 9-1

5. (10 points) Find the matrix representations $R_{B,B}$, $R_{B',B'}$ and an invertible C such that $R_{B',B'} = C^{-1}R_{B,B}C$ for the given linear transformation T .

$T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined as reflection of \mathbb{R}^3 through the plane $x+y-z=0$; $B = E$, $B' = ([1, 0, 1], [1, -1, 0], [1, 1, -1])$.

$$C_{BB'} = \frac{1}{3} \begin{bmatrix} 1 & 1 & 2 \\ 1 & -2 & -1 \\ 1 & 1 & -1 \end{bmatrix}, C_{BB'} = \begin{bmatrix} 1 & 11 & \\ 0 & -1 & 1 \\ 1 & 0 & -1 \end{bmatrix}, R_{B'B'} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \text{ and } R_{BB} = \frac{1}{3} \begin{bmatrix} 1 & -2 & 2 \\ -2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}.$$

Is $C = C_{BB'}$ or $C_{B'B}$? $C_{B'B}$.

From 7-2

6. (5 points) Express $(\sqrt{3} + i)^6$ in the form $a + bi$ for a, b are real numbers.

Answer: $a = -64$, $b = 0$.

From 9-1

7. (10 points) Using the Gram-Schmidt process to transform the basis $\{[1, i, 1 - i], [1 + i, 1 - i, 1]\}$ into an orthogonal basis and then extend it as an orthogonal basis for \mathbb{C}^3 .

Answer: the found orthogonal basis for \mathbb{C}^3 is $\{[1, i, 1 - i], [3 + 3i, 5 - 5i, 2], [-12i, 4, 8 + 8i]\}$

From 9-2

8. (10 points) Find an unitary matrix U and a diagonal matrix D such that $D = U^{-1}AU$. Also find where

$$A = \begin{bmatrix} 1 & -i & 0 \\ i & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

From 9-3

$$D = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}, U = \frac{1}{\sqrt{2}} \begin{bmatrix} i & -i & 0 \\ 1 & 1 & 0 \\ 0 & 0 & \sqrt{2} \end{bmatrix}$$

9. (10 points) Find a Jordan canonical form and a Jordan basis for the given matrix.

$$\begin{bmatrix} i & 0 & 0 & 0 & 0 \\ 0 & i & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \\ 2 & 0 & -1 & 0 & 2 \end{bmatrix}$$

From 9-4

$$J = \begin{bmatrix} \boxed{i} & & & & \\ & \boxed{i} & & & \\ & & 0 & & \\ & 0 & \boxed{\begin{matrix} 2 & 1 \\ 0 & 2 \end{matrix}} & & \\ & & & \boxed{2} & \end{bmatrix}, \text{ basis: } \{\vec{b}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \vec{b}_2 = \begin{bmatrix} 2-i \\ 0 \\ 0 \\ 0 \\ -2 \end{bmatrix}, \vec{b}_3 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \vec{b}_4 = \begin{bmatrix} 0 \\ 0 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \vec{b}_5 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}\}$$

$$J\vec{b}_1 = i\vec{b}_1,$$

$$J\vec{b}_2 = i\vec{b}_2,$$

$$J\vec{b}_3 = 2\vec{b}_3,$$

$$J\vec{b}_4 = 2\vec{b}_4 + \vec{b}_3,$$

$$J\vec{b}_5 = 2\vec{b}_5$$

10. (10 points) Find a polynomial in A that gives the zero matrix.

$$\begin{bmatrix} i & 0 & 0 & 0 & 0 \\ 0 & i & 1 & 0 & 0 \\ 0 & 0 & i & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 2 \end{bmatrix}$$

From 9-4

$$(A - iI)^2(A - 2I)^1$$

11. (5 points) Prove or disprove the following: All 2×2 matrix with determinant 1 is an orthogonal matrix.

From 6-3 #5

Let $A = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$, such that $\det(A) = 1$, but A is NOT orthogonal!

12. (10 points) Find all the possible 2×2 real matrix that is unitarily diagonalizable.

From 9-3 #17

Answer:

Every 2×2 real matrix A can be written as $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Since A is unitarily diagonalizable, A is normal, i.e. $A^*A = AA^*$.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^* \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix}^*$$

$$\begin{bmatrix} a\bar{a} + c\bar{c} & \bar{a}b + \bar{c}d \\ a\bar{b} + c\bar{d} & b\bar{b} + d\bar{d} \end{bmatrix} = \begin{bmatrix} a\bar{a} + b\bar{b} & a\bar{c} + b\bar{d} \\ \bar{a}c + \bar{b}d & c\bar{c} + d\bar{d} \end{bmatrix}$$

Hence: (notice that a, b, c, d are real.)

$$(1). \quad a\bar{a} + c\bar{c} = a\bar{a} + b\bar{b} \implies a^2 + c^2 = a^2 + b^2$$

$$(2). \quad \bar{a}b + \bar{c}d = a\bar{c} + b\bar{d} \implies ab + cd = ac + bd$$

$$(3). \quad a\bar{b} + c\bar{d} = \bar{a}c + \bar{b}d \implies ab + cd = ac + bd$$

$$(4). \quad b\bar{b} + d\bar{d} = c\bar{c} + d\bar{d} \implies b^2 + d^2 = c^2 + d^2$$

by (1) and (4), we have $b = c$ or $b = -c$. And (2)(3) holds for both cases.

13. (5 points) Prove that for $\vec{u}, \vec{v} \in \mathbb{C}^n$, $(\vec{u}^* \vec{v})^* = \overline{\vec{u}^* \vec{v}} = \vec{v}^* \vec{u} = \vec{u}^T \vec{v}$

From 9-2 #40

Answer:

$$\text{Let } \vec{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}, \vec{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$$

$$\begin{aligned} (\vec{u}^* \vec{v})^* &= \left(\begin{bmatrix} \overline{u_1} & \overline{u_2} & \dots & \overline{u_n} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} \right)^* = \left(\sum_{i=1}^n \overline{u_i} v_i \right)^* \\ &= \overline{\sum_{i=1}^n \overline{u_i} v_i} (= \overline{\vec{u}^* \vec{v}}) \\ &= \sum_{i=1}^n u_i \overline{v_i} \\ &= \begin{bmatrix} \overline{v_1} & \overline{v_2} & \dots & \overline{v_n} \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} (= \vec{v}^* \vec{u}) \\ &= \begin{bmatrix} u_1 & u_2 & \dots & u_n \end{bmatrix} \begin{bmatrix} \overline{v_1} \\ \overline{v_2} \\ \vdots \\ \overline{v_n} \end{bmatrix} (= \vec{u}^T \vec{v}) \end{aligned}$$

14. (10 points) Prove the following:
 - (a) Show that every Hermitian matrix is normal.
 - (b) Show that every unitary matrix is normal.
 - (c) Show that, if $A^* = -A$, then A is normal.

From 9-2 #43

Answer:

- (a) Let H are Hermitian matrices, i.e. $H^* = H$. $HH^* = HH = H^*H$.
 (b) Let U are unitary matrices, i.e. $U^*U = I$, i.e. $U^{-1} = U^*$. $UU^* = I + U^*U$.
 (c) If $A^* = -A$, $A * A = (-A)A = -AA = A(-A) = AA^*$.

應數一線性代數期末考SOLUTION, 學號: _____, 姓名: _____, 以下由閱卷人員填寫

[illegible]