

# Understandable Multifunctionality Measures Using Hill Numbers

Jarrett E. K. Byrnes<sup>1\*</sup>, Fabian Roger<sup>2\*</sup>, Robert Bagchi<sup>3</sup>

1 - Department of Biology, University of Massachusetts Boston, Boston, Massachusetts 02125 United States of America

2 - Department of Environmental Systems Science, Institute of Biogeochemistry and Pollutant Dynamics (IBP), ETH Zurich

3 - Department of Ecology and Evolutionary Biology, University of Connecticut, 75 N. Eagleville Road, Storrs, Connecticut, 06269-3043 United States of America.

\* - Shared first authorship

## Abstract

In ecology, multifunctionality measures the simultaneous provision of multiple ecosystem functions. If species diversity describes the variety of species that together build the ecosystem, multifunctionality attempts to describe the variety of functions these species perform. A range of methods have been proposed to quantify multifunctionality, successively attempting to alleviate problems that have been identified with the previous methods. This has led to a proliferation of more-or-less closely related metrics which, however, lack an overarching theoretical framework. Here we borrow from the comprehensive framework of species diversity to derive a new metric of multifunctionality. Analogously to the effective number of species used to quantify species diversity, the metric we propose is influenced both by the number of functions as well as, crucially, the evenness of performance levels across functions. In addition, the effective multifunctionality also considers the average level at which the functions are performed. The result is a measure of the cumulative performance of the system were all functions provided equally. The framework allows for the inclusion of the correlation structure among functions, thus allowing to account for non-independence between functions. We show that the average metric is a special case of the newly proposed metric when all functions are uncorrelated and performed at equal levels. We hope that by providing a new metric of multifunctionality anchored in the rigorous framework of species diversity based on effective numbers, we will overcome the considerable skepticism that the larger community of ecologists has built against indices of multifunctionality. We thereby hope to help popularize this important concept which, as biological diversity, describes a fundamental property of ecosystems and thus lies at the heart of ecology.

## Introduction

The past decade has witnessed the growth of the concept of ecosystem *multifunctionality*, where multifunctionality is defined as a measure of the simultaneous performance of multiple functions. It has moved from a concept in the biodiversity ecosystem function literature to community ecology writ large (Angelini et al. 2015), invasion biology (Ramus et al. 2017), land management (Nelson et al. 2009, Brandt et al. 2014, Binder et al. 2018), and more. The concept of multifunctionality is broad and can even be applied outside of community and ecosystem ecology - or even outside of ecology altogether. Its adoption as a unifying concept, however, has been slow - and for good reason.

Despite the popularity that the concept gained since it was first explicitly proposed over a decade ago (Hector and Bagchi 2007, Gamfeldt et al. 2008), there is still no consensus on how multifunctionality should be quantified. A range of methods have been proposed where each subsequent method attempted to alleviate problems the authors had identified with the previous method, leading to a proliferation of metrics which cannot easily be compared and can lead to inconsistent interpretations of the results (Byrnes et al. 2014a, Gamfeldt and Roger 2017). We do note that this is a difficult problem. Throwing many different ecologists against this problem in a stimulating working group over multiple years (we are thankful for NCEAS for providing this venue) has likely resulted in more ink impregnated on foreheads hitting whiteboards than should be typical, to say nothing of others who have attempted to cut this Gordian knot before and since (Hector and Bagchi 2007, Brandt et al. 2014, Dooley et al. 2015, Rodríguez-Loinaz et al. 2015, Stürck and Verburg 2017, Manning et al. 2018, Meyer et al. 2018, Hölting et al. 2019). In part, this problem is due to the lack of an underpinning body of theory describing how and where multifunctionality should arise. Regardless, this plurality of concepts has hampered a general understanding of multifunctionality and possibly its adoption outside of the subfield of Biodiversity and Ecosystem Function.

Species diversity, for a long time, suffered the same problems that multifunctionality suffers today: there are an uncountable number of metrics to estimate alpha diversity, reaching from the simple count of observed species without regard to their abundance (Richness), wide array of metrics incorporating the relative abundance (Shannon entropy, Simpson index or the Berger-Parker Dominance index to name some of the most common ones), to metrics based on the histograms of the abundance distribution (e.g. Fisher's alpha) (Magurran and McGill 2010). Yet, starting with MacArthur (1965), then Hill (1973), followed by Jost (2006), and most recently Chao et al. (2014a, 2019), a common framework for species diversity has been developed based on information theory. This framework, the effective number of species, encompasses the vast majority of previous metrics and is able to handle a wide variety of different issues in diversity estimation. We propose it can do the same for multifunctionality.

## **The current state of multifunctional affairs**

The definition of multifunctionality presents a challenge. How do we define a metric that captures both the level of performance of a broad suite of functions as well as the distribution of differences in performance among functions? Researchers have sought to capture this question in four ways, after standardizing functions to similar levels in order to prevent apples-to-oranges comparisons.

First, many have opted for simplicity and taking the average of all functions (e.g., Maestre et al. 2012). This approach is, at first glance, appealing, particularly as it provides a metric that can be put on a y-axis while a predictor is on the x-axis. However, it sacrifices crucial information about the system. An arithmetic mean tells us what level of functioning we would expect were we to sample any one function at random. Consider two plots - one where all functions are similar and performing at half their value and one where half of the functions are at their maximum while half are absent. The averaging approach says that they are identical. Geometric averaging (e.g., Hensel and Silliman 2013) appears to get around this problem to some degree, as the geometric mean is less than the arithmetic mean by a function of the variance of the observations (the two means are the same when all observations are the identical).

79 However, given its formulation, one critically low function can dominate the measurement - e.g., if even  
80 one function is 0, the geometric mean will be 0.

81 Second, we have metrics of the Multivariate Diversity Interactions framework (Dooley et al. 2015). This  
82 elegant framework allows us to tease apart the importance of correlations between functions and the  
83 contribution of different drivers to simultaneous change in those functions. It does not, however, provide  
84 a holistic metric of multifunctionality *per se*, much like the overlap approach before it (Hector and Bagchi  
85 2007). This approach from Hector and Bagchi is a generalization of Sørensen-Dice overlap (Dice 1945,  
86 Sørensen 1948). It provides key information on redundancy versus unique contributions of species. While  
87 it gives us rich information about a system, it lacks holistic interpretability.

88 Last, we have the threshold approach (Gamfeldt et al. 2008). This approach defines the “presence” of a  
89 function if it meets a given threshold of performance, and then sums up the number of functions  
90 performing at that threshold. There are clear issues with the arbitrariness of thresholds in the absence of  
91 relevant theory or management goals. To remedy this, the multiple threshold approach (Byrnes et al.  
92 2014a) seeks to balance the goals of measuring the simultaneous performance of multiple ecosystem  
93 functions with the arbitrariness of choosing a threshold of relevance for those functions. The results,  
94 however, can be difficult to interpret. Looking at multiple relationships between diversity and threshold-  
95 based multifunctionality does not provide a metric *per se*. While there are key metrics that can be  
96 extracted from looking at change in slope over different thresholds, the meaning of these quantities are  
97 non-obvious in their links to the concept of multifunctionality. The values can vary dramatically  
98 depending on the exact threshold chosen and on the method used to standardize the functions, even for  
99 simulated communities where all functions perform identically (Gamfeldt and Roger 2017). In the  
100 absence of suitable null-models, and even if the approach yields rich information about multifunctionality  
101 *sensu stricto*, it is unwieldy for most if not all who choose to use it, and we have noticed that those who  
102 do often report a single threshold in their main text anyway. Related, more recent efforts have sought to

103 use dimensionality-reducing techniques which have yielded metrics that, while useful, similarly lack  
104 easily interpretable meaning (Meyer et al. 2018).

## 105 **Hill numbers and effective diversity**

106 Over the last 70 years, ecologists studying how to measure species diversity have shown that the vast  
107 majority of previous diversity metrics can be organized into a common framework (Macarthur 1965, Hill  
108 1973, Jost 2006, Chao et al. 2014a, 2019). This is true for indices such as Shannon entropy, all Simpson  
109 measures, all Renyi entropies, all HCDT or "Tsallis" entropies and species richness (Jost, 2006) and  
110 more. All can be expressed as generalized entropies that can be converted to an effective number of  
111 species of "order"  $q$  which specifies the weighting of proportional abundances. The general formula for  
112 the diversity of order  $q$  for  $S$  species is the following:

$$113 \quad {}^qD = \left( \sum_{i=1}^S p_i^q \right)^{1/(1-q)} \quad (1)$$

114 Here,  $p_i$  is the relative abundance of the  $i$ th species and  $q$  is the weight given to the species' relative  
115 abundances. Species richness, the effective number of species based on Shannon entropy, the effective  
116 number of species based on the Simpson index, and the Berger-Parker dominance index are all effective  
117 numbers of species of order  $q = 0, 1, 2$  and  $\infty$ , respectively. (Note that the formula is undefined for  $q = 1$ ,  
118 but its limit  $q \rightarrow 1$  is  $\exp(-\sum p_i \log p_i)$ ). The effective number of species of order  $q$  is also often referred to  
119 as Hill numbers.

120

121 We propose leveraging this framework for a more meaningful and less *ad hoc* metric of  
122 multifunctionality composed of two parts: (1) the effective *number* of functions that are performed and  
123 (2) the *arithmetic mean performance* of the functions that are measured. Our proposed multifunctionality  
124 index is then the product of both terms. This approach draws on ideas already swirling in the  
125 multifunctionality literature (Brandt et al. 2014, Rodríguez-Loinaz et al. 2015, Stürck and Verburg 2017,

126 Hölting et al. 2019). With this approach, we aim to provide a unifying framework for the measurement of  
127 multifunctionality.

## 128 **Multifunctionality as the product of effective number of functions and average** 129 **level of functions**

130 To define the effective number of functions, we begin with a set of measurements on  $k$  functions that have  
131 been standardized to a common scale (i.e., between 0 and 1 where 0 means no function and 1 means  
132 maximum level of function). Let  $F_i, i \in 1, 2, \dots, K$  show the level of function for function  $i$ . The relative  
133 proportion a function contributes to the whole is defined as

$$134 \quad p_i = \frac{F_i}{\sum F_i} \quad (2)$$

135 We can now substitute the relative proportion into the formula for the effective number of types given in  
136 Eq 1

$$137 \quad {}^qN = \left( \sum_{i=1}^K p_i^q \right)^{1/(1-q)} \quad (3)$$

138 where  ${}^qN$  is the **effective number of functions** for some order  $q$ . *The effective number of functions here*  
139 *translates to the equivalent number of functions were all functions provided at the same level. Adapting*  
140 *from Jost (2006), if  $q = 0$ , this is the total number of functions  $K$ , which is unimportant as it is just the*  
141 *number of functions measured. For  $q = 1$ , the approximation of this function is equivalent to results from*  
142 *Shannon diversity for species shown earlier - which we note has been used as a multifunctionality metric*  
143 *previously (Stürck and Verburg 2017). For  $q > 1$ , functions performing at higher levels are given greater*  
144 *weight. At  $q = 2$ , we get results that are equivalent to the number of functions calculated from Simpson's*  
145 *diversity (inverse Simpson index). If one of our goals is to up-weight high performing functions,  $q=2$  is a*  
146 *reasonable choice, while  $q=1$  is sufficient as it accommodates information about unequal levels of*  
147 *functioning proportional to the relative functional performance. Lower values would upweight low-*  
148

performing functions - which might be desirable in certain contexts. The ability to modulate the sensitivity of the metric to high or low performing functions thus provides a strong tool for both ecologists and managers. In the absence of a justification for a particular value of  $q$ , exploring the robustness of results to different choices of  $q$  could prove fruitful as it has in biodiversity research. Further exploration of how  $q$  relates to management goals or ecological theory of multifunctionality would be a fruitful avenue for future research.

Effective number of functions does not accommodate knowledge about the absolute level of functioning in a system - a true metric of multifunctionality. As long as the distribution of  $p_i$  is the same, a system with high average function and low average function will look the same. To achieve this translation, we need to take into account the level at which the functions are performed: the arithmetic mean of the function values standardized to a common scale, which we define as  $A$ . As we are using standardized values as before,  $A$  will range from 0 to 1.

We can then calculate **effective multifunctionality** of order  $q$  as the product of both terms. This metric represents the cumulative performance of the system were it composed of functions all performing at equal levels

$${}^qM_{ef} = {}^qNA \quad (4)$$

This metric will have a maximum value of  $K$ , the total number of functions measured in the system, as maximum performance is all functions performing at a standardized level of 1. Alternatively, we can standardize by the total number of functions to calculate the fraction of performance achieved by the whole system, of **standardized effective multifunctionality**, which could facilitate future comparisons among studies.


$${}^qM = {}^qNA/K \quad (5)$$

However, care has to be taken when comparing multifunctionality values across systems that measure different sets of functions and the value of such comparisons is a topic that eludes consensus even among authors of this article.

Why must we consider the level of function and effective number of functions together in one metric?

First, when  $A$  is less than one, multiple values of  ${}^qN$  are possible depending on the distribution of performance of functions. Second, the average functioning and the effective number of functions are not independent of each other. The upper limit of  ${}^qN$  is  $K$  by definition.  ${}^qN$  is  $K$  when all functions are performing at the same level, i.e.  $E[A] = F_1, F_2, F_3 \dots F_K$ . This can be achieved for any value of  $A$ , low or high. Thus,  ${}^qN$  has no information about the level of function achieved. The lower limit of  ${}^qN$  (as long as  $q$  is not 0) occurs at the maximum level of unevenness of  $F$  across all functions. This lower limit will vary by different levels of  $A$ . For example,  ${}^qN$  will always be  $K$  when  $A$  is 1. Similarly, if all functions but one have a value of 0, then  ${}^qN$  will be 1, as  $p_i$  for the non-zero function will be 1. If more than 1 function is not 0, then the lower limit is set by a combination of  $A$  and the number of dominant functions, i.e., the functions performing at  $F_i = 1$  (Figure 1). See Appendix 1 for derivation.





**Figure 1.** Relationship between  ${}^qN$  and average level of functions for 4 functions at  $q = 1, 2, 3$ , and 4. Filled areas show the full range of possible values of effective number of functions. Note, areas with a higher order ( $q$ ) do overlap completely with lower orders, save for the additional edge that is visible.

Most importantly, a combined metric satisfies our definition of multifunctionality. High numbers imply both a high level of function and high functional evenness (i.e.,  $p_i$  is close to  $1/K$  for all functions). Low values imply that, even if a single function is being optimized, the assemblage of functions as a whole is not performing well. The relationship can also be easily decomposed into its constituent parts for a more detailed examination of its behavior.

Last, from a convenience standpoint, having a single metric allows us to begin to examine it as any other response variable. In the Biodiversity and Ecosystem Functioning world, we might look at additive partitioning in addition to complementary overlap approaches. In global change biology, we can look at stability, resistance, and resilience. The options are open.

## 204    **A note on standardization**

205    As pointed out in discussions of multifunctionality, how functions are standardized matters (Gamfeldt and  
206    Roger 2017, Manning et al. 2018). First is the choice of direction - what implies positive function? When  
207    services are being examined explicitly, this is hopefully a straightforward choice - although consider  
208    tradeoffs between nutrient cycling rates and storage as a tricky context. Further, not all functions are  
209    equally important, particularly in the case of management. Fortunately, choosing an ‘optimal’ level of  
210    function to link to 1 can alleviate this (e.g., if 25% of function is sufficiently high, 25% or higher can be  
211    considered a ‘1’). Functions can also be upweighted or downweighted in the calculation of  $\text{qN}$  so long as  
212     $\sum p_i = 1$ . Choices for standardization are often best made in the context of a specific system or application  
213    and must be transparently justified.

214

## 215    **Correlated functions and multifunctionality**

216    A great deal of debate in the multifunctionality literature has sprung from the issue of how to deal with  
217    correlations between functions (Bradford et al. 2014a, b, Byrnes et al. 2014b, Manning et al. 2018). As  
218    with species diversity, the rationale is that not all functions are equally different and that a metric of true  
219    multifunctionality should identify “variables that represent independent aspects of ecosystem functioning”  
220    (Manning et al. 2018). As an illustration, consider a scenario where we measure a set of functions, several  
221    of which result from a shared mechanism so that they are inextricably linked (for example, growth rate  
222    and final biomass). If we want to study the circumstances under which overall multifunctionality is  
223    maximized, the results will be disproportionately driven by circumstances maximizing this mechanism  
224    influencing the set of correlated functions; thus a naive application of a multifunctionality measure would  
225    implicitly upweight the importance of this function cluster over uncorrelated functions.

226

To address the issue of correlated functions, Manning et al. (2018) proposed performing a cluster analysis of ecosystem functions and building a dendrogram representing the distance matrix between functions. The performance of single functions is then averaged within clusters and multifunctionality is calculated among clusters and not individual functions. This is conceptually very similar to metrics of phylogenetic and functional species diversity that construct phylogenetic trees or dendrograms based on functional similarities and measure species diversity taking into account the dendrogram structure (e.g. Faith's PD, Allen's H, Petchey and Gaston's FD, Rao's Quadratic entropy etc.). The problem is that, as discussed by Chao et al. (2014a), dendrograms are very sensitive to the type of clustering method used and the shape of a dendrogram can vary substantially between different methods (Poos et al. 2009). For the method proposed by Manning et al 2018, it is also unclear at what depth the dendrogram should be cut (i.e., what clusters are considered as independent). Fortunately, extensions of the Hill numbers framework to account for species similarity (e.g., in terms of traits or genetic relatedness) ,(Chao et al. 2014a, 2019) provide a solution to the challenge of evaluating multifunctionality across correlated functions.

To account for correlation among functions in the effective multifunctionality framework, we need to incorporate several concepts from Chao et al. (2019). First, how similar are functions? For this, we need a distance matrix of some sort. If this is based on *a priori* mechanistic information about the underlying processes driving functions (e.g., physiological or evolutionary constraints), so much the better. We often do not have such a matrix. Other options include a matrix derived from principal components or other methods of constructing distance matrices (see Manning et al. 2018). In the absence of such information, a practical choice could be to look at the correlation matrix among functions,  $R$ . Defining  $D = (1 - R)/2$  would create a distance matrix where a  $d_{ij}$  value of 0 means two functions are inextricably linked (perfectly correlated) while 1 means they trade-off completely ( $r_{ij} = -1$ ). This is one choice, and we suggest that determining the proper way to create a distance matrix between functions for the estimation of effective diversity is an exciting area of research.

With a distance matrix in hand, we need to ask ourselves, how distinct must two functions be before we consider them completely different? This incorporates the threshold  $\tau$  proposed in Chao et al. (2019). Using this threshold, the distance matrix is truncated such that  $d_{ij}(\tau) = \min(d_{ij}, \tau)$ . If, for example,  $\tau = \min(d_{ij})$ , all functions are equally distinct and we recover the results ignoring correlations. If  $\tau = 1$  or the maximum distance in the matrix, we fully incorporate information from the distance matrix with no truncation. If  $\tau = 0.5$ , all negatively correlated functions are as distinct as uncorrelated functions. Chao et al. (2019) recommend using the average weighted pairwise distance between any two functions as  $\tau$

$$d_{mean} = \sum_{i=1}^K \sum_{j=1}^K d_{ij} p_i p_j \quad (6)$$

as setting  $\tau$  to  $d_{mean}$  yields results that are consistent with comprehensive evaluations of all possible values of  $\tau$ .

We can then use our truncated distance matrix,  $d_{ij}(\tau)$ , to define the effective number of underlying functions,  ${}^qN_u$  as follows when we are interested in the effective number of underlying “functions”. This equation takes advantage of  $d_{ij}(\tau)/\tau$  signifying the portion of one function that is distinct from another.

$${}^qN_u = \left( \sum_{i=1}^K p_i \left( \sum_{j=1}^K \left[ 1 - \frac{d_{ij}(\tau)}{\tau} \right] p_j \right)^{q-1} \right)^{1/(1-q)} \quad (7)$$

In essence, the inner part of the equation is the proportion of function  $i$  multiplied by the summed portion of all functions that contribute to the same underlying function. For  $q=1$ , we need to use the limit, as before, so

$${}^qN_u = \exp \left( - \sum_{i=1}^K p_i \log \left( \sum_{j=1}^K \left[ 1 - \frac{d_{ij}(\tau)}{\tau} \right] p_j \right) \right) \quad (8)$$

This effective number of functions can then be used to calculate multifunctionality as above. The average,  $A$ , does not need to be modified, because correlations do not systematically bias estimates of the mean (although uncertainty in that mean could be biased downwards by correlations). The correlations only need to be accounted for in the estimate of  ${}^qN$ .

## Discussion

With the framework above in hand, we are hopeful for a proliferation of literature investigating both new and old ideas in multifunctionality. With its ties to theoretically grounded methods of measuring species diversity and its flexibility to encompass some of the real sticking points in the field, we hope that Ecology will embrace this framework moving forward. It produces intuitive metrics - effective number of functions performing at equal levels, effective multifunction scaled by total level of performance, and standardized versions of both - as well as offers a solution to account for correlated functions. To aid in this advance, see Appendix 2 where we present a worked example in R (R Core Development Team 2008). Over the last decade and a half of its development, the multifunctionality literature has grown slowly, in no small part due to the *ad hoc* nature of the metrics we have developed. We hope that period is at an end.

Further, the metrics presented here are the foundation of a much larger framework that has seen deep exploration in the species diversity world. By embracing this framework for Multifunctionality research, we open up new vistas for ecology. Some are small - what are the consequences and best choices for  $\tau$ , how we calculate distance matrices, at what scale to evaluate correlation between functions, should we incorporate changes in correlation and distance matrices as an ecological response in and of themselves, how we standardize functional measurements, and many more. Other new areas of inquiry are quite large. Can we use this framework to begin to address the problem of unmeasured functions as we deal with unmeasured species (Chao et al. 2014b)? Can we use it to think about turnover in multifunctionality

across space and time as we do with beta diversity (Chao and Ricotta 2019) or think about partitioning landscape multifunctionality into different components (Jost 2007)? Particularly as we think about global change's impacts on ecosystem services at multiple scales, this framework lays out fruitful avenues for future exploration.

Ultimately, we feel that the proliferation of univariate multifunctionality metrics without strong theoretical underpinning has caused a great deal of confusion about how to measure multifunctionality. We hope that this piece will provide the field of multifunctionality with a way out of its current state of division and confusion. Further, we hope it provides food for additional theory that addresses the causes and consequences of ecosystem multifunctionality, something that is currently sorely lacking but highly relevant to policy and management (e.g., the efforts of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services). We have been heartened by the idea, leaving the cradle of the field of biodiversity and ecosystem function, and feel that it has the promise to provide a holistic unifying concept for anyone interested in capturing a snapshot of system dynamics in a single meaningful metric with direct ties to the beautifully developing field of diversity partitioning. Much is to be done on honing the particulars of this approach, but we feel it offers a strong theory-driven unified approach that will enable the field of Multifunctionality research to move forward swiftly.

## **Acknowledgements**

We thank Mark Urban and Morgan Tingley for organizing the workshop that began informal discussions between JEKB and RB leading to this manuscript. We thank Christian Alsterberg for giving reason for FR to travel to Boston where FR and JEKB realized that the three authors had independently come to the same conclusions and should collaborate instead of working in competition. It's how science should be. We also thank funding from multiple seminar series and statistical workshops that have kept the

collaboration going. This work was partially supported by the National Science Foundation as part of the  
PIE-LTER Program (award #1637630).

## Data Access

In addition to the appendices, see [https://github.com/jebyrnes/new\\_multifunc\\_metric/](https://github.com/jebyrnes/new_multifunc_metric/) for additional code,  
notes, and co-author discussions related to this manuscript. The code to implement all methods here is  
part of the *multifunc* package in R and can be found at <https://jebyrnes.github.io/multifunc/> with tutorials.

## References

- Angelini, C., van der Heide, T., Griffin, J. N., Morton, J. P., Derksen-Hooijberg, M., Lamers, L. P. M.,  
Smolders, A. J. P. and Silliman, B. R. 2015. Foundation species' overlap enhances biodiversity  
and multifunctionality from the patch to landscape scale in southeastern United States salt  
marshes. - Proc. R. Soc. B Biol. Sci. 282: 20150421.
- Binder, S., Isbell, F., Polasky, S., Catford, J. A. and Tilman, D. 2018. Grassland biodiversity can pay. -  
PNAS: 201712874.
- Bradford, M. A., Wood, S. A., Bardgett, R. D., Black, H. I. J., Bonkowski, M., Eggers, T., Grayston, S. J.,  
Kandeler, E., Manning, P., Setälä, H. and Jones, T. H. 2014a. Discontinuity in the responses of  
ecosystem processes and multifunctionality to altered soil community composition. - PNAS 111:  
14478–14483.
- Bradford, M. A., Wood, S. A., Bardgett, R. D., Black, H. I. J., Bonkowski, M., Eggers, T., Grayston, S. J.,  
Kandeler, E., Manning, P., Setälä, H. and Jones, T. H. 2014b. Reply to Byrnes et al.: Aggregation  
can obscure understanding of ecosystem multifunctionality. - PNAS 111: E5491–E5491.

346 Brandt, P., Abson, D. J., DellaSala, D. A., Feller, R. and von Wehrden, H. 2014. Multifunctionality and  
347 biodiversity: Ecosystem services in temperate rainforests of the Pacific Northwest, USA. - Biol.  
348 Conserv. 169: 362–371.

349 Byrnes, J. E. K., Gamfeldt, L., Isbell, F., Lefcheck, J. S., Griffin, J. N., Hector, A., Cardinale, B. J.,  
350 Hooper, D. U., Dee, L. E. and Emmett Duffy, J. 2014a. Investigating the relationship between  
351 biodiversity and ecosystem multifunctionality: challenges and solutions. - Methods Ecol. Evol. 5:  
352 111–124.

353 Byrnes, J., Lefcheck, J. S., Gamfeldt, L., Griffin, J. N., Isbell, F. and Hector, A. 2014b. Multifunctionality  
354 does not imply that all functions are positively correlated. - PNAS 111: E5490.

355 Chao, A. and Ricotta, C. 2019. Quantifying evenness and linking it to diversity, beta diversity, and  
356 similarity. - Ecology 100: e02852.

357 Chao, A., Chiu, C.-H. and Jost, L. 2014a. Unifying Species Diversity, Phylogenetic Diversity, Functional  
358 Diversity, and Related Similarity and Differentiation Measures Through Hill Numbers. - Annu.  
359 Rev. Ecol. Evol. Syst. 45: 297–324.

360 Chao, A., Gotelli, N. J., Hsieh, T. C., Sander, E. L., Ma, K. H., Colwell, R. K. and Ellison, A. M. 2014b.  
361 Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in  
362 species diversity studies. - Ecol. Monogr. 84: 45–67.

363 Chao, A., Chiu, C.-H., Villéger, S., Sun, I.-F., Thorn, S., Lin, Y.-C., Chiang, J.-M. and Sherwin, W. B.  
364 2019. An attribute-diversity approach to functional diversity, functional beta diversity, and related  
365 (dis)similarity measures. - Ecol. Monogr. 89: e01343.

366 Dice, L. R. 1945. Measures of the Amount of Ecologic Association Between Species. - Ecology 26: 297–  
367 302.

368 Dooley, Á., Isbell, F., Kirwan, L., Connolly, J., Finn, J. A. and Brophy, C. 2015. Testing the effects of  
369 diversity on ecosystem multifunctionality using a multivariate model (Y Buckley, Ed.). - Ecol.  
370 Lett. 18: 1242–1251.

371 Gamfeldt, L. and Roger, F. 2017. Revisiting the biodiversity–ecosystem multifunctionality relationship. -



372 Nat. Ecol. Evol. 1: 1–7.  
 373 Gamfeldt, L., Hillebrand, H. and Jonsson, P. R. 2008. Multiple functions increase the importance of  
 374 biodiversity for overall ecosystem functioning. - Ecology 89: 1223–1231.  
 375 Hector, A. and Bagchi, R. 2007. Biodiversity and ecosystem multifunctionality. - Nature 448: 188–190.  
 376 Hensel, M. J. S. and Silliman, B. R. 2013. Consumer diversity across kingdoms supports multiple  
 377 functions in a coastal ecosystem. - PNAS 110: 20621–20626.  
 378 Hill, M. O. 1973. Diversity and Evenness: A Unifying Notation and Its Consequences. - Ecology 54:  
 379 427–432.  
 380 Hölting, L., Beckmann, M., Volk, M. and Cord, A. F. 2019. Multifunctionality assessments – More than  
 381 assessing multiple ecosystem functions and services? A quantitative literature review. - Ecol.  
 382 Indic. 103: 226–235.  
 383 Jost, L. 2006. Entropy and diversity. - Oikos 113: 363–375.  
 384 Jost, L. 2007. Partitioning Diversity into Independent Alpha and Beta Components. - Ecology 88: 2427–  
 385 2439.  
 386 Macarthur, R. H. 1965. Patterns of Species Diversity. - Biol. Rev. 40: 510–533.  
 387 Maestre, F. T., Quero, J. L., Gotelli, N. J., Escudero, A., Ochoa, V., Delgado-Baquerizo, M., Garcia-  
 388 Gomez, M., Bowker, M. A., Soliveres, S., Escolar, C., Garcia-Palacios, P., Berdugo, M.,  
 389 Valencia, E., Gozalo, B., Gallardo, A., Aguilera, L., Arredondo, T., Blones, J., Boeken, B., Bran,  
 390 D., Conceicao, A. A., Cabrera, O., Chaieb, M., Derak, M., Eldridge, D. J., Espinosa, C. I.,  
 391 Florentino, A., Gaitan, J., Gatica, M. G., Ghiloufi, W., Gomez-Gonzalez, S., Gutierrez, J. R.,  
 392 Hernandez, R. M., Huang, X., Huber-Sannwald, E., Jankju, M., Miriti, M., Moneris, J., Mau, R.  
 393 L., Morici, E., Naseri, K., Ospina, A., Polo, V., Prina, A., Pucheta, E., Ramirez-Collantes, D. A.,  
 394 Romao, R., Tighe, M., Torres-Diaz, C., Val, J., Veiga, J. P., Wang, D. and Zaady, E. 2012. Plant  
 395 Species Richness and Ecosystem Multifunctionality in Global Drylands. - Science 335: 214–218.  
 396 Magurran, A. E. and McGill, B. J. 2010. Biological Diversity: Frontiers in Measurement and Assessment.  
 397 - OUP Oxford.

398 Manning, P., van der Plas, F., Soliveres, S., Allan, E., Maestre, F. T., Mace, G., Whittingham, M. J. and  
 399 Fischer, M. 2018. Redefining ecosystem multifunctionality. - *Nat. Ecol. Evol.* 2: 427–436.  
 400 Meyer, S. T., Ptacnik, R., Hillebrand, H., Bessler, H., Buchmann, N., Ebeling, A., Eisenhauer, N., Engels,  
 401 C., Fischer, M., Halle, S., Klein, A.-M., Oelmann, Y., Roscher, C., Rottstock, T., Scherber, C.,  
 402 Scheu, S., Schmid, B., Schulze, E.-D., Temperton, V. M., Tschardtke, T., Voigt, W., Weigelt, A.,  
 403 Wilcke, W. and Weisser, W. W. 2018. Biodiversity–multifunctionality relationships depend on  
 404 identity and number of measured functions. - *Nat. Ecol. Evol.* 2: 44–49.  
 405 Nelson, E., Mendoza, G., Regetz, J., Polasky, S., Tallis, H., Cameron, Dr., Chan, K. M., Daily, G. C.,  
 406 Goldstein, J., Kareiva, P. M., Lonsdorf, E., Naidoo, R., Ricketts, T. H. and Shaw, Mr. 2009.  
 407 Modeling multiple ecosystem services, biodiversity conservation, commodity production, and  
 408 tradeoffs at landscape scales. - *Front. Ecol. Environ.* 7: 4–11.  
 409 Poos, M. S., Walker, S. C. and Jackson, D. A. 2009. Functional-diversity indices can be driven by  
 410 methodological choices and species richness. - *Ecology* 90: 341–347.  
 411 R Core Development Team 2008. R: A Language and Environment for Statistical Computing.  
 412 Ramus, A. P., Silliman, B. R., Thomsen, M. S. and Long, Z. T. 2017. An invasive foundation species  
 413 enhances multifunctionality in a coastal ecosystem. - *PNAS* 114: 8580–8585.  
 414 Rodríguez-Loinaz, G., Alday, J. G. and Onaindia, M. 2015. Multiple ecosystem services landscape index:  
 415 A tool for multifunctional landscapes conservation. - *J. Environ. Manage.* 147: 152–163.  
 416 Sørensen, T. 1948. A method of establishing groups of equal amplitude in plant sociology based on  
 417 similarity of species content and its application to analyses of the vegetation on Danish commons.  
 418 - *Biol. Skr.* 5: 1–34.  
 419 Stürck, J. and Verburg, P. H. 2017. Multifunctionality at what scale? A landscape multifunctionality  
 420 assessment for the European Union under conditions of land use change. - *Landsc. Ecol.* 32: 481–  
 421 500.