

Laser-Driven Proton Sources Using Low-Density Foam Targets And Their Applications to Alpha Particle Generation

Anton Moroz¹, Katarzyna Batani², Gerard Malka¹, Dimitri Batani¹, Jeffrey Commenge¹, Howel Larreur¹, Lidia Giuditta Pavan¹, Alessandro Milani¹, Didier Raffestin¹, Sebastian Busch³, Artem Martynenko³, Olga Rosmej³, Parysatis Tavana³, Karim Zarouk³, Dieter Hoffman^{3,8}, Evgeny Filippov⁴, Alejandro Huerta⁴, Luca Volpe^{4,5}, Yair Ferber⁶, Noaz Nissim⁶, Massimo Alonzo⁷, Fabrizio Consoli⁷, Claudio Verona¹², Benoit Grau¹², Massimiliano Sciscio⁷, Wei Kang⁹, Daniele Margarone¹⁰, Marco Tosca¹⁰, Sergey Pikuz¹¹, Angelo Raso¹², Bing Liu¹³

1 Celia, University of Bordeaux, France

2 IPPLM Warsaw, Poland

3 GSI Darmstadt, Germany

4 CLPU, Salamanca, Spain

5 Universidad Politecnica Madrid, Spain

6 Soreq NRC, Israel,

7 ENEA Frascati, Italy

8 Xi'an Jiaotong, University, China

9 Peking University, Beijing, China,

10 ELI Beamlines, Prague, Czech Republic

11 HB11 Energy, Australia

12 University of Roma Tor Vergata, Italy

13 ENN, China

In a recent experiment at the PHELIX facility (GSI), we investigated proton generation via the interaction of high-intensity picosecond laser pulses with low-density foam targets. A primary objective was to study the process of super-ponderomotive acceleration of electrons in plasmas with very long scale length and the effect they have on proton acceleration in laser-produced plasmas. This talk presents the results on the evaluation of laser-accelerated proton energy spectra and simulation alpha particle yields produced by a boron nitride target. We discuss the estimated reaction rates based on data from multiple diagnostics, considering the interaction probabilities and target parameters. Our findings suggest that cold foams near the critical density regime increase the generation of proton beams in the low-energy range which is well-suited for triggering p-¹¹B fusion reactions. These results highlight the potential of this approach for producing high-brightness α -particle sources and advancing aneutronic fusion research.

This work has been carried out within the framework of the COST Action CA21128- PROBONO “PROton BORon Nuclear fusion: from energy production to medical applications”, supported by COST (European Cooperation in Science and Technology - www.cost.eu)

This project has received funding from the European Union's Horizon Europe Research and Innovation Programme under the Marie Skłodowska-Curie Grant Agreement 101081465