

The Adolescent Scar of Early-Life Disaster

José Elías Durán Roa

October 5, 2026

Tables and instrument appendix — draft for discussion

OVERVIEW

The 27 February 2010 earthquake, of magnitude 8.8, struck a large part of Chile. Fourteen years later the ELPI, Chile's early-childhood panel, returned to children who were between six months and four years old that day and, for the first time, asked the adolescents themselves about their mental health. We find that the earthquake left a scar whose depth depends on the age at which it struck. Adolescents who lived through it at age two report fewer anxiety and depression symptoms at ages 15–18 than adolescents struck as infants, when attachment and stress regulation are forming, or at ages three and four, when children form their first lasting memories, a U-shaped profile whose trough is consistent with the window of infantile amnesia (Terr 1988; Bauer 2007; Alberini and Travaglia 2017). Girls explain between 81 and 95 percent of this profile, and their caregivers do not perceive these problems, reporting no gradient at any age.

Beyond mental health, the 2024 wave lets us follow the developmental damage that Gillmore (2026) documents after the same earthquake. His cognitive result survives the passage to adolescence, since the vocabulary loss he measures at ages 7–11, of 0.17 standard deviations, is still there at ages 15–18, at 0.24. His non-cognitive result does not survive it, since the caregiver-reported behavior problems he finds at ages 7–11 are gone at ages 15–18 (Table 2). By contrast, the mental-health effects documented in this paper appear in the symptoms that adolescents report about themselves, while for the same adolescents the caregiver reports used by Gillmore (2026) show no effect.

The magnitude is moderate. Measured with the PHQ-4, a four-question screener of low mood, loss of interest, nervousness and worry that the adolescents answer in private, exposure at age two lowers the score by 0.39 points on its 0-to-12 scale relative to exposure before age one, or 0.147 standard deviations, statistically significant at the 5 percent level (column 3 of Panel A in Table 3). The index averages 4.31 points in the sample, so the gap corresponds to 9 percent of the mean and to about a third of the 1.11-point distance between girls and boys, and the distance by mother's education is essentially zero. The anxiety half of the index moves the most, since the probability of a positive GAD-2 screen falls by 7.3 percentage points relative to a base of 27.3 percent.

Among girls the PHQ-4 gap doubles to 0.300 standard deviations and appears in both halves of the index, with a fall of 13.7 percentage points in the anxiety screen and of 11.7 in the depression screen, while among boys every estimate is close to zero and statistically insignificant (Table 5). Among caregivers the same design finds nothing at any age (Table 4), which is consistent with symptoms that adolescents feel and adults do not observe. Figure 1 draws the full age profile.

EMPIRICAL STRATEGY

Every adolescent of the 2024 wave lived through the earthquake as a child, and identification rests on the two things that vary across them, how hard their municipality was shaken and how old they were when it struck, with age grouped as 0–11 months (the omitted group), 12–23, 24–35 and 36–59. We compare the outcome gap between two exposure ages within the municipalities the earthquake hit against the same gap within the municipalities it did not hit, so municipality fixed effects keep every comparison local and birth-cohort fixed effects absorb everything a cohort shares, including its age at the survey. All tables estimate versions of

$$Y_{imc} = \alpha + \sum_a \beta_a (\text{Bin}_{ai} \times D_m) + \sum_a \gamma_a \text{Bin}_{ai} + \psi_m + \theta_c + X_i' \delta + \varepsilon_{imc}, \quad (1)$$

where Y_{imc} is the outcome of adolescent i of birth cohort c in municipality m ; Bin_{ai} are the exposure-age groups; D_m equals one in the six regions where the official intensity study records destructive shaking (V to IX and the Metropolitan Region; Astroza et al. 2010); ψ_m and θ_c are municipality-of-selection and birth-cohort fixed effects; X_i are household controls; and ε_{imc} is the error term. The coefficients of interest are the β_a , the effect of having lived the earthquake in an affected municipality at age a relative to the children exposed before age one. The main effects γ_a absorb the outcome differences across exposure-age groups that are common to affected and unaffected municipalities, differences that exist mechanically because a child who was older at the earthquake is also older at the 2024 interview, and β_a is therefore identified only from how that age profile differs between the two zones. Standard errors are clustered by municipality of selection, and estimates use the 2024 sampling weights.

Table 1
Descriptive statistics.

	(1) Non-Earthquake	(2) Earthquake	(3) Difference
Girl	0.489 (0.500)	0.489 (0.500)	−0.000 (0.011)
Age at the earthquake (months)	26.930 (12.710)	27.098 (12.614)	0.169 (0.254)
Mother's age (2010)	28.980 (7.066)	29.676 (7.116)	0.695*** (0.196)
Mother's schooling (2010)	10.312 (3.182)	10.311 (3.180)	−0.002 (0.211)
HH size (2010)	4.865 (1.641)	4.948 (1.686)	0.083 (0.074)
Rural household (2010)	0.080 (0.271)	0.094 (0.292)	0.014 (0.035)
Children of the mother (2012)	2.165 (1.072)	2.155 (1.075)	−0.010 (0.029)
Prior mental health, mother (2012)	0.091 (0.288)	0.125 (0.331)	0.034*** (0.012)
Prior mental health, father (2012)	0.023 (0.150)	0.027 (0.161)	0.004 (0.004)
Prior mental health, relative (2012)	0.123 (0.328)	0.152 (0.359)	0.029* (0.017)
Observations	2,349	7,654	10,003

Notes: ELPI uses sampling weight at national level. The sample are the 10,003 adolescents of the 2024 wave, born January 2006–August 2009, aged 6 to 49 months on the day of the earthquake and 15 to 18 at the 2024 interview. Sex and ages are fixed at birth and the 2010 rows come from the pre-earthquake baseline; the 2012 rows are the family size and the lifetime conditions reported in the 2012 wave, the controls of the main specification, and are missing for adolescents without that interview. Mental health rows mark a reported diagnosis of depression, anxiety, bipolar disorder, schizophrenia, a personality disorder or an autism spectrum disorder. Earthquake municipalities are those of the six regions where the official intensity study records destructive shaking (V, VI, VII, VIII, IX and Metropolitan; Astroza et al. 2010). Column 3 is the difference from a regression on the earthquake indicator; standard errors clustered at the municipality of selection appear in parentheses—number of clusters: 116. * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table 2

The impact of the earthquake on cognitive and non-cognitive development in the medium- and long-term.

	Medium-term effect (Gillmore) 2017 wave		Long-term effect 2024 wave	
	(1)	(2)	(3)	(4)
<i>Panel A: Peabody</i>				
Affected×Earthquake	−0.174** (0.0686)	−0.280*** (0.0866)	−0.240** (0.104)	−0.412** (0.180)
Observations	14,069	14,069	13,779	13,779
<i>Panel B: CBCL2</i>				
Affected×Earthquake	−0.129* (0.0763)	−0.220** (0.105)	0.084 (0.0884)	−0.066 (0.125)
Observations	11,568	11,568	11,038	11,038
Municipality FE	✓	✓	✓	✓
Birth Cohort FE	✓	✓	✓	✓
Child-Mother-HH control	✓	✓	✓	✓
Municipality linear trends	×	✓	×	✓

Notes: Each column represents a separate regression. The dependent variable is the test score measured in standard deviations. Peabody is the Peabody Test Score and CBCL is the Child Behavior Check List measure, oriented so that higher means fewer problems. Columns 1 and 2 reproduce Columns 3 and 4 of Table 3 in Gillmore (2026): children exposed between in utero and age four, with scores measured seven years later, at ages 7–11; our replication of his Column 3 on the public ELPI files gives −0.160 for Peabody and −0.103 for CBCL2. Columns 3 and 4 estimate the same two specifications on the 2024 wave: the same exposed children measured fourteen years later, at ages 15–18, against the comparison group conceived after the earthquake and measured in 2017; pooling the two waves adds survey-wave fixed effects, as in Gillmore’s short-term table. Standard errors clustered at the municipal level appear in parentheses—number of clusters: 145 in Columns 1 and 2 and 116 in Columns 3 and 4. The estimates use sampling weights. Child/Mother/HH characteristics include gender, age at the moment of the test (in months), birth order, mothers’ age and schooling, father present, number of household members, prior mental health condition for mother, father and relative. * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.




Figure 1. Estimates by Age at Exposure

Notes: Each marker is a β_a coefficient of equation (1) with the controls of column 3 of Table 3, with its 95 percent confidence interval. The reference group, drawn at zero by construction, are the children struck at 0–11 months — already born; children in utero during the earthquake are not in the 2024 wave, which follows only the cohort born 2006–2009. The PHQ-4 is in standard deviations and the GAD-2 screen in probability units. Standard errors clustered by municipality of selection in parentheses.

Table 3

Age at Exposure to the Earthquake and Self-Reported Mental Health at Ages 15–18

	(1)	(2)	(3)	(4)
<i>Panel A. Mental health: PHQ-4 index, four questions on low mood, loss of interest, nervousness and worry (0–12, z)</i>				
Exposed at age 1	–0.051 (0.080)	–0.044 (0.078)	–0.045 (0.078)	–0.030 (0.079)
Exposed at age 2	–0.144** (0.072)	–0.153** (0.069)	–0.147** (0.069)	–0.120 (0.076)
Exposed at ages 3–4	–0.038 (0.064)	–0.048 (0.060)	–0.044 (0.060)	–0.011 (0.063)
<i>Panel B. The anxiety half of the index: positive GAD-2 screen (its two anxiety questions, subscore of 3 or more)</i>				
Exposed at age 1	–0.014 (0.032)	–0.012 (0.032)	–0.012 (0.033)	–0.018 (0.035)
Exposed at age 2	–0.073** (0.032)	–0.076** (0.032)	–0.073** (0.032)	–0.064* (0.035)
Exposed at ages 3–4	–0.015 (0.027)	–0.018 (0.028)	–0.016 (0.028)	–0.015 (0.030)
<i>Panel C. The depression half of the index: positive PHQ-2 screen (its two depression questions, subscore of 3 or more)</i>				
Exposed at age 1	–0.017 (0.039)	–0.014 (0.038)	–0.014 (0.038)	–0.020 (0.041)
Exposed at age 2	–0.046 (0.042)	–0.050 (0.041)	–0.048 (0.041)	–0.073 (0.045)
Exposed at ages 3–4	–0.008 (0.032)	–0.011 (0.031)	–0.010 (0.031)	–0.017 (0.035)
Municipality fixed effects	Yes	Yes	Yes	Yes
Birth-cohort fixed effects	Yes	Yes	Yes	Yes
Pre-earthquake controls	No	Yes	Yes	Yes
2012 controls	No	No	Yes	Yes
Excludes the Metropolitan Region	No	No	No	Yes
Observations	9,996	9,996	9,996	6,430
Selection municipalities (clusters)	116	116	116	67

Notes: Ordinary least squares estimates of equation (1) on the adolescents of the 2024 ELPI wave; seven of the 10,003 adolescents of Table 1 drop for incomplete PHQ-4 answers. Panels B and C disaggregate the Panel A index into its two pairs of questions, expressed as clinical screens; Appendix Table A1 lists every question verbatim. Exposure-age groups give the child's age when the earthquake struck, defined in months of age as 0–11 (before age 1, the omitted group), 12–23 (age 1), 24–35 (age 2) and 36–59 (ages 3–4). Each coefficient compares the gap between the age group in the row and the omitted group in affected municipalities with the same gap in unaffected municipalities. The affected zone comprises the six regions where the official intensity study records destructive shaking (V, VI, VII, VIII, IX and Metropolitan; Astroza et al. 2010). Column 1 includes exposure-age, municipality-of-selection and birth-cohort fixed effects; column 2 adds pre-earthquake controls (sex, age in months, and the 2010 household baseline: mother's education and age, household size, rurality); column 3 adds 2012 controls (number of children of the mother, family history of mental illness), with missing-value indicators; column 4 estimates column 3 without the Metropolitan Region. Estimates use the 2024 sampling weights. The sample is the same in every panel; column 4 has 6,430 adolescents in 67 municipalities. Standard errors clustered by municipality of selection in parentheses. *** Significant at the 1 percent level. ** Significant at the 5 percent level. * Significant at the 10 percent level.

Table 4
Caregiver-Reported Outcomes

	(1)	(2)	(3)	(4)
<i>Panel A. CBCL internalizing score in 2024: caregiver checklist (T, z)</i>				
Exposed at age 1	−0.057 (0.099)	−0.047 (0.096)	−0.055 (0.095)	−0.024 (0.100)
Exposed at age 2	−0.121 (0.074)	−0.121 (0.075)	−0.109 (0.075)	−0.131 (0.084)
Exposed at ages 3–4	−0.086 (0.069)	−0.084 (0.068)	−0.077 (0.069)	−0.069 (0.074)
Observations	9,970	9,970	9,970	6,412
<i>Panel B. Within-child change in the same CBCL score, 2017 to 2024 (z by wave)</i>				
Exposed at age 1	−0.030 (0.131)	−0.011 (0.125)	−0.012 (0.124)	−0.020 (0.128)
Exposed at age 2	−0.009 (0.104)	−0.004 (0.102)	−0.002 (0.102)	−0.045 (0.110)
Exposed at ages 3–4	−0.040 (0.109)	−0.049 (0.106)	−0.050 (0.106)	−0.093 (0.112)
Observations	7,169	7,169	7,169	4,901
Municipality fixed effects	Yes	Yes	Yes	Yes
Birth-cohort fixed effects	Yes	Yes	Yes	Yes
Pre-earthquake controls	No	Yes	Yes	Yes
2012 controls	No	No	Yes	Yes
Excludes the Metropolitan Region	No	No	No	Yes
Selection municipalities (clusters)	116	116	116	67

Notes: Same design and specifications as Table 3, with the caregiver’s report of the child as the outcome. The CBCL internalizing score is reported by the main caregiver, in T units, standardized within age. Panel B uses the change between the 2017 measurement (ages 8–12) and the 2024 measurement (ages 15–18) of the same child, each standardized within its wave and age; with two periods this is equivalent to a child fixed-effects estimate. Neither panel shows an exposure-age gradient: the scar of Table 3 appears only in what adolescents report in private. Exposure-age groups give the child’s age when the earthquake struck, defined in months of age as 0–11 (before age 1, the omitted group), 12–23 (age 1), 24–35 (age 2) and 36–59 (ages 3–4). Each coefficient compares the gap between the age group in the row and the omitted group in affected municipalities with the same gap in unaffected municipalities. Column 1 includes exposure-age, municipality-of-selection and birth-cohort fixed effects; column 2 adds pre-earthquake controls (sex, age in months, and the 2010 household baseline: mother’s education and age, household size, rurality); column 3 adds 2012 controls (number of children of the mother, family history of mental illness), with missing-value indicators; column 4 estimates column 3 without the Metropolitan Region. Estimates use the 2024 sampling weights. Standard errors clustered by municipality of selection in parentheses. *** Significant at the 1 percent level. ** Significant at the 5 percent level. * Significant at the 10 percent level.

Table 5

The effect of exposure at age two on adolescent mental health, by sex, mother's education and area of residence.

	Sex		Mother's education		Household in 2010	
	Girls (1)	Boys (2)	Secondary (3)	Tertiary (4)	Urban (5)	Rural (6)
Panel A: PHQ-4 index						
Exposed at age 2	−0.300*** (0.100)	−0.016 (0.098)	−0.107 (0.074)	−0.201 (0.188)	−0.139* (0.076)	−0.238 (0.172)
Observations	4,961	5,035	7,755	2,053	9,005	991
Panel B: Anxiety, positive GAD-2 screen						
Exposed at age 2	−0.137*** (0.047)	−0.026 (0.048)	−0.082** (0.034)	−0.021 (0.098)	−0.068** (0.034)	−0.113 (0.092)
Observations	4,961	5,035	7,755	2,053	9,005	991
Panel C: Depression, positive PHQ-2 screen						
Exposed at age 2	−0.117** (0.057)	0.016 (0.045)	−0.030 (0.043)	−0.082 (0.089)	−0.046 (0.045)	−0.073 (0.084)
Observations	4,961	5,035	7,755	2,053	9,005	991
Municipality FE	✓	✓	✓	✓	✓	✓
Birth Cohort FE	✓	✓	✓	✓	✓	✓
Pre-earthquake controls	✓	✓	✓	✓	✓	✓
2012 controls	✓	✓	✓	✓	✓	✓

Notes: Each column represents a separate regression of equation (1) with the controls of column 3 of Table 3, estimated on the subsample named in the column heading. The coefficient reported is the effect of exposure at age two in an affected municipality, relative to exposure at 0–11 months. The PHQ-4 index is measured in standard deviations; the GAD-2 and PHQ-2 screens equal one for a subscore of three or more on the corresponding pair of questions. Mother's education is measured in 2010, and secondary means twelve years of schooling or less. Weighting girls and boys by their shares of the sample, girls account for 95 percent of the age-two coefficient and for 81 percent of the gap between exposure at ages 3–4 and at age two. The PHQ-4 index averages 4.88 points for girls and 3.76 for boys. Standard errors clustered by municipality of selection appear in parentheses. The estimates use sampling weights. * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table E1

Exploratory outcomes I: mental health and well-being.

	Age 1 (1)	Age 2 (2)	Ages 3–4 (3)	Mean (4)	Observations (5)
Panel A: Mental health					
PHQ-4 symptoms (z)	−0.045 (0.078)	−0.147** (0.069)	−0.044 (0.060)	0.000	9,996
Positive anxiety screen, GAD-2 (0/1)	−0.012 (0.033)	−0.073** (0.032)	−0.016 (0.028)	0.273	9,996
Positive depression screen, PHQ-2 (0/1)	−0.014 (0.038)	−0.048 (0.041)	−0.010 (0.031)	0.346	9,996
Internalizing problems, CBCL (z)	−0.026 (0.082)	−0.071 (0.076)	−0.044 (0.071)	0.000	9,552
Externalizing problems, CBCL (z)	−0.010 (0.079)	−0.087 (0.061)	−0.056 (0.061)	0.000	9,552
Attention problems, CBCL (z)	−0.068 (0.088)	−0.098 (0.074)	−0.082 (0.061)	0.000	9,552
Panel B: Well-being					
Resilience (z)	0.061 (0.077)	0.141* (0.079)	0.098 (0.068)	0.000	9,996
Satisfaction with life in general (z)	−0.003 (0.067)	0.069 (0.067)	0.031 (0.060)	0.000	9,996
Satisfaction with family life (z)	0.086* (0.050)	0.098* (0.056)	0.116* (0.062)	0.000	9,996
Satisfaction with friends (z)	0.048 (0.068)	0.033 (0.063)	0.030 (0.067)	0.000	9,996
Satisfaction with school (z)	−0.014 (0.067)	0.032 (0.070)	0.076 (0.073)	0.000	9,561
Satisfaction with oneself (z)	0.055 (0.069)	0.157** (0.074)	0.123* (0.064)	0.000	9,996
Satisfaction with the neighborhood (z)	0.008 (0.077)	0.040 (0.083)	−0.011 (0.070)	0.000	9,996
Satisfaction with one's body (z)	0.044 (0.078)	0.109 (0.073)	0.053 (0.056)	0.000	9,996
Poor self-rated health (z)	−0.086 (0.066)	−0.072 (0.077)	−0.054 (0.070)	0.000	9,996

Notes: Each row is a separate regression of equation (1) with the controls of column 3 of Table 3. Columns 1 to 3 report the coefficient on exposure at the age named in the column interacted with the affected zone, relative to exposure at 0–11 months; column 4 reports the sample mean of the outcome. The design identifies differences across exposure ages, not the level of the effect, so a flat row is consistent both with no effect and with an effect common to all exposure ages. Variables marked (z) are standardized; Appendix Tables A1 to A10 list the items behind every index. Standard errors clustered by municipality of selection appear in parentheses. The estimates use sampling weights. * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table E2

Exploratory outcomes II: cognition, school and peers.

	Age 1 (1)	Age 2 (2)	Ages 3–4 (3)	Mean (4)	Observations (5)
<i>Panel A: Cognition and school</i>					
Vocabulary, TVIP (z)	−0.085 (0.089)	−0.086 (0.079)	0.034 (0.077)	0.000	9,990
Attends school (0/1)	0.009* (0.005)	0.006 (0.006)	0.001 (0.013)	0.956	10,003
Grade average last year (1–7)	−0.031 (0.055)	−0.002 (0.048)	0.009 (0.046)	5.87	9,561
Doing well at school (z)	−0.032 (0.072)	0.138** (0.067)	0.119* (0.070)	0.000	9,176
Likes going to school (z)	−0.068 (0.079)	0.025 (0.101)	0.056 (0.106)	0.000	9,176
Likes the teachers (z)	−0.082 (0.074)	0.021 (0.077)	0.037 (0.086)	0.000	9,561
Expects a university degree (0/1)	0.002 (0.043)	0.039 (0.036)	0.056 (0.034)	0.756	8,978
Does not know how far they will study (0/1)	0.009 (0.031)	0.014 (0.022)	−0.004 (0.022)	0.102	9,996
<i>Panel B: Peers and victimization</i>					
Peer victimization and class climate (z)	0.028 (0.060)	−0.047 (0.071)	0.037 (0.063)	0.000	9,176
Cyber-victimization, last 12 months (0/1)	−0.015 (0.029)	−0.007 (0.023)	−0.005 (0.025)	0.150	9,985
Dating violence, ever dated (0/1)	0.025 (0.019)	0.005 (0.020)	0.028 (0.021)	0.051	7,101

Notes: Each row is a separate regression of equation (1) with the controls of column 3 of Table 3. Columns 1 to 3 report the coefficient on exposure at the age named in the column interacted with the affected zone, relative to exposure at 0–11 months; column 4 reports the sample mean of the outcome. The design identifies differences across exposure ages, not the level of the effect, so a flat row is consistent both with no effect and with an effect common to all exposure ages. Variables marked (z) are standardized; Appendix Tables A1 to A10 list the items behind every index. Standard errors clustered by municipality of selection appear in parentheses. The estimates use sampling weights. * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table E3

Exploratory outcomes III: risk behaviors and daily life.

	Age 1 (1)	Age 2 (2)	Ages 3–4 (3)	Mean (4)	Observations (5)
Panel A: Risk behaviors					
Smoked tobacco, last 12 months (0/1)	0.012 (0.016)	−0.002 (0.015)	−0.022 (0.024)	0.128	9,995
Drank alcohol, last 12 months (0/1)	−0.038 (0.032)	−0.037 (0.040)	−0.078** (0.031)	0.398	9,995
Used cannabis, last 12 months (0/1)	0.013 (0.014)	0.014 (0.015)	0.008 (0.021)	0.105	9,995
Used other drugs, last 12 months (0/1)	0.014 (0.015)	0.007 (0.013)	0.012 (0.013)	0.039	9,995
No contraception at last sex (0/1)	0.005 (0.073)	0.015 (0.074)	−0.000 (0.068)	0.107	3,256
Panel B: Daily life					
Hours of sleep on weekdays	0.140 (0.104)	0.010 (0.102)	−0.017 (0.112)	7.90	9,295
Stays up on the phone in bed (z)	0.074 (0.084)	0.105 (0.065)	0.057 (0.072)	0.000	9,996
Time on social media (z)	0.098 (0.067)	0.131* (0.070)	0.063 (0.075)	0.000	9,996
Days of physical activity, last week	−0.056 (0.145)	−0.158 (0.143)	−0.033 (0.154)	2.85	9,996
Time reading (z)	−0.027 (0.064)	−0.032 (0.066)	−0.090 (0.077)	0.000	9,996
Days eating junk food, last week	0.090 (0.093)	−0.001 (0.092)	−0.038 (0.086)	1.81	9,996
Portions of fruit per day	0.006 (0.145)	0.255* (0.152)	0.016 (0.152)	2.77	9,994

Notes: Each row is a separate regression of equation (1) with the controls of column 3 of Table 3. Columns 1 to 3 report the coefficient on exposure at the age named in the column interacted with the affected zone, relative to exposure at 0–11 months; column 4 reports the sample mean of the outcome. The design identifies differences across exposure ages, not the level of the effect, so a flat row is consistent both with no effect and with an effect common to all exposure ages. Variables marked (z) are standardized; Appendix Tables A1 to A10 list the items behind every index. Standard errors clustered by municipality of selection appear in parentheses. The estimates use sampling weights. * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table E4

Exploratory outcomes IV: family, adversity, caregiver and household.

	Age 1 (1)	Age 2 (2)	Ages 3–4 (3)	Mean (4)	Observations (5)
Panel A: Family and social life					
Parental knowledge of activities (z)	0.043 (0.062)	0.065 (0.066)	0.027 (0.063)	0.000	9,508
Relationship with main caregiver (z)	0.081 (0.075)	0.120* (0.068)	0.102 (0.073)	0.000	9,970
Confides in nobody (0/1)	−0.001 (0.023)	−0.020 (0.018)	−0.016 (0.019)	0.086	9,996
Activities with family, last week (0–5)	0.131* (0.075)	0.138 (0.085)	0.114 (0.075)	3.73	9,985
Takes part in a group or organization (0/1)	−0.044 (0.035)	−0.036 (0.037)	0.014 (0.033)	0.696	9,996
Panel B: Adverse experiences reported by the caregiver					
Number of adverse experiences (0–10)	−0.092 (0.104)	−0.264** (0.126)	−0.144 (0.098)	1.25	9,834
Witnessed fights at home (0/1)	−0.068** (0.027)	−0.099*** (0.027)	−0.078*** (0.025)	0.159	9,671
Abrupt separation from caregiver (0/1)	−0.019 (0.035)	−0.059 (0.039)	−0.065* (0.036)	0.346	9,797
Housing problems (0/1)	0.001 (0.018)	−0.010 (0.019)	0.014 (0.017)	0.036	9,885
Not enough to eat (0/1)	0.014 (0.025)	0.008 (0.023)	0.030 (0.025)	0.090	9,798
Panel C: Caregiver and household					
Caregiver depressive symptoms, CES-D (z)	−0.064 (0.069)	−0.071 (0.056)	−0.021 (0.073)	0.000	9,978
Parenting stress, PSI total (z)	−0.072 (0.070)	0.013 (0.057)	−0.032 (0.069)	0.000	9,629
Parental distress, PSI (z)	−0.126* (0.071)	0.014 (0.054)	−0.048 (0.072)	0.000	9,629
Dysfunctional interaction, PSI (z)	−0.058 (0.070)	0.025 (0.066)	−0.034 (0.072)	0.000	9,629
Parenting sense of competence (z)	−0.074 (0.075)	−0.106 (0.066)	−0.110 (0.070)	0.000	9,978
Log household income per capita	−0.087 (0.054)	−0.084 (0.054)	−0.101* (0.059)	12.35	9,967

Notes: Each row is a separate regression of equation (1) with the controls of column 3 of Table 3. Columns 1 to 3 report the coefficient on exposure at the age named in the column interacted with the affected zone, relative to exposure at 0–11 months; column 4 reports the sample mean of the outcome. The design identifies differences across exposure ages, not the level of the effect, so a flat row is consistent both with no effect and with an effect common to all exposure ages. Variables marked (z) are standardized; Appendix Tables A1 to A10 list the items behind every index. Standard errors clustered by municipality of selection appear in parentheses. The estimates use sampling weights. * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

APPENDIX. INSTRUMENTS: EVERY ITEM AND ITS SCALE

Table A1
Mental-Health Instruments: Items and Scales

	Question, verbatim (1)	Response scale (2)
<i>Panel A. PHQ-4, answered by the adolescent in private (2024 wave)</i>		
d4_1	¿Te has sentido bajoneado(a), deprimido(a), irritable o desesperanzado(a)?	0 = Nunca; 1 = Algunos días; 2 = Más de la mitad de los días; 3 = Casi todos los días
d4_2	¿Has sentido poco interés o placer al hacer las cosas?	0 = Nunca; 1 = Algunos días; 2 = Más de la mitad de los días; 3 = Casi todos los días
d4_3	¿Te has sentido muy nervioso(a), angustiado(a) o con los nervios de punta?	0 = Nunca; 1 = Algunos días; 2 = Más de la mitad de los días; 3 = Casi todos los días
d4_4	¿No has podido dejar de preocuparte o controlar la preocupación?	0 = Nunca; 1 = Algunos días; 2 = Más de la mitad de los días; 3 = Casi todos los días
<i>Panel B. Child Behavior Checklist, answered by the main caregiver about the adolescent</i>		
cbc12	Total problems scale. The survey releases the computed scores, not the copyrighted items.	T score, standardized in the analysis
cbc12 _i	Internalizing problems: the anxious/depressed, withdrawn and somatic-complaints syndrome scales.	T score, standardized
cbc12 _e	Externalizing problems: the rule-breaking and aggressive-behavior syndrome scales.	T score, standardized
cbc12 _6	Attention problems syndrome scale.	T score, standardized

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. The PHQ-4 score used in the paper is the sum of the four items rescaled to 0–3 each (0–12 in total) and standardized within age in years. The positive PHQ-2 (depression) and GAD-2 (anxiety) screens are the survey-provided indicators for a subscore of three or more on the corresponding pair of items.

Table A2
Well-Being Outcomes: Items and Scales

	Question, verbatim (1)	Response scale (2)
<i>Panel A. Resilience: Brief Resilience Scale (all six items worded positively)</i>		
d2_1	Tiendo a recuperarme rápidamente después de haber vivido situaciones difíciles	1 = Muy en desacuerdo; 2 = En desacuerdo; 3 = Neutral; 4 = De acuerdo; 5 = Muy de acuerdo
d2_2	Tengo facilidad para superar eventos estresantes	<i>Same scale</i>
d2_3	No me toma mucho tiempo recuperarme de un evento estresante	<i>Same scale</i>
d2_4	Es fácil para mí recuperarme cuando algo malo sucede	<i>Same scale</i>
d2_5	Usualmente supero situaciones difíciles con poca dificultad	<i>Same scale</i>
d2_6	Me toma poco tiempo recuperarme de las adversidades en mi vida	<i>Same scale</i>
<i>Panel B. Satisfaction, one item per domain, scale 1 to 7</i>		
d3_6	Pensando en una escala de 1 a 7, donde 1 es nada satisfecho(a) y 7 es muy satisfecho(a), ¿qué tan satisfecho(a) estás con tu vida en general?	1 = Nada Satisfecho(a); 2 = -; 3 = -; 4 = -; 5 = -; 6 = -; 7 = Muy Satisfecho(a)
d3_1	Pensando en una escala de 1 a 7, donde 1 es nada satisfecho(a) y 7 es muy satisfecho(a), ¿qué tan satisfecho(a) estás con tu vida familiar?	<i>Same scale</i>
d3_2	Pensando en una escala de 1 a 7, donde 1 es nada satisfecho(a) y 7 es muy satisfecho(a), ¿qué tan satisfecho(a) estás con tus amigos(as)?	<i>Same scale</i>
d3_3	Pensando en una escala de 1 a 7, donde 1 es nada satisfecho(a) y 7 es muy satisfecho(a), ¿qué tan satisfecho(a) estás con tu experiencia en el colegio?	<i>Same scale</i>
d3_4	Pensando en una escala de 1 a 7, donde 1 es nada satisfecho(a) y 7 es muy satisfecho(a), ¿qué tan satisfecho(a) estás contigo mismo(a)?	<i>Same scale</i>
d3_5	Pensando en una escala de 1 a 7, donde 1 es nada satisfecho(a) y 7 es muy satisfecho(a), ¿qué tan satisfecho(a) estás con el barrio donde vives en general?	<i>Same scale</i>
d3_7	Pensando en una escala de 1 a 7, donde 1 es nada satisfecho(a) y 7 es muy satisfecho(a), ¿qué tan satisfecho(a) estás con tu cuerpo?	<i>Same scale</i>
<i>Panel C. Self-rated health</i>		
d1	Considerando tu salud física, dirías que esta es:	1 = Muy mala; 2 = Mala; 3 = Regular; 4 = Buena; 5 = Muy buena

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. The resilience score is the mean of the six items, standardized; the ELPI 2024 version words all six positively, unlike the original scale, so none is reverse-coded. Each satisfaction item is standardized on its own; life satisfaction is the item on life in general. Self-rated health is standardized with its sign reversed, so that higher values mean worse health.

Table A3
Cognition and School: Items and Scales

	Question, verbatim (1)	Response scale (2)
<i>Panel A. Cognition, assessed by the interviewer</i>		
tvip	Test de Vocabulario en Imágenes Peabody (TVIP), the Spanish adaptation of the Peabody Picture Vocabulary Test: receptive vocabulary.	Standard score on Hispanic norms, standardized
<i>Panel B. School, answered by the adolescent</i>		
e1	Actualmente, ¿%nombre% asiste a algún establecimiento educacional, nivelación de estudios, jardín infantil, sala cuna u otro programa no convencional de educación parvularia?	1 = Sí; 2 = No
e3_asiste	¿Qué promedio de notas tuviste el año pasado?	1 = Menos de 4.0; 2 = Entre 4.0 y 4.4; 3 = Entre 4.5 y 4.9; 4 = Entre 5.0 y 5.4; 5 = Entre 5.5 y 5.9; 6 = Entre 6.0 y 6.4; 7 = Entre 6.5 y 7.0
e2_asiste	¿Cómo te va en el colegio?	1 = Muy mal; 2 = Mal; 3 = Más o menos; 4 = Bien; 5 = Muy bien
e5	¿Cuánto te gusta ir al colegio?	1 = No me gusta nada; 2 = No me gusta; 3 = No me gusta ni me disgusta; 4 = Me gusta; 5 = Me gusta mucho
e6	¿Cuánto te agradan tus profesores y profesoras?	1 = No me gustan nada; 2 = No me gustan; 3 = No me gustan ni me disgustan; 4 = Me gustan; 5 = Me gustan mucho
e7	Pensando en el futuro, ¿cuál es el nivel de educacional más alto que crees que vas a completar?	1 = 8° año de Educación Básica; 2 = 4° año de Educación Media Técnica...; 3 = 4° año de Educación Media Cientí...; 4 = Una carrera en un Centro de Form...; 5 = Una carrera en un Instituto Prof...; 6 = Una carrera en la Universidad; 7 = Estudios de Postgrado; 88 = No sabe; ...

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. Attendance is one when the adolescent attends an educational establishment. The grade average converts each range to its midpoint on the Chilean 1–7 scale; the items on how school goes, liking school and liking the teachers are standardized. The expected degree is one for a university degree or postgraduate studies, among adolescents who name a level; not knowing is one for answer 88.

Table A4
Peers and Victimization: Items and Scales

	Question, verbatim (1)	Response scale (2)
<i>Panel A. Peer victimization and class climate (frequency scale; standardized mean; item g1_3 reverse-coded)</i>		
g1_1	Mis compañeros(as) se burlan de mí, me dicen sobrenombres	1 = Nunca; 2 = Pocas veces; 3 = Casi siempre; 4 = Siempre
g1_2	Me siento solo(a) en el curso	<i>Same scale</i>
g1_3	Lo paso bien con mis compañeros(as) de clase	<i>Same scale</i>
g1_4	Mis compañeros(as) son muy agresivos(as)	<i>Same scale</i>
g1_5	Mis compañeros(as) pelean mucho	<i>Same scale</i>
g1_6	A mis compañeros(as) les gusta hacer sufrir a los demás	<i>Same scale</i>
g1_7	La paso mal en la sala de clase	<i>Same scale</i>
g1_8	A mis compañeros(as) les gusta poner sobrenombres	<i>Same scale</i>
<i>Panel B. Cyber-victimization, last 12 months (any item)</i>		
g2_1	¿En los últimos 12 meses has recibido mensajes ofensivos por medio de WhatsApp u otros servicios de mensajería?	1 = Sí; 2 = No; 99 = Prefiero no responder
g2_2	¿En los últimos 12 meses han publicado fotografías o videos vergonzosos de ti, a través de redes sociales?	<i>Same scale</i>
g2_3	¿En los últimos 12 meses has recibido llamadas al celular insultantes o amenazantes?	<i>Same scale</i>
g2_4	¿En los últimos 12 meses has recibido mensajes de correo electrónico insultantes o amenazantes?	<i>Same scale</i>
g2_5	¿En los últimos 12 meses has leído o tenido conocimiento de conversaciones en grupos de chat en las que se expresan frases ofensivas sobre ti?	<i>Same scale</i>
g2_6	¿En los últimos 12 meses has visitado o tenido conocimiento sobre páginas web donde se burlan de ti, o se descarga información personal sin tu permiso?	<i>Same scale</i>
<i>Panel C. Dating violence, among adolescents who ever dated (any item)</i>		
g9_1	Violencia física	0 = No; 1 = Sí
g9_2	Violencia psicológica	<i>Same scale</i>
g9_3	Violencia sexual	<i>Same scale</i>

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. Cyber-victimization is one if any of the listed items is answered yes. The dating-violence items are the options of a multiple-response question on the forms of violence suffered in a relationship; the indicator is one if any is marked.

Table A5**Risk Behaviors: Items and Scales**

	Question, verbatim (1)	Response scale (2)
<i>Panel A. Substance use, last 12 months</i>		
g11	¿Has fumado cigarrillos de tabaco alguna vez en los últimos 12 meses?	1 = Sí; 2 = No
g12_1	Cerveza	<i>Same scale</i>
g12_2	Vino	<i>Same scale</i>
g12_3	Licores solos o mezclados con refrescos (pisco, vodka, ron, gin)	<i>Same scale</i>
g12_4	Cualquier otra bebida que contenga alcohol	<i>Same scale</i>
g13_1	Cannabis (Marihuana, hachís, creppy)	<i>Same scale</i>
g13_2	Cocaína (coca, polvo, nieve, jale)	<i>Same scale</i>
g13_3	Éxtasis, MDMA	<i>Same scale</i>
g13_4	Anfetaminas, metanfetaminas, tusi o speed	<i>Same scale</i>
g13_5	Pasta base (bazuca, pasturri, mono, marciano)	<i>Same scale</i>
g13_6	Opiáceos (heroína, metadona, ketamina, fentanil)	<i>Same scale</i>
g13_7	LSD (ácidos, tripi, alucinógenos)	<i>Same scale</i>
g13_8	Inhalables (pegamentos, disolventes, aerosoles)	<i>Same scale</i>
g13_9	Medicamentos no recetados (clonazepam, u otros tranquilizantes)	<i>Same scale</i>
<i>Panel B. Sexual behavior</i>		
g6	¿Has tenido relaciones sexuales voluntarias? Esto también se conoce como “hacer el amor” o “tener sexo”. Incluye relaciones con o sin penetración.	1 = Sí; 2 = No
g8	En tu última relación sexual voluntaria, ¿tú o tu pareja utilizaron algún método anticonceptivo?	1 = Sí; 2 = No

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. Drank alcohol is one if any of the four beverage items is answered yes; cannabis is item g13_a alone, and other drugs is one if any of items g13_b to g13_i is answered yes. No contraception is one when no method was used at the last voluntary sexual relation, among adolescents who report one.

Table A6
Daily Life: Items and Scales

	Question, verbatim (1)	Response scale (2)
a8_a	Pensando en una semana de lunes a viernes, en promedio ¿a qué hora te despiertas?	Open numeric answer (clock time)
a8_b	Pensando en una semana de lunes a viernes, en promedio ¿a qué hora te duermes?	Open numeric answer (clock time)
a8_e	Pensando en la última semana, ¿acostumbas a quedarte despierto después de acostarte mirando tu celular o Tablet?	1 = Nunca; 2 = Una vez a la semana; 3 = Más de una vez a la semana; 4 = Todos los días
a4	Pensando en un día de lunes a viernes, en promedio ¿Cuánto tiempo al día usas aplicaciones como Instagram, TikTok, WhatsApp, Youtube, Facebook u otras?	1 = Nada; 2 = Menos de 1 hora; 3 = Entre 1 y menos de 2 horas; 4 = Entre 2 y menos de 3 horas; 5 = Entre 3 y menos de 4 horas; 6 = Entre 4 y menos de 5 horas; 7 = Más de 5 horas.
a2	Durante la última semana, ¿cuántos días hiciste deporte o actividad física por al menos 60 minutos?	Open numeric answer (days, 0–7)
a7	Pensando en un día de lunes a viernes, en promedio ¿Cuánto tiempo lees libros, revistas o cómics en algún dispositivo electrónico, por ejemplo: Tablet, celular o computador?	1 = Nada; 2 = Menos de 1 hora; 3 = Entre 1 y menos de 2 horas; 4 = Entre 2 y menos de 3 horas; 5 = Entre 3 y menos de 4 horas; 6 = Entre 4 y menos de 5 horas; 7 = Más de 5 horas.
a10_h	Comida chatarra (completos, hamburguesas, pizzas, papas fritas, etc.)?	Open numeric answer (days, 0–7)
a11	Pensando en una semana de lunes a domingo, ¿cuántas veces al día comes frutas?	Open numeric answer (portions per day)

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. Hours of sleep on weekdays are the survey's own computation from the waking and sleeping times (variable tot_hrs_sueno_sem), kept between 3 and 14. Staying up on the phone, time on social media and time reading are standardized; the days of physical activity and of junk food and the daily portions of fruit enter in their own units.

Table A7
Family and Social Life: Items and Scales

	Question, verbatim (1)	Response scale (2)
<i>Panel A. Parental knowledge of the adolescent's activities (standardized mean of eight items)</i>		
f2_1	¿Sabes en tu casa qué haces durante tu tiempo libre?	1 = Nunca; 2 = Casi nunca; 3 = A veces; 4 = Muchas veces; 5 = Siempre
f2_2	¿Con qué frecuencia saben en tu casa quiénes son tus amigos y/o amigas?	Same scale
f2_3	¿Con qué frecuencia saben en tu casa qué tareas escolares tienes?	Same scale
f2_4	¿Sabes en tu casa en qué gastas tu dinero?	Same scale
f2_5	¿Sabes en tu casa cuándo tienes pruebas, exámenes o trabajos para el colegio/instituto, centro de formación técnica o universidad?	Same scale
f2_6	¿Sabes en tu casa cómo te va en los distintos ramos del colegio/instituto, centro de formación técnica o universidad?	Same scale
f2_7	¿Sabes en tu casa dónde vas cuando sales con tus amigos y/o amigas?	Same scale
f2_8	¿Sabes en tu casa qué haces después del colegio/instituto, centro de formación técnica o universidad?	Same scale
<i>Panel B. Relationships and time with the family</i>		
f5_1	¿Cómo es tu relación con %NOMBRE RESPONS- ABLE PRINCIPAL%?	1 = Muy mala; 2 = Mala; 3 = Más o menos; 4 = Buena; 5 = Muy buena
f7	En la última semana, ¿tú y alguien de tu familia vieron juntos televisión o vieron una película, serie o videos por Internet?	1 = Sí; 2 = No
f8	En la última semana, ¿tú y alguien de tu familia pasaron tiempo juntos en la casa, por ejemplo, jugando o conversando?	Same scale
f9	En la última semana, ¿tú y alguien de tu familia salieron o hicieron alguna actividad juntos fuera de la casa?	Same scale
f10	En la última semana, ¿tú y alguien de tu familia, visitaron juntos a amigos, amigas o familiares?	Same scale
f11	En la última semana, ¿tú y alguien de tu familia, compartieron al menos una comida al día, por ejemplo, desayuno, almuerzo once o cena?	Same scale
f1 _9	Option "Nadie" (nobody) of a multiple-response question on whom the adolescent tells about their problems.	Marked = 1; not marked = 0
<i>Panel C. Social participation</i>		
a1a _99	Option "No participa en ninguna organización o grupo" of a multiple-response question listing sports clubs, religious, artistic, civic and student groups.	Takes part = 1 when not marked

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. Parental knowledge is the mean of items f2_1 to f2_8 when at least six are answered. Activities with the family count the yes answers to items f7 to f11. The relationship with the main caregiver is standardized.

Table A8**Adverse Experiences Reported by the Caregiver: Items and Scales**

	Question, verbatim (1)	Response scale (2)
aces_1	Vio, escuchó o fue víctima de los siguientes hechos de violencia en su vecindario: agresiones, atentados u otros actos violentos	1 = Sí; 2 = No; 88 = No sabe; 99 = No responde
aces_2	Ha presenciado peleas o amenazas entre los integrantes del hogar	<i>Same scale</i>
aces_3	Ha debido enfrentar una separación abrupta y prolongada de su responsable principal o figuras significativas, por ejemplo, muerte o separación.	<i>Same scale</i>
aces_4	Se separó de sus personas responsables por estar en un hogar de acogida, Centro de Mejor Niñez o algún programa residencial	<i>Same scale</i>
aces_5	Ha tenido problemas de vivienda, por ejemplo, vivir en la calle, no tener un lugar estable para vivir, mudarse seguido.	<i>Same scale</i>
aces_6	Ha sido discriminado, por ejemplo, “bullying”, o lo(la) han molestado o excluido(a) por su raza, etnia, identidad de género, orientación sexual, religión, diferencias en el aprendizaje o discapacidades.	<i>Same scale</i>
aces_7	No tuvo suficiente qué comer o hubo riesgo de que la comida se fuera a acabar antes de que se pudiera comprar más	<i>Same scale</i>
aces_8	Vivió o vive con alguien que tiene o tenía algún problema mental (por ejemplo, depresión, esquizofrenia, psicosis, trastorno bipolar, trastorno de ansiedad, entre otros)	<i>Same scale</i>
aces_9	Vivió o vive con alguien que estuvo privado de libertad	<i>Same scale</i>
aces_10	Vivió o vive con personas que tenían un problema de consumo excesivo de alcohol, drogas o medicinas	<i>Same scale</i>

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. The main caregiver answers each item about the adolescent's whole life. The count adds the yes answers when at least eight items are answered.

Table A9
Caregiver Measures I: Items and Scales

	Question, verbatim (1)	Response scale (2)
<i>Panel A. Caregiver depressive symptoms, CES-D, ten items on the last week</i>		
cesd_p1a	Me molestaron cosas que usualmente no me molestan	1 = Casi nunca o ninguna vez (menos ...; 2 = Pocas veces (entre 1 y 2 días); 3 = Varias veces (entre 3 y 4 días); 4 = Casi todo el tiempo (entre 5 y 7...
cesd_p1b	Tuve dificultad para mantener mi atención en lo que hacía	<i>Same scale</i>
cesd_p1c	Me sentí con esperanza sobre mi futuro	<i>Same scale</i>
cesd_p1d	Me sentí deprimido/a	<i>Same scale</i>
cesd_p1e	Sentí que todo lo que hacía me costaba un gran esfuerzo	<i>Same scale</i>
cesd_p1f	Me sentí con miedo	<i>Same scale</i>
cesd_p1g	Dormí mal en la noche (descansé poco)	<i>Same scale</i>
cesd_p1h	Me sentí contento/a	<i>Same scale</i>
cesd_p1i	Me sentí solo/a	<i>Same scale</i>
cesd_p1j	Sentí que no tenía ganas de hacer nada	<i>Same scale</i>
<i>Panel B. Parenting stress and household income</i>		
psi	Parenting Stress Index, short form, answered by the caregiver: parental distress, dysfunctional parent-child interaction and difficult child, twelve items each. The survey releases the scores, not the items.	Total 36–180 and subscales 12–60, standardized
ytotal _pc	Total household income per capita in the month before the interview, computed by the survey from the income module.	Chilean pesos; enters in logs

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. The CES-D score is the survey's raw score (0–30, positive items reversed), standardized.

Table A10**Caregiver Measures II: Parenting Sense of Competence**

	Question, verbatim (1)	Response scale (2)
pscs_p1	Los problemas de cuidar a un(a) adolescente o joven son fáciles de resolver una vez que sabes cómo tus acciones lo pueden afectar	1 = Muy en desacuerdo; 2 = Moderadamente en desacuerdo; 3 = Ni de acuerdo ni en desacuerdo; 4 = Moderadamente de acuerdo; 5 = Muy de acuerdo
pscs_p2	Cumplo mis expectativas respecto al dominio que tengo en el cuidado del adolescente o joven	<i>Same scale</i>
pscs_p3	Yo sería un buen ejemplo a seguir para que una madre o padre aprendan lo necesario para ser buenos padres	<i>Same scale</i>
pscs_p4	Ser cuidador es manejable, y los problemas se resuelven fácilmente	<i>Same scale</i>
pscs_p5	Si alguien puede encontrar la respuesta a lo que le molesta al adolescente o joven, ese/esa soy yo	<i>Same scale</i>
pscs_p6	Un problema difícil de ser cuidador(a) es no saber si estás haciendo un buen trabajo	<i>Same scale</i>
pscs_p7	Teniendo en cuenta el tiempo que he sido un(a) cuidador(a), me siento muy cómodo(a) en este rol	<i>Same scale</i>
pscs_p8	Creo que tengo todas las habilidades necesarias para ser un(a) buen(a) cuidador(a) para el adolescente o joven	<i>Same scale</i>
pscs_p9	A pesar de que ser cuidador(a) puede ser gratificante, me siento frustrado(a) por como es el adolescente o joven en este momento	<i>Same scale</i>
pscs_p10	No sé por qué, pero a veces cuando se supone que debo tener el control, más bien siento que estoy siendo manipulado(a)	<i>Same scale</i>
pscs_p11	Mi padre/madre estaba mejor preparado(a) que yo para ser un(a) buen(a) cuidador(a)	<i>Same scale</i>
pscs_p12	A veces siento como si no estuviera logrando terminar nada	<i>Same scale</i>
pscs_p13	Voy a la cama de la misma manera que me despierto por la mañana, sintiendo que no he logrado mucho	<i>Same scale</i>
pscs_p14	Mis talentos e intereses están en otras áreas, no en ser un/a cuidador/a	<i>Same scale</i>
pscs_p15	Si ser cuidador(a) de un adolescente o joven fuera más interesante, estaría más motivado(a) a hacer un mejor trabajo como cuidador(a)	<i>Same scale</i>
pscs_p16	Ser cuidador/a me pone tenso/a y ansioso/a	<i>Same scale</i>
pscs_p17	Ser un/a buen/a cuidador/a es una recompensa en sí mismo	<i>Same scale</i>

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. Parenting Sense of Competence Scale, answered by the main caregiver; the analysis uses the survey's total score, standardized, with higher values meaning more competence.

Table A11
Family Mental-Health Controls: Items and Scales

	Question, verbatim (1)	Response scale (2)
<i>Panel A. Family history of mental illness, asked about the mother (2012)</i>		
b55o_madre	Depresión. Le mencionaré una lista de enfermedades, ¿me podrí...	1 = sí; 2 = no; 9 = no sabe
b55p_madre	Ansiedad. Le mencionaré una lista de enfermedades, ¿me podría...	<i>Same scale</i>
b55q_madre	Trastorno bipolar o bipolaridad. Le mencionaré una lista de e...	<i>Same scale</i>
b55r_madre	Esquizofrenia. Le mencionaré una lista de enfermedades, ¿me p...	<i>Same scale</i>
b55s_madre	Trastornos de la personalidad. Le mencionaré una lista de enf...	<i>Same scale</i>
b55t_madre	Trastornos de la esfera autista (autismo o síndrome de asperg...	<i>Same scale</i>
<i>Panel B. Number of children of the mother (2012)</i>		
b71	¿Cuántos hijos nacidos vivos tiene la madre biológica del niño (a) seleccio...	Open numeric answer (number of children); special codes: 88 = no responde; 99 = no sabe

Notes: Question wording and response labels are reproduced verbatim in the original Spanish of the questionnaire, from the variable and value labels of the public microdata; wording the microdata truncates at 80 characters is marked with an ellipsis. The same six items are asked about the father (b55o_padre to b55t_padre) and about another relative (b55o_ofam to b55t_ofam). Each of the three indicators of family mental illness is one if any of the six conditions is reported, and the three enter the 2012 controls of the main specification together with the number of children.