

1 The partitioning of gross primary production for young *Eucalyptus tereticornis*
2 trees under experimental warming and altered water availability

3 John E. Drake^{1,2*}, Mark G. Tjoelker¹, Michael J. Aspinwall^{1,3}, Peter B. Reich^{1,4}, Sebastian
4 Pfautsch^{1,5}, Craig V. M. Barton¹

5 ¹Hawkesbury Institute for the Environment, Western Sydney University, Locked Bag 1797,
6 Penrith NSW 2751, Australia. ORCHID ID:

7 ²Forest and Natural Resources Management, SUNY-ESF, 1 Forestry Drive, Syracuse, NY,
8 13210 USA.

9 ³Department of Biology, University of North Florida, 1 UNF Drive, Jacksonville, FL, 32224
10 USA.

11 ⁴Department of Forest Resources, University of Minnesota, 1530 Cleveland Ave N., St Paul,
12 MN, 55108 USA.

13 ⁵School of Social Science and Psychology (Urban Studies), Western Sydney University, Locked
14 Bag 1797, Penrith, NSW 2751, Australia.

15 *Corresponding author. Email: jedrake@esf.edu, telephone: 415-470-6574

16 ORCHID IDs

17 John Drake: <https://orcid.org/0000-0003-4274-4780>

18 Mark Tjoelker: <https://orcid.org/0000-0003-4607-5238>

19 Michael Aspinwall: <https://orcid.org/0000-0003-0199-2972>

20 Peter Reich: <https://orcid.org/0000-0003-4424-662X>

21 Sebastian Pfautsch: <https://orcid.org/0000-0002-4390-4195>

22 Craig Barton: NA

23 Heading for social media: Carbon allocation in trees measured in unique whole-tree chambers.

24 Article accepted 20 Nov 2018

25 Word counts:

26 Main text (Intr-Ack): 6999

27 Summary: 199

28 Introduction: 1349

29 Materials and Methods: 2534

30 Results: 1327

31 Discussion: 1670

32 Acknowledgements: 119

33 8 figures, 1 tables, and supporting information.

Summary

- The allocation of carbon (C) is an important component of tree physiology that influences growth and ecosystem C storage. Allocation is challenging to measure, and its sensitivity to environmental changes such as warming and altered water availability is uncertain.
- We exposed young *Eucalyptus tereticornis* trees to +3°C warming and elimination of summer precipitation in the field using whole-tree chambers. We calculated C allocation terms using detailed measurements of growth and continuous whole-crown CO₂ and H₂O exchange measurements.
- Trees grew from small saplings to nearly 9-m-height during this 15 month experiment. Warming accelerated growth and leaf area development, and increased the partitioning of Gross Primary Production (GPP) to aboveground respiration and growth, while decreasing partitioning belowground. Eliminating summer precipitation reduced C gain and growth, but did not impact GPP partitioning. Trees utilized deep soil water and avoided strongly negative water potentials.
- Warming increased growth respiration, but maintenance respiration acclimated homeostatically. The increasing growth in the warmed treatment resulted in higher rates of respiration, even with complete acclimation of maintenance respiration. Warming-induced stimulations of tree growth likely involve increased C allocation aboveground, particularly to leaf area development, while reduced water availability may not stimulate allocation to roots.

Key words. Allocation, respiration, partitioning, warming, drought, climate change

56 **Introduction**

57 The carbon (C) economy of trees and forests depends not only on the amount of C fixed via
58 photosynthesis, but how that fixed C is used. Ecosystem C storage is affected by the allocation of C to
59 long-lived C pools such as wood relative to C allocation to pools with higher turnover rates such as fine
60 roots (DeLucia *et al.*, 2005). C allocation also affects the acquisition of light, nutrients, and water, which
61 influences ecosystem C cycling and tree survival (Litton *et al.*, 2007; Epron *et al.*, 2012; Franklin *et al.*,
62 2012; De Kauwe *et al.*, 2014; Trugman *et al.*, 2018). The allocation of C belowground affects soil C and
63 nutrient cycling (Högberg *et al.*, 2001; Epron *et al.*, 2012) in part because belowground C allocation can
64 affect soil organic matter decomposition and the acquisition of limiting nutrients by trees (Drake *et al.*,
65 2011; Finzi *et al.*, 2015). The importance of C allocation and the relative difficulty of its study contribute
66 to its role as an important unknown for modeling the biogeochemistry of ecosystems (Roux *et al.*, 2001;
67 Franklin *et al.*, 2012; Dietze *et al.*, 2014; De Kauwe *et al.*, 2014).

68 Terminology regarding allocation has been a source of some confusion. Here, we follow Litton *et al.*
69 *et al.* (2007) and use ‘allocation’ as a term of broad definition encompassing three specific aspects of study:
70 (1) *ratios* of biomass pool sizes, (2) *fluxes* of C to a given component, and (3) *partitioning*, the C flux to a
71 given component as a fraction of GPP. These areas of study are similar, but not equivalent. For example,
72 old and large trees have a large wood mass fraction relative to small young trees (Poorter *et al.*, 2015), but
73 this reflects the low turnover of woody tissues relative to leaves, not a higher partitioning of GPP to wood
74 in old trees (Duursma & Falster, 2016). Thus it is often inappropriate to infer C partitioning from biomass
75 ratios (Reich, 2002; Litton *et al.*, 2007). Surprisingly, partitioning of photosynthate is relatively poorly
76 understood despite its direct relevance to ecosystem models (Epron *et al.*, 2012; Franklin *et al.*, 2012; De
77 Kauwe *et al.*, 2014).

78 Several schemes have been used to conceptualize and model C allocation. The simplest approach
79 is to assume that trees partition a constant fraction of fixed C to each use (e.g., growth, respiration). This
80 is supported by linear relationships between production terms in some systems (Gower *et al.*, 2001).
81 However, fixed allocation schemes cannot capture ontogenetic effects (Poorter *et al.*, 2015; Duursma &
82 Falster, 2016) or dynamic temporal responses (De Kauwe *et al.*, 2014; Doughty *et al.*, 2014). Another
83 approach is to assume a functional balance between tree organs via allometric relationships, Huber values,
84 or root to leaf mass fractions (Landsberg & Waring, 1997; Mäkelä *et al.*, 2008; Feng *et al.*, 2012). Finally,
85 there is the concept that trees increase C partitioning towards the acquisition of the primary limiting
86 resource (McMurtrie & Dewar, 2013). This approach appears sensible and has been implemented in
87 several models (e.g., Running & Gower, 1991; Friedlingstein *et al.*, 1999), but direct evidence supporting

88 this concept is scarce, given the challenges involved in measuring allocation (Poorter & Sack, 2012;
89 Poorter *et al.*, 2015). However, optimization approaches have been used to constrain dynamic allocation
90 schemes with some success (Franklin *et al.*, 2012; McMurtrie & Dewar, 2013).

91 Temperature is a fundamental aspect of climate that affects many aspects of tree physiology,
92 including photosynthesis, respiration, and growth (Way & Oren, 2010; Lu *et al.*, 2013). Several lines of
93 evidence suggest that warming may increase tree C allocation aboveground at the expense of C allocation
94 belowground. Experimental warming of forest soil increased aboveground biomass while reducing or not
95 affecting belowground biomass (Strömberg & Linder, 2002; Melillo *et al.*, 2011), which has been
96 attributed to a warming-induced increase in soil N availability (Melillo *et al.*, 2002). A recent ^{13}C -CO₂
97 labeling study indicated that warming increased allocation aboveground and reduced C allocation
98 belowground in beech saplings via a direct effect on tree physiology, without an altered soil N cycle
99 (Blessing *et al.*, 2015). Also, environmental gradients in mean annual temperature are strongly correlated
100 with the distribution of biomass; forests have a lower root mass fraction (i.e., the proportion of live forest
101 biomass contained in roots) in warm climates than in cold climates (Reich *et al.*, 2014). Meta-analyses of
102 warming experiments generally find an increase in aboveground plant growth (Rustad *et al.*, 2001) that is
103 slightly larger than the increase in belowground plant growth (Lu *et al.*, 2013), although such experiments
104 have exclusively involved small stature vegetation given the logistical challenges of warming tall forests.
105 Thus, prior research suggests a shift in C allocation aboveground with experimental warming, although
106 direct tests in the field with large trees have not yet been performed.

107 Water availability also impacts tree growth and physiology (Mencuccini, 2003; Nemani *et al.*,
108 2003; Farooq *et al.*, 2009; Martínez-Vilalta *et al.*, 2009; Yang *et al.*, 2018) and the effects of drought are
109 of particular concern (Burke *et al.*, 2006; Sillmann *et al.*, 2013; Jump *et al.*, 2017). While it appears
110 sensible that trees would increase C allocation to roots in dry regions or during drought periods (Poorter
111 *et al.*, 2012), there is limited support for this idea. Reich *et al.* (2014) found no correlation between root
112 mass fraction and aridity across a global dataset of >6,200 forests. Additionally, Amazonian forests
113 responded to droughts in 2005 and 2010 with a shift away from fine-root growth and increased C
114 partitioning to aboveground growth and respiration, particularly in the year following the drought
115 (Doughty *et al.*, 2014, 2015). However, drought has been observed to increase root mass fractions for
116 small plants grown in artificial conditions (Reich, 2002; Poorter *et al.*, 2012), and increased C allocation
117 belowground under drought is consistent with some ^{13}C -CO₂ labeling studies (Hommel *et al.*, 2016), but
118 not others (Hartmann *et al.*, 2015). While the simple expectation of increased allocation to roots during
119 drought is appealing, C allocation responses to drought are likely more complex and merit further study.

120 Interactions between temperature and drought effects may also be important for tree C allocation.
121 Warmer temperatures may exacerbate tree H₂O loss during drought and increase mortality risk (Allen *et*
122 *al.*, 2015). If warmer temperatures reduce C allocation belowground, then the ability of trees to acquire
123 soil water may also be impaired. However, an open top chamber experiment with young oak saplings
124 found no interaction between experimental warming and drought on tree transpiration or biomass (Kuster
125 *et al.*, 2013). Other studies have found limited evidence for interactive effects on plant growth rates (e.g.,
126 Edler *et al.*, 2015; Taeger *et al.*, 2015), although there are also exceptions (Munir *et al.*, 2015; Leon-
127 Sanchez *et al.*, 2016). A six-year warming and precipitation redistribution experiment with two tree
128 species found complex growth responses (Volder *et al.*, 2013) with a strongly interactive effect on the
129 relative growth rate of *Quercus stellata* monocultures. Thus, it is challenging to generalize how the
130 interactive effects of drought and warming affect tree physiology and growth.

131 To address these knowledge gaps concerning, temperature, water availability, and C allocation,
132 we studied C allocation in young eucalypt trees as they grew from roughly 1 to 9 meters in height in an
133 experiment that manipulated both temperature and water availability. ~~We used whole-tree chambers in the~~
134 field in southeastern Australia to grow *Eucalyptus tereticornis* trees under experimental warming of +3 °C
135 for more than one year, crossed with a summer drought for three months. We continuously measured
136 whole-crown CO₂ and H₂O exchange and measured aboveground biomass production every 2 weeks.
137 From these intensive measurements, we derived GPP, aboveground net primary production (NPP_a),
138 aboveground autotrophic respiration (R_a) and the residual C that must have been partitioned belowground
139 for each fortnightly interval. We use these data to test the predictions that warming decreases C
140 partitioning belowground, while drought increases C partitioning belowground.

Commented [JED1]: A sentence was removed here, at the editor's request.

Materials and Methods

Site and experiment

We implemented a warming and drought experiment using 12 whole-tree chambers (WTCs) in Richmond, New South Wales (Australia; 33°36'40"S, 150°44'26.5"E). The WTCs were large cylindrical structures topped with a cone (3.25 m in diameter, 9 m in height, volume of ~53 m³) that enclosed a single tree rooted in soil. The WTCs controlled atmospheric CO₂ concentration, air temperature (T_{air}), relative humidity (RH), and irrigation while continuously measuring the net exchange of CO₂ and H₂O between entire tree crowns and the atmosphere (Barton *et al.*, 2010; Duursma *et al.*, 2011; Barton *et al.*, 2012; Duursma *et al.*, 2014; Drake *et al.*, 2016b; Aspinwall *et al.*, 2016).

The roots of each tree were compartmentalized with a barrier extending vertically belowground to 100-cm-depth. A cemented layer of manganese nodules and clay was present at 90–100 cm depth, providing a natural horizontal barrier at the bottom of the rooting volume. Thus, the rooting volume of each tree was isolated from surrounding trees. However, some trees extended roots through this layer and acquired deep soil water in a previous experiment (Duursma *et al.*, 2011). Soil was collected from an adjacent paddock and placed in the chambers in two layers (0–25 cm and from 25 cm to the hard layer) on 10 July 2012. Soils at the site were an alluvial formation of low-fertility sandy loam (Clarendon sand).

Nursery seedlings of a local provenance of *Eucalyptus tereticornis* Sm. were established in 25 L pots inside the WTCs using the same soil. *Eucalyptus tereticornis* was chosen because it is a widespread and abundant tree across eastern Australia (Drake *et al.*, 2015). Six potted trees were placed in each chamber on 5 December 2012; a single tree was selected based on size similarity within each treatment and planted in the chamber center on 12 March 2013. Trees assigned to the ambient and warmed temperature treatments had equivalent height and basal diameter when potted seedlings were placed into the WTCs in December 2012 (heights of $41.5 \pm \text{SE of } 0.8$ and 40.2 ± 1.8 cm; diameters of 2.4 ± 0.1 and 2.5 ± 0.1 mm in ambient and warmed treatment, respectively). Thus, the experiment began with small seedlings, but subsequent growth over the 15-month study period was rapid and trees quickly grew into larger size classes.

Six chambers tracked ambient T_{air} and six chambers tracked ambient $T_{\text{air}} + 3$ °C warming ($n = 6$; 'ambient' and 'warmed', respectively); treatments started on 12 December 2012 (Drake *et al.* 2016b; Aspinwall *et al.* 2016). The average warming achieved was +2.9 °C (\pm sd of 0.3 across 466 days) for T_{air} , +2.2 °C (\pm sd of 0.4) for soil temperature at 5-cm-depth, +2.0 °C (\pm sd of 0.4) for soil temperature at 20-cm-depth, and +1.4 °C (\pm sd of 0.2) for soil temperature at 50-cm-depth. Trees were irrigated equally every 15 d with half the mean monthly rainfall, which we consider the control treatment. A water exclusion treatment was applied to half of the trees on 12 February 2014, resulting in a 2x2 factorial

design between the experimental treatments of warming and drought ($n = 3$; abbreviated A-Con, A-Dry, W-Con, and W-Dry hereafter). Trees assigned to the drought treatment received no irrigation from 12 February 2014 through 5 May 2014, representing a summer drought of nearly three months. A rainfall record from 1881–2014 from this area (Australian Bureau of Meteorology station 67021, station is <2 km from site) indicates that only three years (~2% of the record) had <20 mm precipitation during the months of the drought treatment, and 80% of the years had >100 mm precipitation. This suggests that our drought was relatively strong in terms of the surface water addition during these months.

Plant water status and soil water content

Predawn leaf water potentials (Ψ_{L-PD}) were measured monthly prior to the drought and every one to two weeks during the drought treatment. Three leaves were measured per tree on each date using a Scholander-type pressure chamber (1505D-EXP; PMS Instrument Company, OR, USA). Leaves were placed in sealed and humidified plastic bags, placed in a dark cool box, and measured within one hour of collection in a nearby laboratory.

Soil volumetric water content was measured by three sensors in each chamber (CS650 time-domain reflectometers; Campbell Scientific, Logan, UT, USA). Sensors were installed horizontally at three depths: in the surface soil (10-cm-depth), at 30-cm-depth, and just above the hard layer of cemented manganese (~100-cm-depth). Soil temperature was measured with thermocouples at 5, 20, and 50-cm within the center of each chamber.

A single neutron probe tube per chamber (503DR, Hydroprobe, Instrotek, NC, USA) was used to measure soil water content to a depth of 425 cm (at 25 or 50 cm steps) approximately every two weeks (Duursma *et al.*, 2011). Note that high neutron probe counts in deep soil (150–400 cm depth) partially reflect a change in soil texture towards a higher clay content.

Whole tree crown flux measurements

An automated system measured the net exchange of CO_2 and H_2O between each crown and its chamber airspace (Barton *et al.*, 2010). Measurements began on 13 September 2013 when suspended plastic floors were sealed around the stem of each tree at ~45 cm height, when the trees were ~3 m tall. Flux measurements finished on 26 May 2014, when the trees were nearly 9 m tall. We report >70,000 hourly flux observations aggregated into >3000 daily sums across 12 trees.

We partitioned the net CO_2 fluxes into the components of GPP and R_a using an analytical technique common to eddy-covariance research (Reichstein *et al.*, 2005). We used direct measurements of whole-crown R_a and its temperature dependence at night to predict R_a for each hourly measurement as a function of T_{air} . For daylight hours, we then calculated GPP as the sum of the measured net CO_2 flux and the predicted R_a given the measured T_{air} . We assumed GPP was zero when PPFD = 0; in such conditions,

the measured net C flux was used as the measure of R_a . Note that the chamber airtspaces were continuously well-mixed and R_a fluxes were directly measured at night, avoiding some of the issues inherent in eddy covariance partitioning. The underlying flux data and the partitioning approach were published previously (Drake *et al.*, 2016ab).

Final harvest

The dry mass of all trees was measured destructively at the end of the experiment (26 May 2014), 15 months after seedlings were planted. At this time, the trees had heights of 8.8 ± 0.14 meters and diameters of 6.6 ± 0.2 cm. Total tree dry mass was measured as the sum of five components: leaves, branches, stem, coarse roots, and fine roots.

The crown of each tree was divided into three equal heights. All branches were cut flush to the stem and all leaves were separated from branches. A random subsample of 100 leaves per layer was measured for total leaf area (LI-3100C leaf area meter, LiCor, Lincoln, NE, USA), dry mass, and specific leaf area (SLA). The stem was cut into three segments and a single 1-cm-thick cross-section (cookie) was sampled for bark depth, wood density, and bark density at the stem base, between the first and second layers, and between the second and third crown layers. Bark and wood density was measured on cookie subsamples (Thomas *et al.*, 2007). Wood and bark densities were similar (0.44 and 0.37 g cm³ for wood and bark, respectively). Bark depth increased with stem diameter ($\log_{10}(\text{bark depth, mm}) = -1.48 + 1.23 \times \log_{10}(\text{diameter, cm})$, $P < 0.001$, $r^2 = 0.92$) while wood and bark density decreased with stem diameter (wood density = $0.50 - 0.001 \times \text{diameter}$, $P = 0.007$, $r^2 = 0.17$; bark density = $0.45 - 0.001 \times \text{diameter}$, $P < 0.001$, $r^2 = 0.48$; densities in g cm³, diameter in cm). The reduction in density with increasing stem diameter reflected increasing densities in the upper layers of the tree crowns. Warming and drought treatments did not alter these relationships (ANCOVA, $P > 0.05$). Total stem, branch, and leaf mass were measured directly after drying at 70 °C; some samples required >2 weeks of drying to reach a stable dry mass.

Fine roots were measured using soil cores. The soil surface area was divided into four equal quadrants and two 50-mm-diameter cores were taken within each quadrat on 29 May 2014, just after the crown harvest. Cores were separated into two depths: (1) 0–25 cm and (2) from 25 cm to the hard layer, which varied from 70 to 100 cm depth. Samples within each quadrat and depth category were composited (eight samples per chamber). Fine roots were isolated by washing samples through 2-mm and then 1-mm brass sieves; fine roots were defined as all roots < 2-mm-diameter. Fine root dry mass was measured after drying at 70 °C. Total fine root dry mass was calculated as the product of fine root density in the soil (g m⁻³) and soil volume (m³) in each layer.

Coarse roots were destructively harvested by fully excavating the soil volume of each chamber. Soil was shoveled out of the chamber onto a conveyor belt that transported the soil to a series of 5-mm

steel sieves. Roots were collected by hand, washed, sorted into two size categories (2-10 mm, > 10 mm diameter), and weighed after drying at 70 °C. During the excavation, several small roots (~1 cm diameter) were observed to have penetrated through the hard layer and into deeper soil (Drake, *personal observation*).

Growth measurements

Aboveground biomass was estimated every two weeks for each tree as the sum of leaf, branch, wood, and bark mass. The paragraphs below describe the methodologies for each component in detail. Aboveground net primary production (NPP_a) was estimated as the fortnightly difference in aboveground biomass plus fortnightly litterfall, assuming a constant biomass C fraction of 0.5 of dry mass. Allocation to reproduction was not explicitly measured, but can be considered zero as these trees did not produce any reproductive structures.

Tree height and stem volume were measured fortnightly; diameter was measured at 30-cm-intervals along each tree stem from a basal height of 15-cm (prior to floor installation) or 65-cm (after floor installation) to the tree apex. The volume of stem wood and bark was estimated for each stem segment as the frustum of a cone, corrected for bark depth (*see above*). Wood and bark mass were calculated as the product of volume and density. This approach assumes that the wood and bark density did not change through time in this experiment.

An allometric relationship was developed to predict branch wood mass from branch diameter. Four branches from each experimental tree were destructively sampled near the end of the study (13 May 2014 and 22 May 2014); branch mass was strongly correlated with branch diameter ($\log_{10}(\text{branch mass}) = -1.299 + 2.722 \times \log_{10}(\text{branch diameter})$, $P < 0.001$, $r^2 = 0.91$, branch mass in g, branch diameter in mm, $n = 48$ branches). This allometry did not differ between treatments (ANCOVA, $P > 0.1$) and was used to predict total branch mass on three dates when the diameter of all branches was measured (24 Oct 2013, 15 Jan 2014, and 22 May 2014). Total branch mass and stem volume were strongly correlated in a chamber specific manner (log-log ANCOVA, $P < 0.001$, $r^2 = 0.95$), which was used to estimate branch mass as a function of stem volume.

Standing leaf area and leaf mass production were estimated as previously at this site (Barton *et al.*, 2012; Drake *et al.*, 2016b). Standing leaf area was measured for each tree by counting all the leaves and multiplying by a tree-specific mean leaf size measured across the crown of each tree with a handheld leaf area meter (LI-3000; $n = 86$ to 102 leaves per tree). These measurements were performed prior to chamber floor installation (9 Sept 2013) and at the beginning of the drought treatment (10 Feb 2014). A third direct measurement of standing leaf area was calculated from the final harvest data (26 May 2014) by multiplying total crown leaf dry mass by SLA weighted by the leaf dry mass in each layer. Litterfall was collected, dried, and weighed fortnightly for each tree, although relatively few leaves fell as litter

(~5% of the total leaf mass). Total tree leaf mass was estimated for each set of fortnightly size measurements by dividing leaf area by the crown-weighted SLA measured at harvest.

Calculating C partitioning

A major goal of this study was to calculate the partitioning of photosynthetically fixed C into components for each fortnightly interval. We quantified GPP, NPP_a , and R_a separately, as described above. We calculated the residual between GPP and the sum of NPP_a and R_a :

$$GPP = NPP_a + R_a + \text{residual} \quad (\text{eq. 1})$$

The residual term is a mass-balance calculation of all C put belowground to root production, respiration, and exudation, but this term is also affected by measurement error in GPP, NPP_a , and R_a . We calculated the partitioning of GPP directly for each fortnightly interval as NPP_a/GPP , R_a/GPP , and residual/GPP.

Growth and maintenance R_a

Given the evidence for thermal acclimation of tissue-specific respiration rates to experimental warming in this experiment (Drake *et al.*, 2016b; Aspinwall *et al.*, 2016), we investigated growth and maintenance respiration as drivers of R_a (McCree, 1970; Tjoelker *et al.*, 1999; Amthor, 2000; Adu-Bredu & Hagihara, 2003). Separating growth and maintenance components of R_a is not possible in all C-cycle studies, but we were able to couple direct measures of tree growth with continuous measurements of integrated whole-tree above ground respiration to enable quantification of growth and maintenance components. We evaluated the relationship between R_a per unit tree C and relative growth rate (RGR); the slope reflects the growth component of R_a , while the y-intercept reflects the maintenance component of R_a . If R_a does not acclimate to warming, we expect the warmed treatment to have a higher y-intercept than the ambient treatment. If R_a acclimates homeostatically, we expect the ambient and warmed treatments to have equivalent intercepts. We also directly estimated coefficients associated with growth and maintenance components of R_a (Amthor, 2000);

$$R_a = R_g + R_m = g_r G + m_r W \quad (\text{eq. 2})$$

where R_g is the growth respiration rate (gC d^{-1}), R_m is the maintenance respiration rate (gC d^{-1}), G is biomass growth (gC d^{-1}), W is the standing biomass weight (gC), g_r is the growth respiration coefficient ($\text{gC respired per gC growth}$), and m_r is the maintenance respiration coefficient ($\text{gC respired per gC standing biomass d}^{-1}$).

Data analysis

Data were analyzed following a completely randomized design with the single treatment of warming ($n = 6$ for 6 months, then $n = 3$ for the drought period). Longitudinal analyses were performed using the 'lme'

306 function within the ‘NLME’ R package with a random tree effect and fixed effects of date, temperature
307 treatment, and water treatment. Treatment means were estimated after adjustment for other terms in the
308 model (i.e. least square means, or LS means) with the ‘LSMEANS’ package in R v.3.2.2 (R Development
309 Core Team, 2012; Pinheiro et al., 2013). Analyses were evaluated to test assumptions of residual
310 normality and homoscedasticity; transformations were often necessary. Datasets that were not
311 longitudinal were analyzed as a 2x2 ANOVA using the ‘lm’ function in R. Equation 2 was fit using the
312 ‘NLME’ R package with a random tree effect.

Results

Growth

Experimental warming increased the rates of diameter and height growth (Fig. 1a,b), particularly during the Austral winter and spring. Trees in the warmed treatment were larger than trees in the ambient treatment when the CO₂ and H₂O flux measurements began (13 Sept 2013; vertical dashed line in Fig. 1). On that date, warming had increased diameter by 21% ($P < 0.01$; Fig. 1a), height by 19% ($P < 0.01$; Fig. 1b), total leaf area by 53% ($P < 0.01$; Fig. 1c), and stem volume by 79% ($P < 0.01$; Fig. 1d). During the warm summer, the diameter and height of the ambient and warmed treatment trees converged (Fig. 1a,b), but total stem volume continued to be larger in the warmed treatment (Fig. 1d). This effect was driven by a difference in stem taper- the warmed trees had wider stem diameters throughout the crown than the ambient trees (*not shown*). The drought treatment reduced tree diameter but not height growth (Fig. 1a,b), modestly reduced total leaf area (Fig. 1c), and reduced stem volume increment (Fig. 1d). Notably, there was no interactive effect of warming and drought on growth (e.g., $P > 0.4$ for volume increment).

CO₂ and H₂O fluxes

Experimental warming increased photosynthetic C uptake and H₂O loss via transpiration early in the experiment (Fig. 2a,c). This was expected, given the strong increase in tree growth and total leaf area with experimental warming during this period (Fig. 1). However, the rates of C uptake and H₂O loss converged between the ambient and warmed treatments during the summer (January; Fig. 2a,c), despite the fact that the warmed trees were larger and had more leaf area. This may have been influenced by warming-induced reductions in photosynthetic rates per unit leaf area (Drake *et al.*, 2016b).

We imposed an experimental drought in which all surface irrigation was withheld from trees in the dry treatments for nearly three months. Total C uptake during the drought period was reduced 25% while total H₂O loss was reduced 32% (Fig. 2b,c; main effects of drought, $P < 0.01$; no interaction with warming, $P > 0.5$). Thus, the drought strongly and significantly reduced whole-crown fluxes of C uptake and H₂O loss. On the other hand, these fluxes were maintained at moderate values during the drought, despite the complete lack of water addition.

Final harvest

The final biomass did not significantly differ between the warming or drought treatments (Fig. 3a). The lack of difference in final mass between the ambient and warmed treatments may have arisen from the convergence of tree diameter and height across treatments (Fig. 1a,b). The difference in stem volume between ambient and warmed trees (Fig. 1d) was apparent in the harvest biomass (Fig. 3a), but was not statistically significant at this level of replication ($n = 3$; $P > 0.05$). The only biomass component that was affected by the experimental treatments at harvest was fine root biomass, for which there was a significant

interaction between warming and drought ($P < 0.05$). The A-Dry trees had higher fine root biomass than the A-Con trees, while the W-Dry trees had slightly lower fine root biomass than the W-Con trees (Fig. 3a). This interaction was also present in the tree root mass ratios; experimental drought increased the root mass ratio, but only in the ambient temperature treatment ($P < 0.05$; Fig. 3b).

Plant and soil water status

The drought reduced soil volumetric water content from 10-100 cm depth to values approaching $0.05 \text{ m}^3 \text{ m}^{-3}$ (Fig. 4a-c). Pre-dawn leaf water potentials ($\Psi_{\text{L-PD}}$) were reduced in the dry treatments relative to the control treatments (Fig. 4d; $P < 0.01$). However, this effect was modest; $\Psi_{\text{L-PD}}$ was -0.29 ± 0.02 in the control and -0.48 ± 0.05 in the dry treatments. Thus, the drought trees had moderate $\Psi_{\text{L-PD}}$ (Fig. 4d) and moderate rates of transpiration (Fig. 2c) despite extremely dry surface soils.

Trees likely utilized deep soil water during the drought treatment. We observed a few roots of approximately 1-cm-diameter penetrating through the cemented manganese layer at ~ 100 cm depth during the complete soil excavation (JE Drake, *personal observation*). Neutron probe measurements down to 400-cm-depth indicated that soil water was removed from the profile in the dry treatment chambers during the drought, particularly from 50- to 200-cm-depth (Fig. S1). Thus, trees in the dry treatments likely transpired deep soil water during the summer drought, consistent with a previous drought study of *Eucalyptus saligna* at this site (Duursma *et al.* 2011).

Fluxes of GPP, NPP_a, R_a and allocation belowground

We derived gross primary production (GPP) and its partitioning to aboveground net primary production (NPP_a), aboveground autotrophic respiration (R_a), and the residual, which we attribute to C allocation belowground as well as measurement error.

GPP was increased by experimental warming early in the experiment ($+22\%$, $P < 0.01$), but GPP between ambient and warmed treatments converged beginning in mid-summer (late January; Fig. 5a). The drought treatment reduced GPP in both temperature treatments (-15% , $P < 0.01$). These results follow the net C flux measurements (Fig. 2a,c). The response of NPP_a (Fig. 5b) closely followed the results for GPP, with a warming effect early in the experiment ($+36\%$, $P < 0.01$) and a reduction with drought in both temperature treatments (-25% , $P < 0.01$). The response of R_a (Fig. 5c) also followed GPP, with a stimulation by warming early in the experiment ($+39\%$, $P < 0.01$) and a modest reduction with drought that was equivalent across temperature treatments (-13% , $P < 0.05$). The allocation of C belowground, as measured by the residual, was decreased by experimental warming throughout the experiment (-11% , $P < 0.05$) and was unchanged by the drought treatment ($+3\%$, $P > 0.1$; Fig. 5d).

GPP partitioning

Given these flux measurements, we derive the partitioning of GPP into three components; NPP_a/GPP , R_a/GPP , and $\text{Residual}/\text{GPP}$ (Figs. 6-7). Warming increased NPP_a/GPP in a way that was stronger early in the experiment (+11%, $P = 0.01$) relative to the entire experiment (+3%; $P > 0.1$; Fig. 6ab). Similarly, warming increased R_a/GPP (+12%; $P < 0.1$) but decreased $\text{Residual}/\text{GPP}$ (-15%; $P < 0.05$) prior to the drought (Figs 6-7). Thus, experimental warming increased the partitioning of GPP to aboveground components (Fig. 7a-b) and decreased partitioning belowground (Fig. 7c). The experimental drought had weak effects on partitioning, none of which were statistically significant ($P > 0.1$).

Growth and maintenance respiration

We combined growth and respiratory measurements to infer changes in respiratory C efflux attributable to growth versus maintenance respiration (Amthor, 2000). There was a strong and linear relationship between R_a per unit aboveground tree C and relative growth rate (RGR; Fig. 8). Neither the slope nor the intercept of this relationship were affected by experimental treatments (all $P > 0.1$). Thus, we present a common relationship across all measurements. The y-intercept of this relationship was positive (mean of 0.0213, 95% CI of 0.0157 to 0.0268), indicating significant R_a in the absence of aboveground growth, reflecting maintenance respiration. The lack of a warming effect on this y-intercept is consistent with respiratory temperature acclimation; trees in the ambient and warmed treatments expended similar amounts of C on maintenance respiration, despite the increased temperature in the warmed treatment. The slope of the relationship (Fig. 8) was strongly positive (mean of 0.0059, 95% CI of 0.0053 to 0.0065), indicating that much of the R_a observed at the crown scale was attributable to construction respiration. Observations during the drought period followed the general relationship, with lower values on both axes (Fig. 8). Thus, the experimental drought reduced R_a primarily via a reduction in growth respiration.

We also directly estimated coefficients for growth and maintenance respiration by fitting equation 2 to the fortnightly dataset of standing biomass, growth rate, and respiration. We estimate the growth respiration to consume approximately 0.3 gC per gram of biomass C produced, and maintenance respiration to consume approximately 0.015 gC per g of standing biomass C per day (Table 1). These coefficients did not differ across the ambient and warmed treatment ($P > 0.3$).

404 **Discussion**

405 *Summary*

406 We studied the experimental effects of warming and drought on the C allocation of *Eucalyptus*
407 *tereticornis* trees using a combination of growth and whole-crown flux measurements. This was the first
408 study to directly measure allocation changes of relatively large trees (as large as 9 meters tall) in an
409 ecologically-relevant field setting with detailed and continuous measurements. Although much smaller
410 than mature trees, the individuals in our experiment were larger than those used in the vast majority of
411 manipulative field experiments involving climate warming. Experimental warming increased the
412 proportion of GPP that was allocated to aboveground uses and decreased the proportion of GPP that was
413 allocated belowground. This was consistent with a reduced root mass fraction in the warmed treatments at
414 the final harvest. The experimental drought reduced CO₂ and H₂O fluxes but did not affect the allocation
415 of C, perhaps because tree access to deep soil water prevented them from experiencing strongly negative
416 water potentials. There were no interactions between warming and drought on C partitioning terms, so we
417 discuss the impacts of warming and drought separately.

418 *Effects of experimental warming on C allocation*

419 Experimental warming strongly affected several aspects of tree C allocation. Warming increased the
420 fractional partitioning of GPP to aboveground uses, including growth and respiration, at the expense of C
421 partitioning belowground. This observation is consistent with some soil warming experiments (e.g.,
422 Melillo *et al.*, 2002, 2011) that attributed this effect to a warming-induced stimulation of soil nutrient
423 availability. However, it is also possible that experimental warming directly stimulated the activity of
424 meristems aboveground, such that a smaller remainder of fixed C was available for transport and use
425 belowground. Such a mechanism would imply an aboveground priority in tree C allocation, consistent
426 with previous work on forest C budgets and elevated atmospheric CO₂ treatments (Palmroth *et al.*, 2006).
427 This mechanism also makes sense given the structural arrangement of tree phloem, as aboveground
428 tissues have the opportunity to remove sucrose from the phloem before belowground tissues (Lemoine *et*
429 *al.*, 2013; Furze *et al.*, 2018). The mechanisms regarding soil nutrient availability and aboveground
430 metabolic activity are not mutually exclusive. For example, enhanced N supply from soil N
431 mineralization may have enabled the increased aboveground metabolism in the warmed treatment, which
432 may have resulted in the larger consumption of GPP aboveground in the warmed relative to the ambient
433 treatment.

434 We also acknowledge that warming may have influenced allocation indirectly via ontogenetic
435 drift. A meta-analysis documented a decline in root to shoot ratios as trees grew larger, possibly reflecting

an ontogenetic effect on belowground allocation (Mokany *et al.*, 2006), although we hesitate to infer allocation directly from root to shoot ratios (Reich, 2002; Litton *et al.*, 2007). In this experiment, the allocation terms (e.g., NPP_a/GPP , R_a/GPP , and $\text{residual}/\text{GPP}$) were not significantly correlated with any metric of tree size, suggesting that the warming effect is unlikely to reflect ontogeny.

We previously demonstrated that aboveground autotrophic respiration acclimated nearly homeostatically to experimental warming in this experiment, both at the leaf-scale (Aspinwall *et al.*, 2016) and at the whole-crown scale (Drake *et al.*, 2016b). As such, the demonstration that warming increased R_a (Fig. 5) may appear contradictory. We emphasize that our previous presentations of autotrophic respiration were expressed per unit leaf area (Drake *et al.*, 2016b; Aspinwall *et al.*, 2016), while the current study shows the total fluxes per tree (Fig. 5c). Experimental warming increased R_a primarily by increasing growth and tree size early in the experiment. Furthermore, the common relationship between relative growth rate and R_a per unit tree mass for the ambient and warmed treatment is indicative of homeostatic acclimation of maintenance respiration in this experiment (Fig. 8). The stimulation of whole-crown R_a by warming was primarily attributable to an increase in respiration to support growth. Thus, we suggest that this study is in agreement with previous published work from this experiment (Drake *et al.*, 2016b; Aspinwall *et al.*, 2016), where homeostatic acclimation of respiration to experimental warming prevented a warming-induced increase in maintenance respiration, while a warming effect on growth stimulated growth respiration and increased whole-crown R_a .

The increased allocation of C aboveground in the warmed treatment, combined with homeostatic acclimation of maintenance respiration, likely contributed to the observed warming-induced stimulation in growth during the first half of this experiment (Fig. 1). Experimental warming had neutral or negative effects on leaf-level photosynthetic rates in this study (Drake *et al.*, 2016b; Aspinwall *et al.*, 2016), so a warming-induced stimulation of growth was somewhat surprising. We suggest that an increase in C partitioning aboveground (Fig. 6a) was associated with accelerated leaf development early in the experiment in these young and rapidly growing trees (Fig. 1c), such that trees exposed to the warmed treatment had higher rates of crown-scale photosynthesis (Fig. 5a) primarily through a warming effect on total crown leaf area. That is, the warmed treatment got a head start in leaf area production, which compounded over time. This is consistent with nutrient fertilization studies, in which increases in leaf area rather than changes in leaf N or leaf function, often dominates the growth responses of rapidly growing plants (Sinclair & Horie, 1989; Gastal & Lemaire, 2002; Lovelock *et al.*, 2004; Wang *et al.*, 2012), although there are exceptions (Santiago *et al.*, 2012). This is also consistent with other experimental manipulations, in which a stimulation of leaf area development early in an experiment can strongly affect exponential growth trajectories (Tjoelker *et al.*, 1998; Kirschbaum, 2011; Drake *et al.*, 2017).

470 *Effects of drought on C allocation*

471 We did not detect any significant effects of drought on C partitioning. The drought appeared to
472 reduce all C fluxes proportionally, such that the ratios of C fluxes to GPP was unchanged. We recognize
473 that our ability to resolve C partitioning belowground was limited by the nature of the measurements
474 based on the residual, and our lack of root biomass measurements through time. The biomass in roots
475 relative to the total at the harvest (root mass ratio; Fig 3b) did indicate an interactive effect of drought and
476 warming, possibly via small differences in the partitioning of GPP to roots that accumulated over time.
477 Perhaps the ambient temperature trees had sufficient carbohydrate reserves to fuel additional root growth
478 in the drought treatment, while warmed temperature trees were consuming more carbohydrates
479 aboveground and were thus unable to increase root growth in the drought. This speculative process may
480 explain the observed interaction between warming and drought on root mass ratio (Fig. 3b).

481 Trees acquired water from deep in the soil profile during the drought. Leaf predawn water
482 potential declined to only approximately -0.5 MPa, which is a moderate value that is not indicative of
483 pronounced water stress. Thus, while we successfully implemented a drought that dried the soils from the
484 surface to 1-m-depth, the trees successfully avoided acute physiological drought stress by reducing
485 growth and transpiration rates while also acquiring deeper soil water. Previous studies have shown that
486 groundwater use enables vegetation to mitigate production declines under conditions of surface moisture
487 limitation (Baldocchi *et al.*, 2010; Barbeta *et al.*, 2015), and several eucalypt species are well-known
488 users of groundwater (Mensforth *et al.*, 1994; Pfautsch *et al.*, 2011, 2015; Eamus *et al.*, 2015; Zolfaghar
489 *et al.*, 2017). Furthermore, Koirala *et al.* (2017) demonstrated correlations between GPP and groundwater
490 table depth that were present over approximately 70% of the vegetated surface of the earth, suggesting
491 that vegetation-groundwater interactions are common and globally relevant. Our study demonstrates that
492 some trees may utilize access to soil water at depth to maintain moderate rates of photosynthetic C uptake
493 and growth during extended droughts that lead to dry surface soils.

494 *Implications for mathematical models*

495 Many ecosystem and earth system models begin their simulation of ecosystem C cycling by predicting
496 GPP as a function of leaf area and environmental drivers. GPP is then partitioned into component terms
497 including R_a and the production of leaf, wood, and root mass. Our observations suggest that tree C
498 allocation of GPP to these terms can be influenced by environmental drivers such as temperature. That is,
499 the observations presented here are not consistent with static partitioning schemes with fixed and constant
500 partitioning of GPP into component fluxes. However when aggregated over longer time periods, our
501 average partitioning coefficients (approximately 30% of GPP to aboveground respiration, 43% of GPP to
502 aboveground growth, and 27% to belowground fluxes) are comparable with values used in some

ecosystem models (De Kauwe *et al.*, 2014). The observations presented here are also not consistent with a dynamic C partitioning scheme based on Sprengel and Leibig's law of the minimum (van der Ploeg *et al.*, 1999), where C would be preferentially allocated to increase the acquisition of the factor most limiting primary production. Furthermore, Aspinwall *et al.* (2016) recently documented strong seasonal variation in carbohydrate storage in these evergreen trees, characterized by the buildup of high starch concentrations during the winter and a drawdown of these reserves during the summer. It appears that these trees partially utilize a non-structural carbohydrate storage reserve to fuel growth and metabolism during the hot summer. Based on these observations, we suggest that a dynamic allocation scheme incorporating a dynamic carbohydrate reserve may be appropriate for future investigation (e.g., Fatichi *et al.*, 2014; Pugh *et al.*, 2016).

Conclusions

We used a novel combination of growth and whole-crown flux measurements to study the effects of warming and drought on the C allocation of young *Eucalyptus tereticornis* trees. These trees did not substantially alter C allocation in response to the drought treatment, as reduced growth, reduced transpiration, and the facultative use of deep soil water allowed the trees to avoid physiological drought stress. Experimental warming increased the proportion of GPP that was allocated to aboveground uses and decreased the proportion of GPP that was allocated belowground. Such a change in tree C allocation may have important implications for tree growth, forest C storage, and soil nutrient cycling in a warmer world. In particular, increased allocation aboveground in a warmer world may stimulate leaf area development and aboveground growth during conditions of sufficient soil resource supply.

Acknowledgements

We thank Burhan Amiji (Western Sydney University) for maintaining the site, for collecting much of the growth and harvest data, and for his excellent research support. This experiment was made possible through a collaboration with Sune Linder and the Swedish University of Agricultural Sciences, who designed, built, and generously provided the whole tree chambers. We also gratefully acknowledge Courtney Campany (Colgate University) for his measurements of fine root biomass as well as Renee Smith, and Carrie Drake (Western Sydney University) for their help with the whole-tree harvests. This research was supported by the Australian Research Council (Discovery, DP140103415), a New South Wales government Climate Action Grant (NSW T07/CAG/016), the Hawkesbury Institute for the Environment, and Western Sydney University.

Authorship statement

534 JED co-led the experimental design, contributed to data collection for the chamber flux, and led
535 the data analysis, modeling, and writing. MGT was the senior scientific lead; he co-led the
536 experimental design and made large contributions to analysis, interpretation, and writing. MJA
537 contributed to the experimental design, data collection and interpretation, and writing. PBR
538 contributed to the experimental design, interpretation, and writing. SP contributed to
539 experimental design, data collection and interpretation, and writing. CVMB contributed to the
540 measurements of chamber flux, and contributed to experimental design, data analysis, and
541 writing.

542 Table 1. Estimate of aboveground growth and maintenance respiration coefficients for *Eucalyptus*
543 *tereticornis* trees derived from equation 2, with standard errors (SE), and 95% confidence intervals;
544 parameters were statistically equivalent across ambient and warmed treatments (all $P > 0.3$).

Term	Units	Ambient		Warmed	
		Mean (SE)	95% CI	Mean (SE)	95% CI
g_r (growth respiration rate)	g C respired per g C growth	0.32 (0.02)	0.27-0.37	0.28 (0.03)	0.21-0.36
m_r (maintenance respiration rate)	g C respired per g C standing aboveground biomass per day	0.015 (0.001)	0.012-0.019	0.017 (0.002)	0.013-0.021

Commented [JED2]: Added species name here.

545

546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579

Figure legends:

Figure 1. Growth of *Eucalyptus tereticornis* trees exposed to warming and drought. Trees were either exposed to ambient T_{air} (“A”, blue) or warming of +3 °C (“W”, red), and either a well-watered control (“Con”, solid points) or drought conditions (“Dry”, open points). Stem diameter (a) was measured at 65-cm height, and height reflects total stem length (b). Total leaf area was directly measured on three dates (c) and stem volume was calculated from diameter measurements along the stem of each tree (d). The vertical dashed line denotes when CO₂ and H₂O flux measurements began. Points reflect the mean, error bars denote 1SEM (n = 6 until Feb 2014, when the drought treatment began and n = 3).

Figure 2. Summary of measured CO₂ and H₂O fluxes for twelve *Eucalyptus tereticornis* trees exposed to ambient (A) or warmed (W) air temperatures in 2013 and 2014. All trees were maintained in control well-watered conditions (Con) until mid-Feb, when half of the trees were subjected to a soil drydown (Dry). We show weekly averages of the measured daily net C uptake (a) and the sum of net C uptake for the two measurement periods (pre-drought, drought; b). We also show weekly averages of the measured daily net H₂O loss to transpiration (c) and the H₂O loss to transpiration summed across the two measurement periods (d). In (a) and (c), lines reflect the mean and shaded areas reflect the standard error. The dotted vertical lines in (b) and (d) separate the pre-drought (left) and drought periods (right). These plots reflect >580,000 individual flux measurements at 15-minute resolution.

Figure 3. Biomass components at final harvest for twelve *Eucalyptus tereticornis* trees exposed to ambient (A) or warmed (W) air temperatures and either control well-watered conditions (Con) or a soil drydown treatment (Dry). Note that these data reflect grams of dry mass. Each of the measured biomass components (a) reflects the mean of three trees per treatment, the error bars reflect the standard error of the total measured mass. The root mass ratio (b) reflects the sum of coarse and fine roots relative to total tree mass (boxplot of three observations per category: the thick line reflects the median value and the whiskers reflect the other two values; the box shows the 25th and 75th percentile). Warming reduced the root mass ratio, while the drought treatment increased root mass ratio in the ambient temperature treatment only. The root mass ratio interaction primarily follows the response of fine roots, although stem wood and coarse roots also contributed.

Figure 4. Soil volumetric water content and predawn leaf water potential (Ψ_{pd}) for twelve *Eucalyptus tereticornis* trees exposed to ambient (A) or warmed (W) air temperatures. All trees were maintained in

Commented [JED3]: Added as requested

control well-watered conditions (Con) until mid-Feb, when half of the trees were subjected to a soil drydown (Dry). We show daily averages of the measured volumetric water content in surface soils (~0.1-m-depth; a), an intermediate depth (~0.5-m-depth; b), and in deep soils just above the hard layer of partially cemented manganese nodules (~1-m-depth; c). The horizontal dashed line reflects the volumetric water content at which soil matric potential drops to -1.5 MPa. We also show leaf Ψ_{pd} measured throughout the drydown (d). Points reflect the mean and error bars reflect the standard error (n = 6 or 3). Note that Ψ_{pd} was moderate in all treatments.

Figure 5. Fortnightly C fluxes for twelve *Eucalyptus tereticornis* trees exposed to ambient (A) or warmed (W) air temperatures. All fluxes are presented in units of g C tree⁻¹ fortnight⁻¹. All trees were maintained in control well-watered conditions until mid-Feb (Con), when half of the trees were subjected to a soil drydown (Dry). Solid lines reflect the mean of fortnightly data (i.e., two-week periods) and shaded areas reflect 1SEM. Measurements include gross primary production (GPP; a), aboveground net primary production (NPP_a; b), aboveground autotrophic respiration (R_a; c), and the residual (d). The residual reflects belowground C flux and measurement error. Note that the y-axis scale is twice as large for GPP relative to the other fluxes.

Figure 6. The fractional partitioning of gross primary production (GPP) for twelve *Eucalyptus tereticornis* trees. GPP was partitioned into aboveground net primary production (NPP_a; a-b), aboveground autotrophic respiration (R_a; c-d), and the residual C, which includes belowground C allocation and measurement error (e-f). The shaded areas in a-c reflect ±1SEM. All trees were maintained in control well-watered conditions (Con) until mid-Jan, when half of the trees were subjected to a soil drydown (Dry). Bar charts of flux partitioning terms (b, d, f) represent the mean (±1SEM), and the dotted vertical lines separate the pre-drought (left) and drought periods (right).

Commented [JED4]: Added as requested.

604
605 **Figure 7.** The fractional partitioning of gross primary production (GPP) for twelve *Eucalyptus*
606 *tereticornis* trees grown under ambient and elevated temperature. Data for each tree were averaged across
607 the pre-drought period; each point reflects an individual tree ($n = 6$). GPP was partitioned into
608 aboveground net primary production (NPP_a; a), aboveground autotrophic respiration (R_a; b), and the
609 residual C, which includes belowground C allocation (c). Solid lines reflect linear models fit to each
610 treatment; dashed lines reflect the 95% confidence interval.

611
612 **Figure 8.** Partitioning of aboveground respiration into maintenance and growth components for
613 *Eucalyptus tereticornis* trees in a warming and drought experiment. Each point reflects a tree during a
614 fortnightly growth interval. Note that the y-intercept reflects the maintenance respiration component and
615 the slope reflects the growth respiration component. Neither the slope nor the intercept were affected by
616 experiment treatments (mixed effects model with random intercepts for each chamber, $P > 0.5$). The solid
617 lines reflects models fit to the ambient temperature (A) and warmed temperature (W) data and dashed
618 lines reflect the 95% confidence interval. All of the data were well-described by a single linear function
619 ($Y = 0.021 + 0.0059x$, $r^2 = 0.64$, $P < 0.001$).

Commented [JED5]: Added species names here

620 References

621 **Adu-Bredu S, Hagihara A. 2003.** Long-term carbon budget of the above-ground parts of a
622 young hinoki cypress (*Chamaecyparis obtusa*) stand. *Ecological Research* **18**: 165–175.

623 **Allen CD, Breshears DD, McDowell NG. 2015.** On underestimation of global vulnerability to
624 tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* **6**: 129.

625 **Amthor JS. 2000.** The McCree–de Wit–Penning de Vries–Thornley respiration paradigms: 30
626 years later. *Annals of Botany* **86**: 1–20.

627 **Aspinwall MJ, Drake JE, Company C, Varhammar A, Ghannoum O, Tissue DT, Reich PB,**
628 **Tjoelker MG. 2016.** Convergent acclimation of leaf photosynthesis and respiration to prevailing
629 ambient temperatures under current and warmer climates in *Eucalyptus tereticornis*. *New*
630 *Phytologist* **212**: 354–367.

631 **Baldocchi DD, Ma S, Rambal S, Misson L, Ourcival J-M, Limousin J-M, Pereira J, Papale**
632 **D. 2010.** On the differential advantages of evergreenness and deciduousness in mediterranean
633 oak woodlands: a flux perspective. *Ecological Applications* **20**: 1583–1597.

634 **Barbeta A, Mejía-Chang M, Ogaya R, Voltas J, Dawson TE, Peñuelas J. 2015.** The
635 combined effects of a long-term experimental drought and an extreme drought on the use of
636 plant-water sources in a Mediterranean forest. *Global Change Biology* **21**: 1213–1225.

637 **Barton CVM, Duursma RA, Medlyn BE, Ellsworth DS, Eamus D, Tissue DT, Adams MA,**
638 **Conroy J, Crous KY, Liberloo M, et al. 2012.** Effects of elevated atmospheric [CO₂] on
639 instantaneous transpiration efficiency at leaf and canopy scales in *Eucalyptus saligna*. *Global*
640 *Change Biology* **18**: 585–595.

641 **Barton CVM, Ellsworth DS, Medlyn BE, Duursma RA, Tissue DT, Adams MA, Eamus D,**
642 **Conroy JP, McMurtrie RE, Parsby J, et al. 2010.** Whole-tree chambers for elevated
643 atmospheric CO₂ experimentation and tree scale flux measurements in south-eastern Australia:
644 The Hawkesbury Forest Experiment. *Agricultural and Forest Meteorology* **150**: 941–951.

645 **Blessing CH, Werner RA, Siegwolf R, Buchmann N. 2015.** Allocation dynamics of recently
646 fixed carbon in beech saplings in response to increased temperatures and drought. *Tree*
647 *Physiology* **35**: 585–598.

648 **Burke EJ, Brown SJ, Christidis N. 2006.** Modeling the recent evolution of global drought and
649 projections for the twenty-first century with the Hadley Centre climate model. *Journal of*
650 *Hydrometeorology* **7**: 1113–1125.

651 **De Kauwe MG, Medlyn BE, Zaehle S, Walker AP, Dietze MC, Wang Y-P, Luo Y, Jain AK, El-**
652 **Masri B, Hickler T, et al. 2014.** Where does the carbon go? A model–data intercomparison of
653 vegetation carbon allocation and turnover processes at two temperate forest free-air CO₂
654 enrichment sites. *New Phytologist* **203**: 883–899.

655 **DeLucia EH, Moore DJ, Norby RJ. 2005.** Contrasting responses of forest ecosystems to rising
656 atmospheric CO₂: Implications for the global C cycle. *Global Biogeochemical Cycles* **19**:
657 GB3006.

- 658 **Dietze MC, Sala A, Carbone MS, Czimczik CI, Mantooth JA, Richardson AD, Vargas R.**
 659 **2014.** Nonstructural carbon in woody plants. *Annual Review of Plant Biology* **65**: 667–687.
- 660 **Doughty CE, Malhi Y, Araujo-Murakami A, Metcalfe DB, Silva-Espejo JE, Arroyo L,**
 661 **Heredia JP, Pardo-Toledo E, Mendizabal LM, Rojas-Landivar VD, et al. 2014.** Allocation
 662 trade-offs dominate the response of tropical forest growth to seasonal and interannual drought.
 663 *Ecology* **95**: 2192–2201.
- 664 **Doughty CE, Metcalfe D, Girardin C a. J, Farfan Amezcuita F, Galiano Cabrera D, Huaraca**
 665 **Huasco W, Silva-Espejo JE, Araujo-Murakami A, Costa D, C M, et al. 2015.** Drought impact
 666 on forest carbon dynamics and fluxes in Amazonia. *Nature* **519**: 78–140.
- 667 **Drake JE, Aspinwall MJ, Pfautsch S, Rymer PD, Reich PB, Smith RA, Crous KY, Tissue**
 668 **DT, Ghannoum O, Tjoelker MG. 2015.** The capacity to cope with climate warming declines
 669 from temperate to tropical latitudes in two widely distributed Eucalyptus species. *Global Change*
 670 *Biology* **21**: 459–472.
- 671 **Drake JE, Gallet-Budynek A, Hofmockel KS, Bernhardt ES, Billings SA, Jackson RB,**
 672 **Johnsen KS, Lichter J, McCarthy HR, McCormack ML, et al. 2011.** Increases in the flux of
 673 carbon belowground stimulate nitrogen uptake and sustain the long-term enhancement of forest
 674 productivity under elevated CO₂. *Ecology Letters* **14**: 349–357.
- 675 **Drake J, Tjoelker M, Aspinwall MJ. 2016a.**
 676 https://figshare.com/articles/Drake_NewPhyt_2016_WTC3_RtoGPP_forfigshare_zip/3122104
 677 [Drake_NewPhyt_2016_WTC3_RtoGPP_forfigshare.zip](#).
 678 doi.org/10.6084/m9.figshare.3122104.v1
- 679 **Drake JE, Tjoelker MG, Aspinwall MJ, Reich PB, Barton CVM, Medlyn BE, Duursma RA.**
 680 **2016b.** Does physiological acclimation to climate warming stabilize the ratio of canopy
 681 respiration to photosynthesis? *New Phytologist* **211**: 850–863.
- 682 **Drake JE, Vårhammar A, Kumarathunge D, Medlyn BE, Pfautsch S, Reich PB, Tissue DT,**
 683 **Ghannoum O, Tjoelker MG. 2017.** A common thermal niche among geographically diverse
 684 populations of the widely distributed tree species *Eucalyptus tereticornis*: No evidence for
 685 adaptation to climate-of-origin. *Global Change Biology* **23**: 5069–5082.
- 686 **Duursma RA, Barton CVM, Eamus D, Medlyn BE, Ellsworth DS, Forster MA, Tissue DT,**
 687 **Linder S, McMurtrie RE. 2011.** Rooting depth explains [CO₂] x drought interaction in
 688 *Eucalyptus saligna*. *Tree Physiology* **31**: 922–931.
- 689 **Duursma RA, Barton CVM, Lin Y-S, Medlyn BE, Eamus D, Tissue DT, Ellsworth DS,**
 690 **McMurtrie RE. 2014.** The peaked response of transpiration rate to vapour pressure deficit in
 691 field conditions can be explained by the temperature optimum of photosynthesis. *Agricultural*
 692 *and Forest Meteorology* **189**: 2–10.
- 693 **Duursma RA, Falster DS. 2016.** Leaf mass per area, not total leaf area, drives differences in
 694 above-ground biomass distribution among woody plant functional types. *New Phytologist* **212**:
 695 368–376.

- 696 **Eamus D, Zolfaghar S, Villalobos-Vega R, Cleverly J, Huete A. 2015.** Groundwater-
697 dependent ecosystems: recent insights from satellite and field-based studies. *Hydrol. Earth*
698 *Syst. Sci.* **19**: 4229–4256.
- 699 **Edler B, Buerger J, Breitsameter L, Steinmann H-H, Isselstein J. 2015.** Growth responses to
700 elevated temperature and reduced soil moisture during early establishment of three annual
701 weeds in four soil types. *Journal of Plant Diseases and Protection* **122**: 39–48.
- 702 **Epron D, Bahn M, Derrien D, Lattanzi FA, Pumpanen J, Gessler A, Högberg P, Maillard P,**
703 **Dannoura M, Gérant D, et al. 2012.** Pulse-labelling trees to study carbon allocation dynamics:
704 a review of methods, current knowledge and future prospects. *Tree Physiology* **32**: 776–798.
- 705 **Farooq M, Wahid A, Kobayashi N, Fujita D, Basra SMA. 2009.** Plant drought stress: effects,
706 mechanisms and management. *Agronomy for Sustainable Development* **29**: 185–212.
- 707 **Fatichi S, Leuzinger S, Koerner C. 2014.** Moving beyond photosynthesis: from carbon source
708 to sink-driven vegetation modeling. *New Phytologist* **201**: 1086–1095.
- 709 **Feng L, Reffye P de, Dreyfus P, Auclair D. 2012.** Connecting an architectural plant model to a
710 forest stand dynamics model—application to Austrian black pine stand visualization. *Annals of*
711 *Forest Science* **69**: 245–255.
- 712 **Finzi AC, Abramoff RZ, Spiller KS, Brzostek ER, Darby BA, Kramer MA, Phillips RP. 2015.**
713 Rhizosphere processes are quantitatively important components of terrestrial carbon and
714 nutrient cycles. *Global Change Biology* **21**: 2082–2094.
- 715 **Franklin O, Johansson J, Dewar RC, Dieckmann U, McMurtrie RE, Brännström Å,**
716 **Dybzinski R. 2012.** Modeling carbon allocation in trees: a search for principles. *Tree Physiology*
717 **32**: 648–666.
- 718 **Friedlingstein P, Joel G, Field CB, Fung IY. 1999.** Toward an allocation scheme for global
719 terrestrial carbon models. *Global Change Biology* **5**: 755–770.
- 720 **Furze ME, Trumbore S, Hartmann H. 2018.** Detours on the phloem sugar highway: stem
721 carbon storage and remobilization. *Current Opinion in Plant Biology* **43**: 89–95.
- 722 **Gastal F, Lemaire G. 2002.** N uptake and distribution in crops: an agronomical and
723 ecophysiological perspective. *Journal of Experimental Botany* **53**: 789–799.
- 724 **Gower ST, Krankina O, Olson RJ, Apps M, Linder S, Wang C. 2001.** Net primary production
725 and carbon allocation patterns of boreal ecosystems. *Ecological Applications* **11**: 1395–1411.
- 726 **Hartmann H, McDowell NG, Trumbore S. 2015.** Allocation to carbon storage pools in Norway
727 spruce saplings under drought and low CO₂. *Tree Physiology* **35**: 243–252.
- 728 **Högberg P, Nordgren A, Buchmann N, Taylor AF, Ekblad A, Högberg MN, Nyberg G,**
729 **Ottosson-Löfvenius M, Read DJ. 2001.** Large-scale forest girdling shows that current
730 photosynthesis drives soil respiration. *Nature* **411**: 789–792.

- 731 **Hommel R, Siegwolf R, Zavadlav S, Arend M, Schaub M, Galiano L, Haeni M, Kayler ZE,**
 732 **Gessler A. 2016.** Impact of interspecific competition and drought on the allocation of new
 733 assimilates in trees. *Plant Biology* **18**: 785–796.
- 734 **Jump AS, Ruiz-Benito P, Greenwood S, Allen CD, Kitzberger T, Fensham R,**
 735 **Martínez-Vilalta J, Lloret F. 2017.** Structural overshoot of tree growth with climate variability
 736 and the global spectrum of drought-induced forest dieback. *Global Change Biology* **23**: 3742–
 737 3757.
- 738 **Kirschbaum MUF. 2011.** Does Enhanced Photosynthesis Enhance Growth? Lessons Learned
 739 from CO₂ Enrichment Studies. *Plant Physiology* **155**: 117–124.
- 740 **Koirala S, Jung M, Reichstein M, de Graaf IEM, Camps-Valls G, Ichii K, Papale D, Ráduly**
 741 **B, Schwalm CR, Tramontana G, et al. 2017.** Global distribution of groundwater-vegetation
 742 spatial covariation. *Geophysical Research Letters* **44**: 2017GL072885.
- 743 **Kuster TM, Arend M, Bleuler P, Günthardt-Goerg MS, Schulin R. 2013.** Water regime and
 744 growth of young oak stands subjected to air-warming and drought on two different forest soils in
 745 a model ecosystem experiment. *Plant Biology* **15**: 138–147.
- 746 **Landsberg JJ, Waring RH. 1997.** A generalised model of forest productivity using simplified
 747 concepts of radiation-use efficiency, carbon balance and partitioning. *Forest Ecology and*
 748 *Management* **95**: 209–228.
- 749 **Lemoine R, La Camera S, Atanassova R, Dedaldechamp F, Allario T, Pourtau N,**
 750 **Bonnemain J-L, Laloi M, Coutos-Thevenot P, Maurousset L, et al. 2013.** Source-to-sink
 751 transport of sugar and regulation by environmental factors. *Frontiers in Plant Science* **4**: 272.
- 752 **Leon-Sanchez L, Nicolas E, Nortes PA, Maestre FT, Querejeta JL. 2016.** Photosynthesis and
 753 growth reduction with warming are driven by nonstomatal limitations in a Mediterranean semi-
 754 arid shrub. *Ecology and Evolution* **6**: 2725–2738.
- 755 **Litton CM, Raich JW, Ryan MG. 2007.** Carbon allocation in forest ecosystems. *Global Change*
 756 *Biology* **13**: 2089–2109.
- 757 **Lovelock CE, Feller IC, Mckee KL, Engelbrecht BMJ, Ball MC. 2004.** The effect of nutrient
 758 enrichment on growth, photosynthesis and hydraulic conductance of dwarf mangroves in
 759 Panama. *Functional Ecology* **18**: 25–33.
- 760 **Lu M, Zhou X, Yang Q, Li H, Luo Y, Fang C, Chen J, Yang X, Li B. 2013.** Responses of
 761 ecosystem carbon cycle to experimental warming: a meta-analysis. *Ecology* **94**: 726–738.
- 762 **Mäkelä A, Valentine HT, Helmisaari H-S. 2008.** Optimal co-allocation of carbon and nitrogen
 763 in a forest stand at steady state. *New Phytologist* **180**: 114–123.
- 764 **Martínez-Vilalta J, Cochard H, Mencuccini M, Sterck F, Herrero A, Korhonen JFJ, Llorens**
 765 **P, Nikinmaa E, Nolè A, Poyatos R, et al. 2009.** Hydraulic adjustment of Scots pine across
 766 Europe. *New Phytologist* **184**: 353–364.

- 767 **McCree KJ. 1970.** An equation for the rate of respiration of white clover grown under controlled
768 conditions. p. 221–229. *Prediction and measurement of photosynthetic productivity.*
769 *Proceedings of the IBP/PP Technical Meeting, Trebon.* PUDOC, Wageningen, the Netherlands.
- 770 **McMurtrie RE, Dewar RC. 2013.** New insights into carbon allocation by trees from the
771 hypothesis that annual wood production is maximized. *New Phytologist* **199**: 981–990.
- 772 **Melillo JM, Butler S, Johnson J, Mohan J, Steudler P, Lux H, Burrows E, Bowles F, Smith**
773 **R, Scott L, et al. 2011.** Soil warming, carbon–nitrogen interactions, and forest carbon budgets.
774 *Proceedings of the National Academy of Sciences* **108**: 9508–9512.
- 775 **Melillo JM, Steudler PA, Aber JD, Newkirk K, Lux H, Bowles FP, Catricala C, Magill A,**
776 **Ahrens T, Morrisseau S. 2002.** Soil warming and carbon-cycle feedbacks to the climate
777 system. *Science* **298**: 2173–2176.
- 778 **Mencuccini M. 2003.** The ecological significance of long-distance water transport: short-term
779 regulation, long-term acclimation and the hydraulic costs of stature across plant life forms. *Plant,*
780 *Cell & Environment* **26**: 163–182.
- 781 **Mensforth LJ, Thorburn PJ, Tyerman SD, Walker GR. 1994.** Sources of water used by
782 riparian *Eucalyptus camaldulensis* overlying highly saline groundwater. *Oecologia* **100**: 21–28.
- 783 **Mokany K, Raison RJ, Prokushkin AS. 2006.** Critical analysis of root : shoot ratios in
784 terrestrial biomes. *Global Change Biology* **12**: 84–96.
- 785 **Munir TM, Perkins M, Kaing E, Strack M. 2015.** Carbon dioxide flux and net primary
786 production of a boreal treed bog: Responses to warming and water-table-lowering simulations of
787 climate change. *Biogeosciences* **12**: 1091–1111.
- 788 **Nemani RR, Keeling CD, Hashimoto H, Jolly WM, Piper SC, Tucker CJ, Myneni RB,**
789 **Running SW. 2003.** Climate-driven increases in global terrestrial net primary production from
790 1982 to 1999. *Science* **300**: 1560–1563.
- 791 **Palmroth S, Oren R, McCarthy HR, Johnsen KH, Finzi AC, Butnor JR, Ryan MG,**
792 **Schlesinger WH. 2006.** Aboveground sink strength in forests controls the allocation of carbon
793 below ground and its [CO₂]-induced enhancement. *Proceedings of the National Academy of*
794 *Sciences* **103**: 19362–19367.
- 795 **Pfautsch S, Dodson W, Madden S, Adams MA. 2015.** Assessing the impact of large-scale
796 water table modifications on riparian trees: a case study from Australia. *Ecohydrology* **8**: 642–
797 651.
- 798 **Pfautsch S, Keitel C, Turnbull TL, Braimbridge MJ, Wright TE, Simpson RR, O'Brien JA,**
799 **Adams MA. 2011.** Diurnal patterns of water use in *Eucalyptus victrix* indicate pronounced
800 desiccation–rehydration cycles despite unlimited water supply. *Tree Physiology* **31**: 1041–1051.
- 801 **van der Ploeg RR, Bořhm W, Kirkham MB. 1999.** On the origin of the theory of mineral
802 nutrition of plants and the law of the minimum. **63**: 1055–1062.
- 803 **Poorter H, Jagodzinski AM, Ruiz-Peinado R, Kuyah S, Luo Y, Oleksyn J, Usoltsev VA,**
804 **Buckley TN, Reich PB, Sack L. 2015.** How does biomass distribution change with size and

- 805 differ among species? An analysis for 1200 plant species from five continents. *New Phytologist*
806 **208**: 736–749.
- 807 **Poorter H, Niklas KJ, Reich PB, Oleksyn J, Poot P, Mommer L. 2012.** Biomass allocation to
808 leaves, stems and roots: meta-analyses of interspecific variation and environmental control.
809 *New Phytologist* **193**: 30–50.
- 810 **Poorter H, Sack L. 2012.** Pitfalls and possibilities in the analysis of biomass allocation patterns
811 in plants. *Frontiers in Plant Science* **3**: 259.
- 812 **Pugh T a. M, Mueller C, Arneth A, Haverd V, Smith B. 2016.** Key knowledge and data gaps in
813 modelling the influence of CO₂ concentration on the terrestrial carbon sink. *Journal of Plant*
814 *Physiology* **203**: 3–15.
- 815 **Reich PB. 2002.** Root-shoot relations: Optimality in acclimation and adaptation or the
816 'Emperor's New Clothes. In: *Plant roots: the hidden half*, eds: Waisel Y, Eshel A, Kafafi U,
817 Marcel Dekker, New York, USA. p 205–220.
- 818 **Reich PB, Luo Y, Bradford JB, Poorter H, Perry CH, Oleksyn J. 2014.** Temperature drives
819 global patterns in forest biomass distribution in leaves, stems, and roots. *Proceedings of the*
820 *National Academy of Sciences* **111**: 13721–13726.
- 821 **Reichstein M, Falge E, Baldocchi D, Papale D, Aubinet M, Berbigier P, Bernhofer C,**
822 **Buchmann N, Gilmanov T, Granier A, et al. 2005.** On the separation of net ecosystem
823 exchange into assimilation and ecosystem respiration: review and improved algorithm. *Global*
824 *Change Biology* **11**: 1424–1439.
- 825 **Roux XL, Lacointe A, Escobar-Gutiérrez A, Dizès SL. 2001.** Carbon-based models of
826 individual tree growth: A critical appraisal. *Annals of Forest Science* **58**: 469–506.
- 827 **Running SW, Gower ST. 1991.** FOREST-BGC, A general model of forest ecosystem
828 processes for regional applications. II. Dynamic carbon allocation and nitrogen budgets. *Tree*
829 *Physiology* **9**: 147–160.
- 830 **Rustad L, Campbell J, Marion G, Norby R, Mitchell M, Hartley A, Cornelissen J, Gurevitch**
831 **J, GCTE-NEWS. 2001.** A meta-analysis of the response of soil respiration, net nitrogen
832 mineralization, and aboveground plant growth to experimental ecosystem warming. *Oecologia*
833 **126**: 543–562.
- 834 **Santiago LS, Wright SJ, Harms KE, Yavitt JB, Korine C, Garcia MN, Turner BL. 2012.**
835 Tropical tree seedling growth responses to nitrogen, phosphorus and potassium addition.
836 *Journal of Ecology* **100**: 309–316.
- 837 **Sillmann J, Kharin VV, Zwiers FW, Zhang X, Brunaugh D. 2013.** Climate extremes indices in
838 the CMIP5 multimodel ensemble: Part 2. Future climate projections. *Journal of Geophysical*
839 *Research-Atmospheres* **118**: 2473–2493.
- 840 **Sinclair TR, Horie T. 1989.** Leaf Nitrogen, Photosynthesis, and Crop Radiation Use Efficiency:
841 A Review. *Crop Science* **29**: 90.

842 **Strömgren M, Linder S. 2002.** Effects of nutrition and soil warming on stemwood production in
843 a boreal Norway spruce stand. *Global Change Biology* **8**: 1194–1204.

844 **Taege S, Sparks TH, Menzel A. 2015.** Effects of temperature and drought manipulations on
845 seedlings of Scots pine provenances. *Plant Biology* **17**: 361–372.

846 **Thomas DS, Montagu KD, Conroy JP. 2007.** Temperature effects on wood anatomy, wood
847 density, photosynthesis and biomass partitioning of *Eucalyptus grandis* seedlings. *Tree*
848 *Physiology* **27**: 251–260.

849 **Tjoelker MG, Oleksyn J, Reich PB. 1998.** Temperature and ontogeny mediate growth
850 response to elevated CO₂ in seedlings of five boreal tree species. *The New Phytologist* **140**:
851 197–210.

852 **Tjoelker MG, Oleksyn J, Reich PB. 1999.** Acclimation of respiration to temperature and CO₂ in
853 seedlings of boreal tree species in relation to plant size and relative growth rate. *Global Change*
854 *Biology* **5**: 679–691.

855 **Trugman AT, Detto M, Bartlett MK, Medvigy D, Anderegg WRL, Schwalm C, Schaffer B,**
856 **Pacala SW. 2018.** Tree carbon allocation explains forest drought-kill and recovery patterns.
857 *Ecology Letters* **21**: 1552–1560.

858 **Volder A, Briske DD, Tjoelker MG. 2013.** Climate warming and precipitation redistribution
859 modify tree–grass interactions and tree species establishment in a warm-temperate savanna.
860 *Global Change Biology* **19**: 843–857.

861 **Wang D, Maughan MW, Sun J, Feng X, Miguez F, Lee D, Dietze MC. 2012.** Impact of
862 nitrogen allocation on growth and photosynthesis of *Miscanthus* (*Miscanthus x giganteus*).
863 *Global Change Biology Bioenergy* **4**: 688–697.

864 **Way DA, Oren R. 2010.** Differential responses to changes in growth temperature between trees
865 from different functional groups and biomes: a review and synthesis of data. *Tree Physiology*
866 **30**: 669–688.

867 **Yang J, Medlyn BE, Kauwe MGD, Duursma RA. 2018.** Applying the concept of
868 ecohydrological equilibrium to predict steady state leaf area index. *Journal of Advances in*
869 *Modeling Earth Systems* **10**: 1740–1758.

870 **Zolfaghar S, Villalobos-Vega R, Zeppel M, Cleverly J, Rumman R, Hingee M, Boulain N, Li**
871 **Z, Eamus D. 2017.** Transpiration of *Eucalyptus* woodlands across a natural gradient of depth-
872 to-groundwater. *Tree Physiology* **37**: 961–975.

873















