# Section 7.3: Trigonometric Substitution

Amarnath Patel

January 29, 2024

## 1 Introduction

Trigonometric substitution is a method for solving integrals and differential equations and is often used when the integrand involves the square root of a quadratic polynomial. It is also useful for integrals involving powers of trigonometric functions.

## 2 Method

The method involves substituting a variable with a trigonometric function. There are three main forms of trigonometric substitution:

- $x = a\sin(\theta)$, used when the integral contains $\sqrt{a^2 - x^2}$
- $x = a\tan(\theta)$, used when the integral contains $\sqrt{a^2 + x^2}$
- $x = a\sec(\theta)$, used when the integral contains $\sqrt{x^2 - a^2}$

## 3 Example

Let's look at an example. Suppose we want to integrate $\int \sqrt{1 - x^2}\, dx$.

For this integral, we can use the substitution $x = \sin(\theta)$:

$$\begin{aligned}
\int \sqrt{1 - x^2}\, dx &= \int \sqrt{1 - \sin^2(\theta)}\, d(\sin(\theta)) \\
&= \int \cos(\theta)\, d(\sin(\theta)) \\
&= \sin(\theta) + C \\
&= x + C
\end{aligned}$$

So, $\int \sqrt{1 - x^2}\, dx = x + C$.

The following table lists the common forms of trigonometric substitution:

**Table of Trigonometric Substitutions**

| Expression | Substitution | Identity |
|---|---|---|
| $\sqrt{a^2 - x^2}$ | $x = a \sin\theta, \quad -\dfrac{\pi}{2} \leqslant \theta \leqslant \dfrac{\pi}{2}$ | $1 - \sin^2\theta = \cos^2\theta$ |
| $\sqrt{a^2 + x^2}$ | $x = a \tan\theta, \quad -\dfrac{\pi}{2} < \theta < \dfrac{\pi}{2}$ | $1 + \tan^2\theta = \sec^2\theta$ |
| $\sqrt{x^2 - a^2}$ | $x = a \sec\theta, \quad 0 \leqslant \theta < \dfrac{\pi}{2}$ or $\pi \leqslant \theta < \dfrac{3\pi}{2}$ | $\sec^2\theta - 1 = \tan^2\theta$ |

**EXAMPLE 1** Evaluate $\displaystyle\int \frac{\sqrt{9 - x^2}}{x^2}\,dx$.

**SOLUTION** Let $x = 3\sin\theta$, where $-\pi/2 \leqslant \theta \leqslant \pi/2$. Then $dx = 3\cos\theta\,d\theta$ and

$$\sqrt{9 - x^2} = \sqrt{9 - 9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3\,|\cos\theta| = 3\cos\theta$$

(Note that $\cos\theta \geqslant 0$ because $-\pi/2 \leqslant \theta \leqslant \pi/2$.) Thus the Inverse Substitution Rule gives

$$\begin{aligned}\int \frac{\sqrt{9 - x^2}}{x^2}\,dx &= \int \frac{3\cos\theta}{9\sin^2\theta}\,3\cos\theta\,d\theta \\ &= \int \frac{\cos^2\theta}{\sin^2\theta}\,d\theta = \int \cot^2\theta\,d\theta \\ &= \int (\csc^2\theta - 1)\,d\theta \\ &= -\cot\theta - \theta + C\end{aligned}$$




**FIGURE 1**
$\sin\theta = \dfrac{x}{3}$

Because this is an indefinite integral, we must return to the original variable $x$. This can be done either by using trigonometric identities to express $\cot\theta$ in terms of $\sin\theta = x/3$ or by drawing a diagram, as in Figure 1, where $\theta$ is interpreted as an angle of a right triangle. Since $\sin\theta = x/3$, we label the opposite side and the hypotenuse as having lengths $x$ and 3. Then the Pythagorean Theorem gives the length of the adjacent side as $\sqrt{9 - x^2}$, so we can simply read the value of $\cot\theta$ from the figure:

$$\cot\theta = \frac{\sqrt{9 - x^2}}{x}$$

(Although $\theta > 0$ in the diagram, this expression for $\cot\theta$ is valid even when $\theta < 0$.) Since $\sin\theta = x/3$, we have $\theta = \sin^{-1}(x/3)$ and so

$$\int \frac{\sqrt{9 - x^2}}{x^2}\,dx = -\cot\theta - \theta + C = -\frac{\sqrt{9 - x^2}}{x} - \sin^{-1}\left(\frac{x}{3}\right) + C \qquad \blacksquare$$

**EXAMPLE 2** Find the area enclosed by the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$




**FIGURE 2**
$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

**SOLUTION** Solving the equation of the ellipse for $y$, we get

$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2} = \frac{a^2 - x^2}{a^2} \qquad \text{or} \qquad y = \pm\frac{b}{a}\sqrt{a^2 - x^2}$$

Because the ellipse is symmetric with respect to both axes, the total area $A$ is four times the area in the first quadrant (see Figure 2). The part of the ellipse in the first quadrant is given by the function

$$y = \frac{b}{a}\sqrt{a^2 - x^2} \qquad 0 \leqslant x \leqslant a$$

and so

$$\tfrac{1}{4}A = \int_0^a \frac{b}{a}\sqrt{a^2 - x^2}\,dx$$

To evaluate this integral we substitute $x = a\sin\theta$. Then $dx = a\cos\theta\,d\theta$. To change

Copyright 2021 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. WCN 02-200-203
Copyright 2021 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.



the limits of integration we note that when $x = 0$, $\sin\theta = 0$, so $\theta = 0$; when $x = a$, $\sin\theta = 1$, so $\theta = \pi/2$. Also

$$\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2\sin^2\theta} = \sqrt{a^2\cos^2\theta} = a\,|\cos\theta| = a\cos\theta$$

since $0 \leqslant \theta \leqslant \pi/2$. Therefore

$$\begin{aligned} A &= 4\frac{b}{a}\int_0^a \sqrt{a^2 - x^2}\,dx = 4\frac{b}{a}\int_0^{\pi/2} a\cos\theta \cdot a\cos\theta\,d\theta \\ &= 4ab\int_0^{\pi/2}\cos^2\theta\,d\theta = 4ab\int_0^{\pi/2}\tfrac{1}{2}(1 + \cos 2\theta)\,d\theta \\ &= 2ab\Big[\theta + \tfrac{1}{2}\sin 2\theta\Big]_0^{\pi/2} = 2ab\left(\frac{\pi}{2} + 0 - 0\right) = \pi ab \end{aligned}$$

We have shown that the area of an ellipse with semiaxes $a$ and $b$ is $\pi ab$. In particular, taking $a = b = r$, we have proved the famous formula that the area of a circle with radius $r$ is $\pi r^2$. ■

**NOTE** Because the integral in Example 2 was a definite integral, we changed the limits of integration and did not have to convert back to the original variable $x$.

**EXAMPLE 5** Evaluate $\displaystyle\int \frac{dx}{\sqrt{x^2 - a^2}}$, where $a > 0$.

**SOLUTION 1** We let $x = a\sec\theta$, where $0 < \theta < \pi/2$ or $\pi < \theta < 3\pi/2$. Then $dx = a\sec\theta\tan\theta\,d\theta$ and

$$\sqrt{x^2 - a^2} = \sqrt{a^2(\sec^2\theta - 1)} = \sqrt{a^2\tan^2\theta} = a\,|\tan\theta| = a\tan\theta$$

Therefore

$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \int \frac{a\sec\theta\tan\theta}{a\tan\theta}\,d\theta = \int \sec\theta\,d\theta = \ln|\sec\theta + \tan\theta| + C$$




**FIGURE 4**
$\sec\theta = \frac{x}{a}$

The triangle in Figure 4 gives $\tan\theta = \sqrt{x^2 - a^2}/a$, so we have

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \ln\left|\frac{x}{a} + \frac{\sqrt{x^2 - a^2}}{a}\right| + C \\ &= \ln\left|x + \sqrt{x^2 - a^2}\right| - \ln a + C \end{aligned}$$

Writing $C_1 = C - \ln a$, we have

$$\boxed{1} \qquad \int \frac{dx}{\sqrt{x^2 - a^2}} = \ln\left|x + \sqrt{x^2 - a^2}\right| + C_1$$