

Swarm Intelligence

Exercise 4

Matthias Kergaßner

Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU)
Department Informatik


19.07.2022 – 22.07.2022



Disclaimer

This set of slides is for personal use only and may not be passed on to third parties.

Problem 13: PageRank Algorithm




Idea and Goal of the PageRank Algorithm

- Random-Walk across the internet
- The more often a page is reached, the higher its PageRank
- Determination of the relevance of websites

Problem 13 (a)

Start at website 1 and run two steps.


Where will one end up with which probability?



Problem 13 (a)

Start at website 1 and run two steps.

Where will one end up with which probability?




Two possibilities:

Problem 13 (a)

Start at website 1 and run two steps.

Where will one end up with which probability?




Two possibilities:

- $1 \rightarrow 2 \rightarrow 3$ with 50% chance

Problem 13 (a)

Start at website 1 and run two steps.

Where will one end up with which probability?



Two possibilities:


- $1 \rightarrow 2 \rightarrow 3$ with 50% chance
- $1 \rightarrow 2 \rightarrow 4$ with 50% chance

Problem 13 (b)

Determine the transition matrix A , such that

$$\pi_{t+1} = \pi_t \cdot A$$

holds.




Problem 13 (b)

Determine the transition matrix A , such that

$$\pi_{t+1} = \pi_t \cdot A$$

holds.




$$A = \begin{pmatrix} 0 & * & 0 & 0 \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \\ * & * & 0 & 0 \end{pmatrix}$$

Problem 13 (b)

Determine the transition matrix A , such that

$$\pi_{t+1} = \pi_t \cdot A$$


holds.



$$A = \begin{pmatrix} 0 & * & 0 & 0 \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \\ * & * & 0 & 0 \end{pmatrix} \rightarrow A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 1 \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$


Problem 13 (c)

Determine the limit π to which this Markov chain converges.



Problem 13 (c)


Determine the limit π to which this Markov chain converges.



Proof of convergence:

Problem 13 (c)

Determine the limit π to which this Markov chain converges.




Proof of convergence:

- A is **irreducible**, since every node is reachable from every other node

Problem 13 (c)

Determine the limit π to which this Markov chain converges.



Proof of convergence:


- A is **irreducible**, since every node is reachable from every other node
- A is **aperiodic**, since node 2 can be revisited after two as well as three steps:

$$2 \rightarrow 4 \rightarrow 2$$

$$2 \rightarrow 3 \rightarrow 4 \rightarrow 2$$

Problem 13 (c)

Determine the limit π to which this Markov chain converges.



Proof of convergence:

- A is **irreducible**, since every node is reachable from every other node
- A is **aperiodic**, since node 2 can be revisited after two as well as three steps:

$$2 \rightarrow 4 \rightarrow 2$$

$$2 \rightarrow 3 \rightarrow 4 \rightarrow 2$$

Thus the Markov chain does converge.

Problem 13 (c)

Determine the limit π to which this Markov chain converges.

Compute (left) eigenvector π corresponding to the largest eigenvalue 1:

Problem 13 (c)

Determine the limit π to which this Markov chain converges.

Compute (left) eigenvector π corresponding to the largest eigenvalue 1:

$$\begin{aligned} \pi \cdot (A - \mathbb{1}) &= (0, \dots, 0) \\ \Leftrightarrow (A - \mathbb{1})^\top \cdot \pi^\top &= (0, \dots, 0)^\top \end{aligned}$$

Problem 13 (c)

Determine the limit π to which this Markov chain converges.

Compute (left) eigenvector π corresponding to the largest eigenvalue 1:

$$\begin{aligned}\pi \cdot (A - \mathbb{1}) &= (0, \dots, 0) \\ \Leftrightarrow (A - \mathbb{1})^\top \cdot \pi^\top &= (0, \dots, 0)^\top\end{aligned}$$

Kernel of $(A - \mathbb{1})^\top$:

$$(A - \mathbb{1})^\top = \begin{pmatrix} -1 & 0 & 0 & \frac{1}{2} \\ 1 & -1 & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & -1 & 0 \\ 0 & \frac{1}{2} & 1 & -1 \end{pmatrix}$$

Problem 13 (c)

Determine the limit π to which this Markov chain converges.

Compute (left) eigenvector π corresponding to the largest eigenvalue 1:

$$\begin{aligned}\pi \cdot (A - \mathbb{1}) &= (0, \dots, 0) \\ \Leftrightarrow (A - \mathbb{1})^\top \cdot \pi^\top &= (0, \dots, 0)^\top\end{aligned}$$

Kernel of $(A - \mathbb{1})^\top$:

$$(A - \mathbb{1})^\top = \begin{pmatrix} -1 & 0 & 0 & \frac{1}{2} \\ 1 & -1 & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & -1 & 0 \\ 0 & \frac{1}{2} & 1 & -1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 0 & -\frac{1}{2} \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -\frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{2} \end{pmatrix}$$

Problem 13 (c)

Determine the limit π to which this Markov chain converges.

Compute (left) eigenvector π corresponding to the largest eigenvalue 1:

$$\begin{aligned}\pi \cdot (A - \mathbb{1}) &= (0, \dots, 0) \\ \Leftrightarrow (A - \mathbb{1})^\top \cdot \pi^\top &= (0, \dots, 0)^\top\end{aligned}$$

Kernel of $(A - \mathbb{1})^\top$:

$$(A - \mathbb{1})^\top = \begin{pmatrix} -1 & 0 & 0 & \frac{1}{2} \\ 1 & -1 & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & -1 & 0 \\ 0 & \frac{1}{2} & 1 & -1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 0 & -\frac{1}{2} \\ 0 & 1 & 0 & -\frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{2} \end{pmatrix} \rightarrow \left\langle \left(\frac{1}{2}, 1, \frac{1}{2}, 1 \right)^\top \right\rangle$$

Problem 13 (c)

Determine the limit π to which this Markov chain converges.

Compute (left) eigenvector π corresponding to the largest eigenvalue 1:

$$\begin{aligned}\pi \cdot (A - \mathbb{1}) &= (0, \dots, 0) \\ \Leftrightarrow (A - \mathbb{1})^\top \cdot \pi^\top &= (0, \dots, 0)^\top\end{aligned}$$

Kernel of $(A - \mathbb{1})^\top$:

$$(A - \mathbb{1})^\top = \begin{pmatrix} -1 & 0 & 0 & \frac{1}{2} \\ 1 & -1 & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & -1 & 0 \\ 0 & \frac{1}{2} & 1 & -1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 0 & -\frac{1}{2} \\ 0 & 1 & 0 & -\frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{2} \end{pmatrix} \rightarrow \left\langle \left(\frac{1}{2}, 1, \frac{1}{2}, 1 \right)^\top \right\rangle$$

Choose normalized eigenvector $\pi = \frac{1}{6} \cdot (1, 2, 1, 2)$

Problem 13 (c)

Determine the limit π to which this Markov chain converges.

Compute (left) eigenvector π corresponding to the largest eigenvalue 1:

$$\begin{aligned}\pi \cdot (A - \mathbb{1}) &= (0, \dots, 0) \\ \Leftrightarrow (A - \mathbb{1})^\top \cdot \pi^\top &= (0, \dots, 0)^\top\end{aligned}$$

Kernel of $(A - \mathbb{1})^\top$:

$$(A - \mathbb{1})^\top = \begin{pmatrix} -1 & 0 & 0 & \frac{1}{2} \\ 1 & -1 & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & -1 & 0 \\ 0 & \frac{1}{2} & 1 & -1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 0 & -\frac{1}{2} \\ 0 & 1 & 0 & -\frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{2} \end{pmatrix} \rightarrow \left\langle \left(\frac{1}{2}, 1, \frac{1}{2}, 1 \right)^\top \right\rangle$$

Choose normalized eigenvector $\pi = \frac{1}{6} \cdot (1, 2, 1, 2)$

Thus the following PageRanks come up:

$$(2, 4, 1, 3)$$

Problem 13 (d)


Remove link from 3 to 4. What is the problem now?

Problem 13 (d)

Remove link from 3 to 4. What is the problem now?

Previously:

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 1 \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$




Problem 13 (d)

Remove link from 3 to 4. What is the problem now?


Previously:

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 1 \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$



Without edge (3 → 4):

$$B =$$




Problem 13 (d)

Remove link from 3 to 4. What is the problem now?


Previously:

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 1 \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$



Without edge (3 → 4):

$$B = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & \mathbf{1} & \mathbf{0} \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$




Problem 13 (d)

Remove link from 3 to 4. What is the problem now?


Previously:

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 1 \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$



Without edge $(3 \rightarrow 4)$:

$$B = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & \mathbf{1} & \mathbf{0} \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$




$p \cdot B^t$ converges to $(0, 0, 1, 0)$ for $t \rightarrow \infty$, although page 3 seems to be rather irrelevant (\rightarrow absorption).

Problem 13 (d)

Remove link from 3 to 4. What is the problem now?


$$B = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & \mathbf{1} & \mathbf{0} \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$



Problem 13 (d)

Remove link from 3 to 4. What is the problem now?

$$B = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & \mathbf{1} & \mathbf{0} \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$




Solution 1: Remove node 3

Problem 13 (d)

Remove link from 3 to 4. What is the problem now?

$$B = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & \mathbf{1} & \mathbf{0} \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$




Solution 1: Remove node 3

$$B_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix} \rightarrow \pi_{B_1} = \frac{1}{5} \cdot (1, 2, 2)$$

Problem 13 (d)

Remove link from 3 to 4. What is the problem now?

$$B = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & \mathbf{1} & \mathbf{0} \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$



Solution 1: Remove node 3


$$B_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix} \rightarrow \pi_{B_1} = \frac{1}{5} \cdot (1, 2, 2)$$

Solution 2: Add edges from 3 to every other node

Problem 13 (d)

Remove link from 3 to 4. What is the problem now?

$$B = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & \mathbf{1} & \mathbf{0} \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}$$



Solution 1: Remove node 3

$$B_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix} \rightarrow \pi_{B_1} = \frac{1}{5} \cdot (1, 2, 2)$$

Solution 2: Add edges from 3 to every other node

$$B_2 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix} \rightarrow \pi_{B_2} = \frac{1}{17} \cdot (3, 6, 4, 4)$$

Problem 13 (d)

Remove link from 3 to 4. What is the problem now?

Additional trick: Ensure **irreducibility** and **aperiodicity**

Problem 13 (d)

Remove link from 3 to 4. What is the problem now?

Additional trick: Ensure **irreducibility** and **aperiodicity**

$$\tilde{A} = \alpha \cdot A + (1 - \alpha) \cdot \begin{pmatrix} 1 & \dots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \dots & 1 \end{pmatrix}$$

Problem 13 (d)

Remove link from 3 to 4. What is the problem now?

Additional trick: Ensure **irreducibility** and **aperiodicity**

$$\tilde{A} = \alpha \cdot A + (1 - \alpha) \cdot \begin{pmatrix} 1 & \dots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \dots & 1 \end{pmatrix} \cdot \frac{1}{n}$$

Problem 13 (d)

Remove link from 3 to 4. What is the problem now?

Additional trick: Ensure **irreducibility** and **aperiodicity**

$$\tilde{A} = \alpha \cdot A + (1 - \alpha) \cdot \begin{pmatrix} 1 & \dots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \dots & 1 \end{pmatrix} \cdot \frac{1}{n}$$

“Good” compromise for $\alpha = 0.85$

Thank you for your attention! Any Questions?

