



FLAME
UNIVERSITY

EVERLASTING
learning

FUNDAMENTALS OF COMPUTER GRAPHICS (CSIT304)

SURFACE REPRESENTATION

CHIRANJOY CHATTOPADHYAY

Associate Professor,
FLAME School of Computation and Data Science


OBJECTIVE

After completing this lecture, students will be able to


- Mathematically represent a surface patches
 - Planar Surface
 - Curved Surface
 - Bi-linear Surface
 - Lofted or Ruled Surface
- Solve related mathematical problems

SURFACE

- Representation of surface
 - Explicit
 - Implicit
 - Parametric




A SURFACE PATCH



DISPLAYING A SURFACE PATCH

- A surface patch can be displayed either as a wire frame



NON-PARAMETRIC SURFACES

- A quadratic surface would have an equation of the form:

$$ax^2 + ey^2 + hz^2 + 2bxy + 2cxz + 2fyz + 2dx + 2gy + 2iz + 1 = 0$$

- Can we represent this as matrix multiplication?

$$\begin{bmatrix} a & b & c & d \\ b & e & f & g \\ c & f & h & i \\ d & g & i & 1 \end{bmatrix}$$

PARAMETRIC SURFACES

- A point on a surface patch is a function of two parameters

$$P(u, v) = au^2 + 2buv + 2cu + dv^2 + 2ev + f = 0$$

- Can we represent this as matrix multiplication?

$$P(u, v) = [u \ v \ 1] \begin{bmatrix} a & b & c \\ b & d & e \\ c & e & f \end{bmatrix} \begin{bmatrix} u \\ v \\ 1 \end{bmatrix}$$

SURFACE EDGES

- The surface has edges given by the four curves for which one of the parameters is either 0 or 1.

$$P(u, v) = au^2 + 2buv + 2cu + dv^2 + 2ev + f = 0$$

$$P(0, v) = ?$$


$$P(1, v) = ?$$

$$P(u, 0) = ?$$

$$P(u, 1) = ?$$

VISUALIZATION

$$P(u, v) = au^2 + 2buv + 2cu + dv^2 + 2ev + f = 0$$



	u	v
P_0	0	0
P_1	0	1
P_2	1	0
P_3	1	1
P_4	$\frac{1}{2}$	0
P_5	$\frac{1}{2}$	1


CURVED SURFACE REPRESENTATION

- A curve is represented as a vector valued function of a single variable
- A surface is represented by a bivariate vector-valued function
- How to represent a curve on the surface?
- How a point on the surface may be represented?


LINEAR COONS SURFACE

- If the four boundary curves $P(u, 0)$, $P(u, 1)$, $P(0, v)$, and $P(1, v)$ are known, then?
- Intuitive solution
- What about the corner points and at the edges?
- Solution?
- Matrix representation of the linear Coons surface

EXAMPLE



PARAMETRIC SURFACE PATCH




$P(u, v) = ?$ [INTERPOLATION]

$P(u, v) = P(u, 0)(1 - v) + P(u, 1)v + P(0, v)(1 - u) + P(1, v)u$ [INCORRECT]

$P(u, v) = P(u, 0)(1 - v) + P(u, 1)v + P(0, v)(1 - u) + P(1, v)u -$
 $P(0, 0)(1 - u)(1 - v) - P(0, 1)(1 - u)v + P(1, 0)u(1 - v) - P(1, 1)uv$

EXAMPLE OF CONSTRUCTING A COON'S PATCH

IDENTIFY THE CORNER POINTS



EXAMPLE OF CONSTRUCTING A COON'S PATCH

(y, v)

(x, u)

	2	3	4	5	6	7
7
8	.	.	10	9		
9	.	14	12	11	10	.
10	.	15	13	14	10	.
11	.	.	10	11		

- Gradients at the corners of the patch

$$P(0,0) = (9,4,12)$$

$$\frac{\partial(P(0,0))}{\partial u} = ((10, 4, 13) - (8, 4, 10))/2 = (1, 0, 1.5)$$

$$\frac{\partial(P(0,1))}{\partial u} =$$

$$\frac{\partial(P(1,0))}{\partial u} =$$

$$\frac{\partial(P(1,1))}{\partial u} =$$

EXAMPLE OF CONSTRUCTING A COON'S PATCH

(y, v)

(x, u)

	2	3	4	5	6	7
7
8	.	.	10	9		
9	.	14	12	11	10	.
10	.	15	13	14	10	.
11	.	.	10	11		

- Gradients at the corners of the patch

$$P(0,0) = (9,4,12)$$

$$\frac{\partial(P(0,0))}{\partial v} = ((9, 5, 11) - (9, 3, 14))/2 = (0, 1, -1.5)$$

$$\frac{\partial(P(0,1))}{\partial v} =$$

$$\frac{\partial(P(1,0))}{\partial v} =$$

$$\frac{\partial(P(1,1))}{\partial v} =$$

EXAMPLE OF CONSTRUCTING A COON'S PATCH

- To find the bounding contours we use the cubic spline patch equation.

$$P(u, 0) = a_3u^3 + a_2u^2 + a_1u + a_0$$

$$a_0 = P_0 = (9, 4, 12)$$

$$a_1 = P'_0 = (1, 0, 1.5)$$

$$a_2 = -3P_0 - 2P'_0 - 3P_1 - P'_1$$

$$a_3 = 2P_0 + P'_0 - 2P_1 + P'_1$$

SURFACE REPRESENTATION

OBJECTIVE

- After completing this lecture, students will be able to
 - Derive mathematical expressions for
 - Bi-cubic surface
 - Surface sub-division
 - Geometric properties of surface

BICUBIC SURFACE PATCH

- Discussed so far simple surface patches from a conceptual point of view.
- Useful patch descriptions
- Boundary points are given
- Need to calculate the basis functions
- Steps:
 - Calculate for a fixed u , calculate for a fixed v
 - Add the two set of curves to generate the surface

BICUBIC SURFACE REPRESENTATION

1. Start with the cubic spline expression and derive for boundary conditions
2. Represent the curve as a function of the extreme points and the blending functions
3. For a fixed u , represent the curve
4. For a fixed v , represent the curve
5. Add them
6. Do the correction
7. Final expression

SUBDIVISION OF SURFACE PATCH

- To construct a surface patch with an interactive algorithm
- Changing the boundary (global) points doesn't change the shape
- Local control is required
- Add more points
- How many points we should add?
- Constraints ?

CONSTRAINT ON SURFACE SUBDIVISION

- The algorithm
 - Takes a surface defined by n points
 - Partitions it into several patches, so that
 - Together they represent the same surface
 - Each patch is now represented with n points
 - They are computed from the original set of points

BEZIER AND B-SPLINE SURFACE

OBJECTIVE

- After completing this lecture, students will be able to
 - Derive the mathematical expressions for
 - Bezier surface
 - B-spline surface
 - Construct Bezier surface
- Calculate expressions for
 - Surface subdivision
 - Degree elevation

BÉZIER CURVE

- Bézier equation

$$P(t) = \sum_{i=0}^n P_i B_i^n(t)$$

$$B_i^n(t) = \text{Bernstein Basis} = \binom{n}{i} t^i (1-t)^{n-i}$$

P_i = Control points

*The corresponding point of t on the new curve is obtained by **translating** the corresponding point of t on the original curve in the **direction of v with a distance $B_i^n(t)$***

BEZIER SURFACE

- $m + 1 \times n + 1$ point grid
- A rectangular Bezier patch is defined as:

$$Q(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_i^m(u) P_{i,j} B_j^n(v)$$

- In Matrix form this can be written as: $Q(u, v) = U N P N^T V^T$

$$U = [u^m, u^{m-1}, \dots, u, 1] \quad V = [v^n, v^{n-1}, \dots, v, 1] \quad N = ?$$

EXAMPLE



- Given the six three-dimensional points

$$\begin{array}{ccc} \mathbf{P}_{10} & \mathbf{P}_{11} & \mathbf{P}_{12} \\ \mathbf{P}_{00} & \mathbf{P}_{01} & \mathbf{P}_{02} \end{array}$$

- Problem:** Generate the corresponding Bezier Surface patch
- Step 1: Find the orders m and n of the surface
- Step 2: Calculate the weight functions $B_i^1(v), B_j^2(u)$
 - $i = 0,1$
 - $j = 0,1,2$
 - $m = 1, n = 2$


SCAFFOLDING CONSTRUCTION

- The scaffolding construction (or **de Casteljau algorithm**)
- Extension of the idea discussed in Bezier curves



SUBDIVIDING RECTANGULAR PATCHES

- To change the surface shape, if there are not enough points
- Just adding points is not a good solution because
 - this changes the shape of the surface
- A better solution is to subdivide the patch into four connected surface patches



Step 1: Select values for u and w

Step 2: Apply de Casteljau Algo (column wise)

Step 3: Apply de Casteljau Algo (row wise)

DEGREE ELEVATION

- In an application it is desired that all involved curves to have the same degree
- Increase the degree of a Bézier curve *without* changing its shape.
- Degree elevation
- Existing degree = n ; elevated degree = $n+1$
- Existing control points = $\{P_0, \dots, P_n\}$; New control points = $\{Q_0, \dots, Q_{n+1}\}$, such that $P_0 = Q_0$ & $P_n = Q_{n+1}$

$$Q_i = \frac{i}{n+1}P_{i-1} + \left(1 - \frac{i}{n+1}\right)P_i; 1 \leq i \leq n$$

DEGREE ELEVATION FOR SURFACE

- Extension of the idea discussed in case of curves
- Bezier patch of degree $m \times n$

- Definition:
$$Q(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_i^m(u) P_{i,j} B_j^n(v)$$

- Degree elevated patch is defined by $(m+2), (n+2)$ control points

$$\mathbf{Q}_{ij} = \left(\frac{i}{m+1}, 1 - \frac{i}{m+1} \right) \begin{bmatrix} \mathbf{P}_{i-1,j-1} & \mathbf{P}_{i-1,j} \\ \mathbf{P}_{i,j-1} & \mathbf{P}_{i,j} \end{bmatrix} \begin{bmatrix} \frac{j}{n+1} \\ 1 - \frac{j}{n+1} \end{bmatrix},$$

for $i = 0, 1, \dots, m+1$, and $j = 0, 1, \dots, n+1$.

EXAMPLE

- Starting with the 2x3 control points

$$\begin{array}{ccc} \mathbf{P}_{10} & \mathbf{P}_{11} & \mathbf{P}_{12} \\ \mathbf{P}_{00} & \mathbf{P}_{01} & \mathbf{P}_{02} \end{array}$$

- Elevation: 1x2 to 2x3

$$\begin{array}{ccc} \mathbf{P}_{10} & \mathbf{P}_{11} & \mathbf{P}_{12} \\ \mathbf{P}_{00} & \mathbf{P}_{01} & \mathbf{P}_{02} \end{array} \rightarrow \begin{array}{ccc} \mathbf{R}_{20} & \mathbf{R}_{21} & \mathbf{R}_{22} \\ \mathbf{R}_{10} & \mathbf{R}_{11} & \mathbf{R}_{12} \\ \mathbf{R}_{00} & \mathbf{R}_{01} & \mathbf{R}_{02} \end{array} \rightarrow \begin{array}{cccc} \mathbf{Q}_{20} & \mathbf{Q}_{21} & \mathbf{Q}_{22} & \mathbf{Q}_{23} \\ \mathbf{Q}_{10} & \mathbf{Q}_{11} & \mathbf{Q}_{12} & \mathbf{Q}_{13} \\ \mathbf{Q}_{00} & \mathbf{Q}_{01} & \mathbf{Q}_{02} & \mathbf{Q}_{03} \end{array}$$

B-SPLINE SURFACE

- B-Spline surface can be represented as:

$$Q(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{j+1}^{i+1} N_k^i(u) M_L^j(v)$$

$$N_1^i(u) = \begin{cases} 1 & \text{if } x_i \leq u \leq x_i + 1 \\ 0 & \text{otherwise} \end{cases}$$

$$N_k^i(u) = \frac{(u - x_i)N_{k-1}^i(u)}{x_{i+k-1} - x_i} + \frac{(x_{i+k} - u)N_{k-1}^{i+1}(u)}{x_{i+k} - x_{i+1}}$$



FLAME
UNIVERSITY

EVERLASTING
learning

THANK YOU