

## Tactics

- `rw[HYPOTHESIS]`: Substitutes the left hand side of an equality in `[HYPOTHESIS]` with the right hand side
- `rel [HYPOTHESIS]`: Substitutes the left hand side of an inequality in `[HYPOTHESIS]` with the right hand side
- `addarith[HYPOTHESIS]`: basic addition and subtraction with the provided hypothesis.
- `have [HYPOTHESIS_NAME] : [STATEMENT] := by [REASONING]`: creates a new hypothesis called `HYPOTHESIS_NAME` for future reference.
- `ring`: used as justification for things related to ring properties such as addition, subtract, multiplication (and sometimes division) with variables or ring elements
- `extra`: used as justification for when an inequality differs by a neutral positive value. Eg:  $y < 1 + y$   
:= by `extra`
- `cancel [ITEM] at [HYPOTHESIS]`: essentially divides both sides of an (in)equality by `[ITEM]` and rewrites the hypothesis. Eg: if `h1 : t^2 = t` then `cancel t at h1` would give `h1 : t = 1`
- `numbers`: Provides justification for any statement involving only numbers.
- `exact [HYPOTHESIS]`: Given `[HYPOTHESIS]` that matches the goal exactly, close the goal using `exact [HYPOTHESIS]`
- `apply [HYPOTHESIS]`: Used to choose a specific value for hypothesis' involving the universal quantifier.

## OR Goals and Hypothesis

- `obtain h1 | h2 := h3`: splits a hypothesis in the form  $x \vee y$  into two separate goals.
- `left`: Given a goal in the form  $a \vee b$ , chooses to prove the left side, `a`
- `right`: Given a goal in the form  $a \vee b$ , chooses to prove the right side `b`

## AND Goals and Hypothesis

- `obtain ⟨ h1 , h2 ⟩ := h3`: splits a hypothesis in the form  $x \wedge y$  into two separate hypothesis's
- `constructor`: Splits a goal involving AND into two separate goals to be closed sequentially

## Existential Quantifier Goals and Hypothesis

- `use [VALUE(S)]`: For goals involving existence, provided `[VALUE(S)]` are the value(s) for the quantities that we must prove the existence of.
- `obtain ⟨ x, hx ⟩ := h`: Creates a value and a hypothesis from another hypothesis that has the existential quantifier in it. Eg. if `h1 :  $\exists a : \mathbb{R}, a * t < 0$`  then, `obtain ⟨ x, hx ⟩ := h1` gives `hx :  $x * t < 0$`

## Lemmas

- `apply ne_of_lt`: Changes any goal in the form of  $x \neq y$  to  $x < y$  (usually paired with `apply ne_of_gt`)
- `apply ne_of_gt`: Changes any goal in the form of  $x \neq y$  to  $y < x$  (usually paired with `apply ne_of_lt`)
- `apply abs_le_of_sq_le_sq`: if  $x^2 \leq y^2$  and  $0 \leq y$ , then  $-y \leq x \wedge x \leq y$
- `apply le_antisymm`: Changes a goal in the form of  $a = b$  into two subgoals  $a \leq b$  and  $a \geq b$
- `rw[mul_eq_zero] at [HYPOTHESIS]`: If you have [HYPOTHESIS] in the form  $xy = 0$ , then `rw[mul_eq_zero] at [HYPOTHESIS]` gives  $[HYPOTHESIS : x = 0 \vee y = 0]$
- `Int.even_or_odd n`: Creates a hypothesis in the form  $\text{Even } n \vee \text{Odd } n$  that any integer  $n$  must be even or odd

## Definitions

- `dsimp[DEFINITION] at *`: rewrites all written occurrences of [DEFINITION] in terms of its mathematical definition.
- `Odd (n : ℤ): n = 2 * k + 1`
- `Even (n : ℤ): n = 2 * k`
- `( · | · )`: Definition of divisibility, used to unpack goals in the form  $a \mid b$