

Covering versus hitting in diversity embeddings: an affine obstruction and matching bounds

Jeff Kline

Version 0.1.0 — release candidate

Abstract

For every prime power q , we construct a unit-length hypergraph on $q^2 + 1$ vertices whose minimum ℓ_1 diversity distortion lies between $q^2/(2q - 1)$ and q , using q^2 supplies and $O(q^3 \log q)$ encoding bits. The square-root lower bound and its finite constant already follow from prior set-function work via exponentially many supplies; priority of the compact realization remains unresolved. Its encoding size contradicts Conjecture 1.7 of Jozefiak and Shepherd in arXiv:2303.04199v1, and the specified connector LP has the same integrality-gap lower bound. The proof compares covering cost with hitting probabilities through incidence moments. Conversely, established universal set-cover approximations give distortion at most $4\sqrt{n}$ for unit-cost pairwise-intersecting supply hypergraphs, even without a common root. The square-root exponent is tight in this class, including families with empty global intersection. Positive costs in this class give an upper bound $2\sqrt{nH_n}$, where H_n is the n th harmonic number. These upper bounds do not settle arbitrary hypergraphs. A constant-factor coverage approximation for a known greedy-bad family separates that algorithm's logarithmic loss from a diversity-embedding obstruction.

1 The covering–hitting principle

A diversity assigns a size to each finite set of points; its ℓ_1 model is the sum of coordinate ranges. Here the size of a terminal set is the minimum cost of a connected supply collection spanning it. A cut indicator asks only whether terminals occur on both sides of a cut. When cheap supplies and expensive demands have comparable crossing probabilities for every cut, no nonnegative cut mixture can approximate both costs well. This is the mechanism behind the example below; its affine incidence structure supplies a small explicit witness. Covering multiplicity alone is insufficient: the probability comparison must hold uniformly over cuts under the same supply and demand measures.

An existential square-root lower bound for diversities already follows from the set-function separation of Badanidiyuru et al. [7] by a rooted construction with exponentially many supplies, as explained in Section 9. The distinction here is the explicit polynomial-size supply representation, its encoding-size consequence, and the specified LP gap. The same prior proof also yields the finite constant $q^2/(2q - 1)$. Priority of the precise compact contribution remains unresolved.

This investigation grew out of range-cost multi-marginal earth moving. Kline [1] established geometric properties of its value, including convex-hull monotonicity and Minkowski additivity. The subsequent prefix-width study [2] expresses the value as the sum, over cuts between adjacent bins, of the largest cumulative mass minus the smallest. In cumulative coordinates this is precisely the coordinate-range ℓ_1 diversity. That connection led to the embedding question studied here and to examining what nonnegative cut sums can detect. The present obstruction uses the standard

cut representation directly; it does not require the later transport-rigidity theorem. These works provide the geometric starting point, while the covering–hitting argument supplies the obstruction.

The matching question has a complete answer for a substantial special class. If supply edges pairwise intersect, every cover is connected. We transfer established universal set-cover approximations to diversity embeddings on all terminal sets, obtaining the square-root exponent for unit costs. These upper bounds do not settle the corresponding question for arbitrary hypergraphs. In general, the ratio between minimum connected cost and minimum cover cost can grow without bound, so the transfer does not give a uniform approximation for arbitrary hypergraphs. Controlling that ratio or finding a different transfer is a separate obstacle from removing the weighted logarithm, and the general square-root direction of Bryant and Tupper [5] remains unresolved here.

2 Definitions and construction

For a connected hypergraph $G = (V, E)$ with nonnegative edge lengths ℓ_e , its *Steiner diversity* is

$$\delta_G(A) = \min_{J \in \mathcal{T}_A} \sum_{e \in J} \ell_e \quad (|A| \geq 2),$$

where $\mathcal{T}_A$ consists of edge subsets inducing a connected subhypergraph containing A . Connectivity is incidence connectivity; vertices outside A may be used. Set $\delta_G(A) = 0$ for $|A| \leq 1$. This is the convention in [3, Definition 1.6].

For a map $f : V \rightarrow \mathbb{R}^d$, the induced ℓ_1 diversity is the sum of coordinate ranges,

$$\Delta_f(A) = \sum_{i=1}^d \left(\max_{v \in A} f_i(v) - \min_{v \in A} f_i(v) \right),$$

with value zero on the empty set. After rescaling, distortion D means $\delta_G(A) \leq \Delta_f(A) \leq D\delta_G(A)$ for every $A \subseteq V$. There is no bound on d in the result below.

Fix a prime power q . Let $X = \mathbb{F}_q^2$ and $V = X \cup \{r\}$, where r is a new root. For $a, b \in \mathbb{F}_q$, put

$$L_{a,b} = \{(x, ax + b) : x \in \mathbb{F}_q\}, \quad e_{a,b} = \{r\} \cup L_{a,b}.$$

Take these q^2 rooted nonvertical lines as the supply edges, all of length one. Vertical lines are not supply edges. For $x \in \mathbb{F}_q$, define

$$Q_x = \{(x, y) : y \in \mathbb{F}_q\}, \quad A_x = \{r\} \cup Q_x.$$

These rooted columns will serve as the demand sets. Write δ_q for the resulting Steiner diversity. The hypergraph has rank $q + 1$ and $q^2(q + 1)$ incidences. Listing vertex identifiers and unit lengths takes $O(q^3 \log q)$ bits.

Every nonempty supply-edge collection is connected through r . Therefore, for $|A| \geq 2$, $\delta_q(A)$ is exactly the minimum number of nonvertical lines covering $A \cap X$. In particular,

$$\delta_q(e_{a,b}) = 1, \quad \delta_q(A_x) = q. \tag{1}$$

The second equality holds because each supply line meets a column in one point, and the q horizontal lines cover the entire plane.

3 An inequality for every cut

For $T \subseteq X$, let

$$c_T(A) = \mathbf{1}\{A \cap T \neq \emptyset, A \cap (V \setminus T) \neq \emptyset\}.$$

Every nontrivial cut can be oriented to have its root-free side in X . On a rooted set $\{r\} \cup B$, its value is simply $\mathbf{1}\{T \cap B \neq \emptyset\}$.

Lemma 1. *For every $T \subseteq X$, a uniformly chosen column Q and a uniformly chosen nonvertical affine line L satisfy*

$$\mathbb{P}(Q \cap T \neq \emptyset) \leq C_q \mathbb{P}(L \cap T \neq \emptyset), \quad C_q = \frac{2q-1}{q}.$$

Proof. Suppose $T \neq \emptyset$. Let $t_x = |T \cap Q_x|$, $t = \sum_x t_x$, and let h be the number of occupied columns. For $Z = |L \cap T|$, point and ordered-pair incidence counts give

$$\mathbb{E}Z = t/q, \quad \mathbb{E}Z^2 = t/q + (t^2 - \sum_x t_x^2)/q^2.$$

Distinct points in different columns lie on exactly one supply line; points in the same column cannot occur together. Cauchy–Schwarz gives $\mathbb{P}(Z > 0) \geq (\mathbb{E}Z)^2/\mathbb{E}Z^2$. Since column hitting has probability h/q , its ratio to line hitting is at most

$$\frac{h}{q} \left(\frac{q}{t} + 1 - \frac{\sum_x t_x^2}{t^2} \right) \leq \frac{h}{q} \left(\frac{q}{t} + 1 - \frac{1}{h} \right) \leq \frac{q+h-1}{q} \leq \frac{2q-1}{q}.$$

Here $\sum_x t_x^2 \geq t^2/h$, $t \geq h$, and $h \leq q$. For $T = \emptyset$, both probabilities vanish. $\square$

A finite ℓ_1 diversity is a nonnegative sum of cut diversities:

$$\Delta = \sum_{T \subseteq X} w_T c_T, \quad w_T \geq 0. \quad (2)$$

Indeed, sort the distinct values in each coordinate of an embedding. Each consecutive gap contributes its length times the cut separating values above that gap from those below it. These contributions telescope to the coordinate range on every set. Complement cuts containing r , then sum over coordinates. Conversely, weighted cut indicators give an ℓ_1 embedding. This is also the cut representation used in [6, 3].

Multiplying Lemma 1 by w_T and summing yields the central inequality, valid for every nonnegative cut sum:

$$\frac{1}{q} \sum_x \Delta(A_x) \leq \frac{C_q}{q^2} \sum_{a,b} \Delta(e_{a,b}). \quad (3)$$

4 The embedding obstruction

Theorem 2. *The minimum ℓ_1 diversity distortion of (V, δ_q) satisfies*

$$\frac{q^2}{2q-1} \leq D_{\min} \leq q.$$

The lower bound already follows from requiring domination on the q sets A_x and the upper bound $\Delta(e_{a,b}) \leq D$ on the q^2 supply edges.

Proof. By (1) and (3), these restricted constraints imply

$$q \leq \frac{1}{q} \sum_x \Delta(A_x) \leq \frac{C_q}{q^2} \sum_{a,b} \Delta(e_{a,b}) \leq C_q D.$$

For the upper bound, use the unit-weight singleton cuts $\Delta_0 = \sum_{p \in X} c_{\{p\}}$. For $|A| \geq 2$, put $s = |A \cap X|$. Then $\Delta_0(A) = s$. Each supply line covers at most q of these points, while choosing one line through each point gives a connected cover. Hence

$$s/q \leq \delta_q(A) \leq s, \quad \delta_q(A) \leq \Delta_0(A) \leq q\delta_q(A).$$

Both diversities vanish on empty sets and singletons. $\square$

Corollary 3. *Conjecture 1.7 of [3], in its arXiv v1 formulation, is false.*

Proof. That conjecture asks for distortion polylogarithmic in the encoding size of the inducing hypergraph. Here $B_q = O(q^3 \log q)$, while the distortion is at least $q/2$. Along unbounded primes, $q/2$ exceeds $C \log(2 + B_q)^k$ for every fixed $C > 0$ and $k \geq 0$. The obstruction concerns existence, independently of running time. $\square$

Since $n = q^2 + 1$, this family's minimum distortion is $\Theta(\sqrt{n})$. This is a statement about the explicit family; no matching upper bound for all diversities is claimed.

5 An explicit connector-LP gap

Give each supply edge capacity $c_{a,b} = 1/q^2$ and each demand edge A_x weight $b_x = 1/q$. These capacities are distinct from the unit lengths used to define δ_q . For a nontrivial cut with root-free side T , its ratio is

$$\phi(T) = \frac{q^{-2} \sum_{a,b} c_T(e_{a,b})}{q^{-1} \sum_x c_T(A_x)}.$$

The denominator is positive because nonempty T meets a column. Lemma 1 gives $\phi(T) \geq 1/C_q = q/(2q-1)$.

The connector relaxation [3, Equation (24)] specializes to

$$\begin{aligned} & \text{minimize} && q^{-2} \sum_{a,b} d_{a,b} \\ & \text{subject to} && q^{-1} \sum_x y_x \geq 1, \\ & && \sum_{e_{a,b} \in J} d_{a,b} \geq y_x \quad (x \in \mathbb{F}_q, J \in \mathcal{T}_{A_x}), \\ & && d_{a,b} \geq 0, \quad y_x \geq 0. \end{aligned} \tag{4}$$

For completeness, this is a relaxation of cut optimization. If a cut has total demand $h > 0$, set $d_e = c_T(e)/h$ and $y_x = c_T(A_x)/h$. A connected connector spanning a demand that crosses the cut must contain a crossing supply edge. Thus all connector inequalities hold, normalization is one, and the objective equals the cut ratio.

Theorem 4. *The optimum of (4) is exactly $1/q$. Its integrality gap on the specified instance is at least $q^2/(2q-1)$.*

Proof. Set $d_{a,b} = 1/q$ and $y_x = 1$. Every column connector contains at least q edges, so this is feasible and its objective is $1/q$. Conversely, for each fixed slope a , the q edges indexed by b form a connector for every column. Therefore

$$\sum_b d_{a,b} \geq y_x \quad \text{for every } x, \quad \sum_b d_{a,b} \geq \frac{1}{q} \sum_x y_x \geq 1.$$

Summing over all q slopes gives an objective of at least $1/q$. Since every cut has ratio at least $1/C_q$, the gap is at least $q/C_q = q^2/(2q-1)$. $\square$

All supply and demand incidences and rational weights have polynomial encoding size. The gap prevents a polylogarithmic rounding guarantee relative to this LP's value on general inputs. It does not establish hardness for all approximation algorithms for hypergraph sparsest cut.

6 A structural certificate and transversal families

Let a finite family $\mathcal{S}$ of nonempty subsets cover a finite set X . Adjoin a root r and use unit-cost supply edges $\{r\} \cup S$. Write $\tau(B)$ for the minimum number of supply sets covering B . Let μ be a full-support probability on supplies and ν a probability on nonempty demand subsets $B \subseteq X$. Define

$$p_x = \mathbb{P}_\mu(x \in S), \quad b_x = \mathbb{P}_\nu(x \in B), \quad K = \mathbb{E}_\nu \tau(B).$$

Theorem 5. Suppose $b_x \leq \gamma p_x$ for all x and, for every $T \subseteq X$, $Z_T = |S \cap T|$ satisfies

$$\mathbb{E} Z_T^2 \leq a \mathbb{E} Z_T + b (\mathbb{E} Z_T)^2, \quad a, b \geq 0.$$

Put $C = a\gamma + b > 0$. Then demand hitting is at most C times supply hitting for every T , and every ℓ_1 diversity embedding has distortion at least K/C . With capacities μ and rooted demand weights ν , the connector-LP gap is also at least K/C . A sufficient incidence condition is $\mathbb{P}_\mu(x, y \in S) \leq \beta p_x p_y$ for $x \neq y$, with $\beta \geq 0$; it gives $C = \gamma + \beta$.

Proof. For nonempty T , put $M = \mathbb{E} Z_T = \sum_{x \in T} p_x > 0$. The union bound and Cauchy-Schwarz bound the ratio of hitting probabilities by

$$\min(\gamma M, 1) \frac{a + bM}{M} \leq a\gamma + b.$$

Use the first branch if $\gamma M \leq 1$ and the second otherwise. Nonempty demands imply $\gamma > 0$. The empty test set is immediate. The pairwise condition gives $\mathbb{E} Z_T^2 \leq M + \beta M^2$ by expansion.

Orient cuts away from r and sum this hitting inequality with nonnegative cut weights. Domination on demands and the distortion bound on unit-cost supplies imply $K \leq CD$. For the LP, minimize $\sum_S \mu(S) d_S$ over $d_S, y_B \geq 0$, with normalization $\mathbb{E}_\nu y_B \geq 1$ and all inequalities $\sum_{S \in J} d_S \geq y_B$, where J covers B , equivalently connects $\{r\} \cup B$. The choice $d_S = 1/K$, $y_B = \tau(B)/K$ is feasible with objective $1/K$. Every positive-demand cut has ratio at least $1/C$. The LP optimum is positive: the full supply collection is a connector for every demand, so $\sum_S d_S \geq 1$ and the objective is at least $\min_S \mu(S) > 0$. This proves the gap claim. $\square$

Corollary 6. Partition X into $k \geq 1$ columns of $q \geq 2$ points. Suppose every supply selects one point per column, with uniform symbols and pairwise independent selections between distinct columns under μ . For uniform column demands, distortion and connector-LP gap are at least

$$\frac{kq}{k+q-1}.$$

If the supplies are a uniform indexed list partitioned into classes of q rows covering X , the LP optimum is $1/q$ and distortion is at most $\min(k, 2q) \leq \sqrt{2|X|}$.

Proof. Every column has covering cost q : each row covers one point, and a supported row exists through each point. For a test set with column occupancies t_i , total $t > 0$ and h occupied columns, the exact second moment is $t/q + (t^2 - \sum_i t_i^2)/q^2$. As in Lemma 1, the ratio of demand hitting to supply hitting is at most

$$\frac{h}{k} \left(\frac{q}{t} + 1 - \frac{\sum_i t_i^2}{t^2} \right) \leq \frac{q+h-1}{k} \leq \frac{q+k-1}{k}.$$

The cut and LP arguments of Theorem 5 now apply with $K = q$. Each resolution class connects every demand. Averaging its LP constraints gives total length at least one; summing classes gives objective at least $1/q$, attained by constant lengths $1/q$ and demands one.

The singleton point-cut sum gives distortion at most k , since a row covers at most k points. Resolution also gives $1 \leq \delta(A) \leq q$ on nonsingletons. A random fair-bit cut crosses such a set with probability between $1/2$ and one; multiplying its expectation by $2q$ gives distortion at most $2q$. Both cut sums vanish on empty sets and singletons. $\square$

These hypotheses have explicit finite-field supply. For a prime power q , an integer $d \geq 2$, and $1 \leq k \leq q^{d-1}$, choose distinct $v_1, \dots, v_k \in \mathbb{F}_q^{d-1}$ and take rows

$$S_{a,b} = \{(i, a + b \cdot v_i) : 1 \leq i \leq k\}, \quad (a, b) \in \mathbb{F}_q \times \mathbb{F}_q^{d-1}.$$

The coefficient vectors $(1, v_i), (1, v_j)$ are linearly independent for $i \neq j$, proving uniform pairwise independence. Fixing b and varying a gives a resolution class. Full evaluation domains have $k = q^{d-1}$, $q^d + 1$ rooted vertices and lower bound $q^d/(q^{d-1} + q - 1)$. Repeated indexed rows on smaller domains may be retained as copies. The balanced case $k = q$ recovers the affine obstruction.

Resolution alone is insufficient. The q constant rows $\{(i, a) : 1 \leq i \leq k\}$ form a resolution, but a test set equal to one row is hit by every column and only a $1/q$ fraction of supplies. Thus the comparison constant can grow with q .

7 Coverage approximation and matching upper bounds

Let $\mathcal{S}$ be a finite family of nonempty sets covering a set X of size $N \geq 1$, with positive costs $c(S)$. Let $f(B)$ be minimum cover cost. A *coverage function* is a nonnegative combination of hitting indicators. The following assignment bounds are established universal set-cover results [4, 13]; we include the elementary arguments to make the diversity transfer self-contained.

Lemma 7. *There is a coverage function g , supported on disjoint blocks partitioning X , with $f \leq g \leq \sqrt{NH_N}f$, where $H_N = \sum_{j=1}^N 1/j$. For unit costs the factor can be replaced by $2\sqrt{N}$.*

Proof. For positive costs, repeatedly choose a supply S_i minimizing $c(S_i)/\sqrt{|B_i|}$, where B_i is its nonempty residual set of unassigned points. Assign all of B_i and define $g(A) = \sum_i c(S_i) \mathbf{1}\{A \cap B_i \neq \emptyset\}$. The charged supplies cover A , so $f \leq g$. Fix a supply S of cost c . At a step whose block hits S , let a_i be the number of remaining points of S and $b_i = |B_i|$. Minimality gives $c(S_i) \leq c\sqrt{b_i/a_i}$. The a_i at hit steps strictly decrease, while all blocks are disjoint. Hence, summing only over hit steps,

$$g(S) \leq c \sum_i \sqrt{b_i/a_i} \leq c \sqrt{\left(\sum_i b_i\right) \left(\sum_i 1/a_i\right)} \leq c\sqrt{NH_N}.$$

Monotonicity and subadditivity of g , applied to an optimal cover of A , give $g(A) \leq \sqrt{NH_N}f(A)$.

For unit costs, set $t = \lceil \sqrt{N} \rceil$. Repeatedly assign all residual points of any supply having at least t of them. There are at most N/t such blocks. Assign each remaining point its own singleton block, associated with any incident supply, and give all blocks weight one. Again $f \leq g$. If R is the remaining set, every supply meets R in at most $t - 1$ points. An optimal cover gives $|A \cap R| \leq (t - 1)f(A)$. For nonempty A , $f(A) \geq 1$, and therefore

$$g(A) \leq N/t + |A \cap R| \leq (N/t + t - 1)f(A) \leq 2\sqrt{N}f(A).$$

The empty set has value zero throughout. $\square$

Lemma 8. *If $g(A) = \sum_T w_T \mathbf{1}\{A \cap T \neq \emptyset\}$ is a coverage function on X , there is a nonnegative cut sum Δ_0 on X such that $g(A) \leq \Delta_0(A) \leq 2g(A)$ for $|A| \geq 2$. If its nonempty atoms form a partition, at most $4N$ cut coordinates suffice.*

Proof. For each atom T , retain its points independently with probability $1/2$ to obtain a random cut side $R_T \subseteq T$. Put $\Delta_0 = 2 \sum_T w_T \mathbb{E} c_{R_T}$. A missed atom contributes zero. If A hits T and has a point outside T , crossing probability is $1 - 2^{-|A \cap T|}$. If $A \subseteq T$ and $|A| \geq 2$, it is $1 - 2^{1-|A|}$. Both probabilities lie in $[1/2, 1]$. Empty and singleton cut values are zero; no comparison to their cover values is asserted.

For the coordinate count, pairwise independent fair bits suffice. On an atom of size m , choose distinct nonzero vectors $v_x \in \mathbb{F}_2^d$, with $d = \lceil \log_2(m + 1) \rceil$, and retain x when $a \cdot v_x = 1$ for uniform $a \in \mathbb{F}_2^d$. An inside point and an outside terminal force a crossing with probability $1/2$; two distinct inside terminals have different bits with probability $1/2$. Thus the same bounds hold using at most $4m$ cuts. For a partition, summing atom sizes gives at most $4N$ coordinates. $\square$

Theorem 9. *Let a finite hypergraph on $n \geq 2$ vertices have nonempty supplies covering all vertices and positive costs. Let f be its minimum cover cost and δ its Steiner diversity. Suppose*

$$f(A) \leq \delta(A) \leq \kappa f(A) \quad (|A| \geq 2), \quad \kappa \geq 1.$$

Then its ℓ_1 diversity distortion is at most $2\kappa\sqrt{nH_n}$, or $4\kappa\sqrt{n}$ for unit costs, with at most $4n$ coordinates. In particular, pairwise-intersecting supplies have $\kappa = 1$, with no common vertex required.

Proof. Apply Lemmas 7 and 8 on the original vertex set. If the assignment factor is D , then on nonsingletons

$$\delta \leq \kappa f \leq \kappa \Delta_0 \leq 2\kappa g \leq 2\kappa Df \leq 2\kappa D\delta.$$

Both diversities vanish otherwise. Under pairwise intersection every nonempty cover is connected, so $f = \delta$ on nonsingletons. $\square$

For common-root supply on $X \cup \{r\}$, one can instead apply the assignment on X and regard its atoms as subsets of the larger vertex set. The same lift proof gives the slightly sharper bounds $4\sqrt{N}$ and $2\sqrt{NH_N}$, with $N = |X|$. It applies to all terminal sets, including those omitting r .

Corollary 10. *The square-root exponent is tight for unit-cost pairwise-intersecting supply hypergraphs, even when their global edge intersection is empty. If a unit-cost supply hypergraph has connected edge-intersection graph of diameter d , its distortion is at most $4 \max(1, d)\sqrt{n}$; in particular diameter two gives $8\sqrt{n}$.*

Proof. For the lower bound, replace the affine example's root by $R = \{r_1, r_2, r_3\}$. For every affine supply line L , use the three edges $L \cup (R \setminus \{r_i\})$, all of unit cost. The resulting $3q^2$ edges pairwise intersect, but no vertex belongs to all of them. There are $n = q^2 + 3$ vertices. On terminal subsets of $X \cup \{r_1\}$, this diversity equals δ_q : projecting a new cover to its affine lines gives one inequality, and replacing every old edge by $L \cup \{r_1, r_2\}$ gives the other. These covers are connected in both systems. Restricting any embedding therefore gives the lower bound $q^2/(2q - 1)$ from Theorem 2. Theorem 9 supplies the matching exponent along this family.

For the diameter claim with $d \geq 1$, connect every edge of a minimum k -edge cover to one chosen cover edge along paths of length at most d in the intersection graph. The connected union uses at most $k + (d - 1)(k - 1) \leq dk$ edges. Thus $\delta \leq df$, and Theorem 9 applies. Diameter zero means there is one supply edge covering all vertices, so $\kappa = 1$ works. This counting argument is specific to unit costs. $\square$

8 What the weighted logarithm does and does not show

The weighted assignment proof loses $\sqrt{H_n}$ through its harmonic Cauchy–Schwarz estimate. Ezra et al. [13, Appendix B, Proposition B.1] give a family attaining $\Omega(\sqrt{n \log n})$ for that greedy algorithm. Consequently a sharper analysis of the unchanged algorithm cannot establish $O(\sqrt{n})$ universally. That is not a lower bound against arbitrary coverage functions. Their bad family consists of disjoint blocks and one transversal, and the following elementary observation applies to all its positive choices of block costs.

Lemma 11. *Let B_i partition a finite set, choose $s_i \in B_i$, and let the supplies be B_i of cost $c_i > 0$ and $S = \{s_i\}_i$ of cost one. Its cover cost f has a coverage approximation of factor $1 + (1 - e^{-1})^{-1} < 2.583$. Its common-root diversity realization has distortion less than 5.165.*

Proof. For a query A , let $J = \{i : A \cap (B_i \setminus \{s_i\}) \neq \emptyset\}$, $F(A) = \sum_{i \in J} c_i$, and $u(A) = \sum_{i \notin J: s_i \in A} c_i$. Petal points force their block purchases, after which the remaining selected points cost either $u(A)$ or one. Thus $f(A) = F(A) + \min(1, u(A))$. Put $z(A) = \sum_{i: s_i \in A} c_i$ and $H(A) = \min(1, z(A))$. Then $f \leq F + H$, $F \leq f$, and $H \leq f$; for the last inequality, charge selected points in forced blocks to F .

Include each s_i in a random hitting set with probability $1 - e^{-c_i}$, independently. Its hitting probability on A is $1 - e^{-z(A)}$. With $\alpha = (1 - e^{-1})^{-1}$, the coverage function $Q(A) = \alpha(1 - e^{-z(A)})$ satisfies $H \leq Q \leq \alpha H$ by concavity on $[0, 1]$ and monotonicity above one. Hence $g = F + Q$ is coverage and $f \leq g \leq (1 + \alpha)f$. The common-root lift doubles the factor. The expected hitting sum is finite; no polynomial support bound is claimed. $\square$

The budgeted-additive-to-coverage factor $e/(e - 1)$ used here is established prior art [8]. The point is its application to the greedy-bad family: that family cannot establish a necessary logarithm for arbitrary coverage or its rooted diversity realization. Whether every positive-cost cover function admits $O(\sqrt{n})$ coverage approximation remains unresolved here. This includes all monotone subadditive functions f with $f(\emptyset) = 0$ and $f(S) > 0$ on nonempty sets: take every nonempty set S as a supply of cost $f(S)$; monotonicity and subadditivity prove that its cover cost is exactly f , though the representation can be exponential.

The non-rooted general question has a separate connectivity issue. Fixing a root in an arbitrary diversity and replacing $\delta(A)$ by $\delta(A \cup \{r\})$ can inflate a close pair far from that root without bound. The cover-cost comparison in Theorem 9 resolves this only under its stated hypothesis. Bryant and Tupper's general n upper bound and proposed square-root direction [5] therefore remain separate from the special-class matching result here. The small-query algorithmic hardness statements in [3, Section 5] do not rule out existential embeddings or algorithms reading explicit supply hypergraphs.

9 Related work and scope of the contribution

The construction uses established ideas from set functions and finite geometry. Badanidiyuru et al. [7, Appendix 6, Theorem 6.1] consider a partition into q groups of size q and the function $f(S) = \max_i |S \cap Q_i|$, obtaining a square-root separation from submodular approximations. Its direct unit-cost rooted representation uses all q^q full transversals. Indeed, every transversal covers at most one point per group, and $f(S)$ such transversals cover S ; adjoining a common root makes every cover connected. For an ℓ_1 embedding of this diversity with distortion D , the function $g(S) = \Delta(\{r\} \cup S)/D$ is coverage, hence submodular, and satisfies $f(S)/D \leq g(S) \leq f(S)$. Their theorem therefore already implies an existential $\Omega(\sqrt{n})$ lower bound for hypergraph Steiner diversities. This is a consequence of their set-function theorem, not a diversity claim stated in that source. Retaining the final inequality in their proof even gives $D \geq q^2/(2q-1)$ before its rounding to $q/2$.

The affine construction replaces q^q full transversals by q^2 supplies and proves the required all-cut comparison for that smaller family. The dense predecessor does not itself contradict a polylogarithmic bound in hypergraph encoding size. Thus neither the existential square-root growth nor its numerical constant is claimed as new here; the precise compact realization and its encoding-size and connector-LP consequences are the contribution whose priority remains to be settled.

Coverage approximation is studied by Devanur et al. [8]. Affine-plane set systems also appear in the streaming lower bounds of Emek and Rosén [9] and Chakrabarti and Wirth [10]. Those streaming statements impose information restrictions; the inequality (3) constrains every cut mixture, with unrestricted support. The affine sample space $(a, b) \mapsto (ax + b)_x$ is the standard pairwise-independent construction of [12, Sections 2.1–2.2, Claim 2.1]. Recent work of Bryant and Tupper [11] connects rooted set functions, strong submodularity and geometric diversity representations. Universal set-cover and disjoint-service lower bounds also need care: they restrict the approximating representation. For example, the function $h(S) = \lceil |S|/q \rceil$ used in [13, Theorem 4.5] has the coverage approximant $g(S) = \mathbf{1}\{S \neq \emptyset\} + |S|/q$ with $h \leq g \leq 2h$. A lower bound for disjoint services therefore does not automatically give an arbitrary-coverage obstruction. These antecedents should be distinguished from the polynomial-encoding counterexample and explicit connector-LP instance established here.

Status. The literature review found no direct antecedent for the precise compact construction and consequence, but it does not establish priority. The conjecture comparison is explicitly to arXiv:2303.04199v1; a reported journal revision has not been checked. This release candidate records the proof without a claim of being the first such result.

Computational assistance. Codex agents assisted with the construction, literature review, proof checking and drafting. The argument above is self-contained and does not rely on a numerical experiment. Agent reviews are not external peer review.

References

- [1] J. Kline. Properties of the d -dimensional earth mover’s problem. *Discrete Applied Mathematics* 265, 128–141, 2019. doi:10.1016/j.dam.2019.02.042.
- [2] J. Kline. *Prefix-width rigidity and dual geometry of multi-marginal earth moving*. Version 0.1.0, 2026. doi:10.5281/zenodo.22675435.
- [3] A. D. Jozefiak and F. B. Shepherd. *Diversity Embeddings and the Hypergraph Sparsest Cut*. 2023. arXiv:2303.04199v1.
- [4] L. Jia, G. Lin, G. Noubir, R. Rajaraman and R. Sundaram. *Universal Approximations for TSP, Steiner Tree, and Set Cover*. STOC 2005, 386–395. Author manuscript.
- [5] D. Bryant and P. Tupper. *Open Problem Statement: Minimal Distortion Embeddings of Diversities in ℓ_1* . 2017. arXiv:1712.01960v1.
- [6] D. Bryant and P. F. Tupper. Diversities and the geometry of hypergraphs. *Discrete Mathematics & Theoretical Computer Science* 16(2), 1–20, 2014. Published article.
- [7] A. Badanidiyuru, S. Dobzinski, H. Fu, R. Kleinberg, N. Nisan and T. Roughgarden. *Sketching Valuation Functions*. SODA 2012, 1025–1035. Appendix reference uses the October 2011 author manuscript.
- [8] N. R. Devanur, S. Dughmi, R. Schwartz, A. Sharma and M. Singh. *On the Approximation of Submodular Functions*. 2013. arXiv:1304.4948.
- [9] Y. Emek and A. Rosén. *Semi-Streaming Set Cover*. Author manuscript, August 2016; preliminary version, ICALP 2014.
- [10] A. Chakrabarti and A. Wirth. *Incidence Geometries and the Pass Complexity of Semi-Streaming Set Cover*. ECCC Report TR15-113, 2015.
- [11] D. Bryant and P. Tupper. *Submodular and strongly submodular functions and diversities*. 2026. arXiv:2608.11468v1.
- [12] M. Luby and A. Wigderson. *Pairwise Independence and Derandomization*. Foundations and Trends in Theoretical Computer Science 1(4), 237–301, 2005. Author PDF.
- [13] T. Ezra, S. Leonardi, M. Pawłowski, M. Russo and S. W. Umboh. *Universal Optimization for Non-Clairvoyant Subadditive Joint Replenishment*. 2024. Author PDF.