

**Example: Exercise 4.13**

$SPACE$	$T_1$	$T_2$	REGULAR	NORMAL
$R_{std}$	✓	✓	✓	✓
$R_{std}^n$	✓	✓	✓	✓
indiscrete	✗	✗	✗	✗
discrete	✓	✓	✓	✓
cofinite	✓	$X$ finite: ✓ $X$ infinite: ✗	$X$ finite: ✓ $X$ infinite: ✗	$X$ finite: ✓ $X$ infinite: ✗
cocountable	✓	$X$ countable: ✓ $X$ uncountable: ✗	$X$ countable: ✓ $X$ uncountable: ✗	$X$ countable: ✓ $X$ uncountable: ✗
$R_{LL}$	✓	✓	✓	✓
$R_{+00}$	✓	✗	✗	✗
LOS	✓	✓	✓	✓

$R$  and  $\mathbb{R}^n$

Since there is a finite distance between points and closed sets (not containing those points), there is always room for enclosing disjoint balls.

indiscrete

Since the only non-empty set is the entire space, there is no separation.

discrete

Since all disjoint subsets are both open and closed, they are self-enclosed.

cofinite/cocountable

First note that all finite sets are closed. Thus, single points can be viewed as closed sets. So assume  $p$  and  $q$  are distinct points in  $X$ . This means that  $X - \{p\}$  and  $X - \{q\}$  are open. Furthermore,  $p \in X - \{q\}$  but  $p \notin X - \{p\}$  and  $q \in X - \{p\}$  but  $q \notin X - \{q\}$ . Thus, cofinite/cocountable is  $T_1$ .

Now assume that there exists disjoint  $U, V \in \mathcal{T}$ . This means that  $X - U$  and  $X - V$  are finite/countable and since  $U \cap V = \emptyset$  it is the case that  $X - (U \cap V) = (X - U) \cup (X - V) = X$

and hence  $X$  is finite/countable. When  $X$  is finite/countable, all subsets are both open and closed, equivalent to the discrete topology, and so cofinite and cocountable are  $T_2$ , regular, and normal. However, if  $X$  is infinite/uncountable then open sets will always intersect and so cofinite and countable are neither  $T_2$ , regular, nor normal.

$\mathbb{R}_{LL}$

Since  $R_{LL}$  is finer than  $\mathbb{R}$ , it has the same separation properties.

$\mathbb{R}_{+00}$

Any two points can be  $T_1$  separated using the basis elements; however, if one point or closed set contains  $0'$  and the other point or closed set contains  $0''$  then there is always overlap between the two containing basis elements.

Lexigraphically Ordered Square

Use the alternate definitions. For any point  $p \in X$ , there exists some containing open set (strip), and it is always possible to use a smaller strip whose closure is contained in the original strip. For any closed set  $A \in X$ ,  $X - A$  is an enclosing open set, and likewise, a smaller open set with contained closure is possible.

#### **Theorem: 4.16**

$X, Y$  are  $T_2 \implies X \times Y$  is  $T_2$ .

*Proof.* Assume that  $X$  and  $Y$  are  $T_2$  and assume  $p_1, p_2 \in X \times Y$  where  $p_1 = (x_1, y_1)$  and  $p_2 = (x_2, y_2)$ . Since  $X$  is  $T_2$ , there exists  $U_1, U_2 \in \mathcal{T}_X$  such that  $x_1 \in U_1$  and  $x_2 \in U_2$  and  $U_1 \cap U_2 = \emptyset$ . Likewise, since  $Y$  is  $T_2$ , there exists  $V_1, V_2 \in \mathcal{T}_Y$  such that  $y_1 \in V_1$  and  $y_2 \in V_2$  and  $V_1 \cap V_2 = \emptyset$ . So  $p_1 \in U_1 \times V_1$  and  $p_2 \in U_2 \times V_2$ . Furthermore,  $U_1 \times V_1, U_2 \times V_2 \in \mathcal{T}_{X \times Y}$  and

$$(U_1 \times V_1) \cap (U_2 \times V_2) = (U_1 \cap U_2) \times (V_1 \cap V_2) = \emptyset$$

Therefore  $X \times Y$  is  $T_2$ . ■

#### **Lemma**

Let  $X$  and  $Y$  be topological spaces and let  $A \subset X$  and  $B \subset Y$ :

$$\overline{A \times B} = \bar{A} \times \bar{B}$$

*Proof.* Assume that  $p \in \overline{A \times B}$ . This means that for all  $U \in \mathcal{T}_{X \times B}$  such that  $p \in U$ :

$$U \cap (A \times B) \neq \emptyset$$

Now assume  $U_1 \in \mathcal{T}_X$  and  $U_2 \in \mathcal{T}_Y$  such that  $p \in U_1 \times U_2 \in \mathcal{T}_{A \times B}$ . Then it must be the case that  $(U_1 \times U_2) \cap (A \times B) = (U_1 \cap A) \times (U_2 \cap B) \neq \emptyset$ . This is only possible if  $U_1 \cap A \neq \emptyset$  and  $U_2 \cap B \neq \emptyset$ .

Therefore  $p \in \bar{A} \times \bar{B}$ .

Assume that  $p \in \bar{A} \times \bar{B}$ . This means that for all  $U_1 \in \mathcal{T}_X$  and  $U_2 \in \mathcal{T}_Y$  such that  $p \in U_1 \times U_2$ :

$$(U_1 \cap A) \times (U_2 \cap B) \neq \emptyset$$

Now assume  $U \in \mathcal{T}_{A \times B}$  such that  $p \in U \in \mathcal{T}_{A \times B}$ . Then there exists  $U_1 \in \mathcal{T}_X$  and  $U_2 \in \mathcal{T}_Y$  such that  $p \in U_1 \times U_2 = U$ . So it must be the case that:

$$U \cap (A \times B) = (U_1 \times U_2) \cap (A \times B) = (U_1 \cap A) \times (U_2 \cap B) \neq \emptyset$$

Therefore  $p \in \overline{A \times B}$ . ■

#### **Theorem: 4.17**

$X, Y$  are regular  $\implies X \times Y$  is regular.

*Proof.* Assume that  $X$  and  $Y$  are regular and assume  $p \in X \times Y$  and  $U \in \mathcal{U}_p$ . Then there exists  $U_1 \in \mathcal{T}_X$  and  $U_2 \in \mathcal{T}_Y$  such that  $p \in U_1 \times U_2 \subset U$ . Now, since  $X$  and  $Y$  are regular, there exists  $V_1 \in \mathcal{T}_X$  and  $V_2 \in \mathcal{T}_Y$  such that  $p \in V_1 \times V_2$ ,  $V_1 \subset \bar{V}_1 \subset U_1$ , and  $V_2 \subset \bar{V}_2 \subset U_2$ . Furthermore, since  $\bar{V}_1$  is closed in  $X$  and  $\bar{V}_2$  is closed in  $Y$ ,  $\bar{V}_1 \times \bar{V}_2$  (and hence  $\overline{V_1 \times V_2}$ ) is closed in  $X \times Y$ . And so:

$$p \in V_1 \times V_2 \subset \overline{V_1 \times V_2} = \bar{V}_1 \times \bar{V}_2 \subset U_1 \times U_2$$

Therefore  $X \times Y$  is regular. ■

#### **Theorem: 4.19**

Every  $T_2$  space is hereditarily  $T_2$ .

*Proof.* Assume that  $X$  is a  $T_2$  topological space and assume that  $Y \subset X$ . Now assume that  $a, b \in Y$ . Thus  $a, b \in X$  and, since  $X$  is  $T_2$ , there exists  $U, V \in \mathcal{T}_X$  such that  $a \in U, b \in V$ , and  $U \cap V = \emptyset$ . Furthermore,  $a \in U \cap Y \in \mathcal{T}_Y$  and  $b \in V \cap Y \in \mathcal{T}_Y$ . And so:

$$(Y \cap U) \cap (Y \cap V) = Y \cap (U \cap V) = Y \cap \emptyset = \emptyset$$

Therefore  $Y$  is also  $T_2$ . ■

#### **Theorem: 4.20**

Every regular space is hereditarily regular.

*Proof.* Assume that  $X$  is a regular topological space and assume that  $Y \subset X$ . Assume that  $p \in Y$ . This means that there exists some  $U_Y \in \mathcal{T}_Y$  such that  $p \in U_Y$ , and hence there exists  $U_X \in \mathcal{T}_X$  such that  $U_Y = U_X \cap Y$  and so  $p \in U_X$ . Now, since  $X$  is regular, there exists  $V_X \in \mathcal{T}_X$  such that  $p \in V_X \subset \bar{V}_X \subset U_X$ , and hence  $p \in V_X \cap Y = V_Y \in \mathcal{T}_Y$ . Furthermore, since  $\bar{V}_X$

is closed in  $X$ ,  $\overline{V_X} \cap Y = W_Y$  is closed in  $Y$ . Finally, since  $\overline{V_Y}$  is the smallest closed set in  $Y$  containing  $V_Y$ :

$$p \in V_Y \subset \overline{V_Y} \subset W_Y \subset U_Y$$

Therefore  $Y$  is regular. ■

### **Lemma**

Let  $X$  be a normal topological space and let  $Y \subset X$  such that  $Y$  is closed in  $X$ . For all  $A \subset Y$ , if  $A$  is closed in  $Y$  then  $A$  is closed in  $X$ .

*Proof.* Assume  $A \subset Y$  such that  $A$  is closed in  $Y$ . This means that  $Y - A \in \mathcal{T}_Y$ , and so there exists  $W \in \mathcal{T}_X$  such that  $W \cap Y = Y - A$ . Furthermore,  $X - W$  is closed in  $X$ . Now:

$$(X - W) \cap Y = (X \cap Y) - (W \cap Y) = Y - (Y - A) = A$$

But  $X - W$  and  $Y$  are closed in  $X$  and therefore  $A$  is also closed in  $X$ . ■

### **Theorem: 4.23**

Let  $X$  be a normal topological space and let  $Y \subset X$  such that  $Y$  is closed in  $X$ .  $Y$  is normal when given the relative topology.

*Proof.* Assume  $A, B \subset Y$  such that  $A$  and  $B$  are closed in  $Y$  and  $A \cap B = \emptyset$ . This means that  $A$  and  $B$  are also closed in  $X$ . Since  $X$  is normal, there exists  $U, V \in \mathcal{T}_X$  such that  $A \subset U, B \subset V$ , and  $U \cap V = \emptyset$ . Finally, since  $A \subset (U \cap Y) \in \mathcal{T}_Y$  and  $B \subset (V \cap Y) \in \mathcal{T}_Y$ :

$$(U \cap Y) \cap (V \cap Y) = (U \cap V) \cap Y = \emptyset \cap Y = \emptyset$$

Therefore  $Y$  is normal. ■