

Quadratic Equations

Definition

The general form of a quadratic equation is given by:

$$ax^2 + bx + c = 0$$

where $a \neq 0$

Note that if $a = 0$ we have a linear equation.

Before we tackle such equations, we need to be clear on two points:

1). $|x| = c$

Ask for a value whose absolute value is c . From either the definition: the high school one or the notion of distance, $x = \pm c$.

$$|x| = 2$$

$$x = \pm 2$$

$$|x| = 0$$

$$x = 0$$

$$|x| = -1$$

no solutions

2). $a = b \iff a^r = b^r?$

It depends! Don't do it without thinking!

3). $x^n = c$, n odd

$$x^3 = 8$$

$$(x^3)^{\frac{1}{3}} = 8^{\frac{1}{3}}$$

$$x = 2$$

$$x^3 = -8$$

$$(x^3)^{\frac{1}{3}} = (-8)^{\frac{1}{3}}$$

$$x = -2$$

4). $x^n = c$, n even

$$x^2 = 4$$

$$(x^2)^{\frac{1}{2}} = 4^{\frac{1}{2}}$$

$$|x| = 2$$

$$|x| = \pm 2$$

$$x^2 = -4$$

$$(x^2)^{\frac{1}{2}} = (-4)^{\frac{1}{2}}$$

no solution in \mathbb{R}

Don't mix up \sqrt{a} , which asks for the principle (positive) root, compared to $x^2 = c$, asking for solutions - they are two different questions!

$$5). (x + d)^2 = c$$

$$(x + 1)^2 = 4$$

$$|x + 1| = 2$$

$$x + 1 = \pm 2$$

$$x + 1 = 2 \text{ and } x + 1 = -2$$

$$x = 1, -3$$

Case 1: $b = 0$

This is the case above:

$$2x^2 - 8 = 0$$

$$2x^2 = 8$$

$$x^2 = 4$$

$$(x^2)^{\frac{1}{2}} = 4^{\frac{1}{2}}$$

$$|x| = 2$$

$$x = \pm 2$$

Case 2: $b \neq 0$

If we can factor by inspection, then we get an easy answer:

Two solutions:

$$x^2 - 5x + 6 = 0$$

$$(x - 2)(x - 3) = 0$$

$$x = 2, 3$$

Repeated solution:

$$x^2 + 2x + 1 = 0$$

$$(x + 1)^2 = 0$$

$$x = -1, -1$$

Otherwise, complete the square!

Start with $a = 1$:

$$x^2 + bx$$

Want to know what constant to add so that it can be rewritten as $(x + d)^2$, because once we do that, we can use the rules above to get at the variable.

$$x^2 + bx + c = (x + d)^2 = x^2 + 2d + d^2$$

So $c = d^2$ and $d = \frac{b}{2}$, so divide by 2 and then square.

Example

$$x^2 + 6x$$

$$b = 6$$

$$d = \frac{6}{2} = 3 \quad c = 3^2 = 9 \quad x^2 + 6x + 9 = (x + 3)^2$$

$$x^2 - 2x$$

$$b = -2$$

$$d = \frac{-2}{2} = -1$$

$$c = (-1)^2 = 1$$

$$x^2 - 2x + 1 = (x - 1)^2$$

$$x^2 - x$$

$$b = -1$$

$$d = -\frac{1}{2}$$

$$c = \left(-\frac{1}{2}\right)^2 = \frac{1}{4}$$

$$x^2 - x + \frac{1}{4} = \left(x - \frac{1}{2}\right)^2$$

When $a \neq 1$, divide it out first:

Example

$$2x^2 + 3x = 2 \left(x^2 + \frac{3}{2}x\right)$$

$$b = \frac{3}{2}$$

$$d = \frac{3}{4}$$

$$c = \frac{9}{16}$$

$$2 \left(x^2 + \frac{3}{2}x + \frac{9}{16}\right) = 2 \left(x + \frac{3}{4}\right)^2$$

$$2x^2 + 4x - 3 = 0$$

$$2x^2 + 4x = 3$$

$$2(x^2 + 2x) = 3$$

$$x^2 + 2x = \frac{3}{2}$$

$$x^2 + 2x + 1 = \frac{3}{2} + 1$$

$$(x + 1)^2 = \frac{5}{2}$$

$$|x + 1| = \sqrt{\frac{5}{2}}$$

$$x + 1 = \pm \sqrt{\frac{5}{2}}$$

$$x = -1 \pm \sqrt{\frac{5}{2}}$$

$$ax^2 + bx + c = 0$$

$$ax^2 + bx = -c$$

$$x^2 + \frac{b}{a}x = -\frac{c}{a}$$

$$x^2 + \frac{b}{a}x + \frac{b^2}{4a^2} = -\frac{c}{a} + \frac{b^2}{4a^2}$$

$$\left(x + \frac{b}{2a}\right)^2 = \frac{b^2 - 4ac}{4a^2}$$

$$x + \frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{2a}$$

$$x = -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

And viola! We have the quadratic formula! So, we can solve quadratics in one of three ways (in order of desiredness):

- 1). Inspection
- 2). Completing the square
- 3). Quadratic formula

Definition

The discriminant of the quadratic equation $ax^2 + bx + c = 0$ is given by:

$$D = b^2 - 4ac$$

4 cases:

- 1). $D > 0$ and a perfect square:

Two rational solutions (probably by inspection):

$$x^2 + 4x + 3$$

$$a = 1, b = 4, c = 3$$

$$x = \frac{-4 \pm \sqrt{4^2 - 4(1)(3)}}{2(1)} = \frac{-4 \pm 2}{2} = -1, -3$$

Note that this provides a factoring: $(x - r_1)(x - r_2) = 0$

$$(x + 1)(x + 3) = 0$$

Another example:

$$3x^2 + 10x + 3$$

$$a = 3, b = 10, c = 3$$

$$x = \frac{-10 \pm \sqrt{10^2 - 4(3)(3)}}{2(3)} = \frac{-10 \pm 8}{6} = -3, -\frac{1}{3}$$

$$\left(x + \frac{1}{3}\right)(x + 3) = 0$$

But if you foil this, you don't quite get the original, need to adjust so that the F works:

$$3\left(x + \frac{1}{3}\right)(x + 3) = 3(0)$$

$$(3x + 1)(x + 3) = 3(0)$$

which is what we would get if we factored by inspection.

2). $D > 0$ and not a perfect square:

Two matching irrational roots, like the previous example.

$$2x^2 + 4x - 3 = 0$$

$$x = -1 \pm \sqrt{\frac{5}{2}}$$

$$[x - (-1 + \sqrt{\frac{5}{2}})][x - (-1 - \sqrt{\frac{5}{2}})] = 0$$

$$[2x - 2(-1 + \sqrt{\frac{5}{2}})][x - (-1 - \sqrt{\frac{5}{2}})] = 0$$

$$[\sqrt{2}x - (-\sqrt{2} + \sqrt{5})][\sqrt{2}x - (-\sqrt{2} - \sqrt{5})] = 0$$

3). $D = 0$

One repeated rational root:

$$x^2 - 2x + 1 = 0$$

$$a = 1, b = 2, c = 1$$

$$x = \frac{2 \pm \sqrt{(-2)^2 - 4(1)(1)}}{2(1)} = \frac{2}{2} = 1$$

$$(x - 1)^2 = 0$$

4). $D < 0$

No real solutions:

$$x^2 + x + 1 = 0$$

$$a = 1, b = 1, c = 1$$

$$1^2 - 4(1)(1) = -3$$

$$x^2 + x = -1$$

$$x^2 + x + \frac{1}{4} = -\frac{3}{4}$$

$$\left(x + \frac{1}{2}\right)^2 = -\frac{3}{4}$$

stuck!