

V. B. Berestetskii E. M. Lifshitz L. P. Pitaevskii

Quantum Electrodynamics

Course of Theoretical Physics
Volume IV

New English translation from the fourth corrected Russian edition

Contents

Introduction	1
1 Uncertainty Relations in the Relativistic Domain	1
I Photon	5
2 Quantization of the Free Electromagnetic Field	5
3 Photons	9
4 Gauge invariance	12
5 The electromagnetic field in quantum theory	13
6 Angular momentum and parity of the photon	15
7 Spherical waves of photons	17
8 Polarization of the photon	22
9 The two-photon system	27
II Bosons	31
10 Wave equation for particles with spin 0	31
11 Particles and antiparticles	35
12 Strictly neutral particles	38
13 The transformations C, P, T	41
14 Wave equation for spin-1 particles	47
15 Wave equation for particles with higher integer spins	50
16 Helicity states of a particle ¹	52
III Fermions	59
17 Four-dimensional spinors	59
18 Relation of spinors to 4-vectors	61
19 Inversion of spinors	65
20 The Dirac equation in spinor representation	69
21 Symmetric form of the Dirac equation	71
22 Algebra of Dirac matrices	76
23 Plane waves	79
24 Spherical waves	82
25 The connection between spin and statistics	85

26	Charge conjugation and time reversal of spinors	88
27	Internal symmetry of particles and antiparticles	93
28	Bilinear forms	95
29	Polarisation density matrix	99
30	Two-component fermions	103
31	Wave equation for a spin-3/2 particle	107
IV	A Particle in an External Field	109
32	The Dirac equation for an electron in an external field	109
33	Expansion in powers of $1/c$	113
34	Fine structure of hydrogen atom levels	116
35	Motion in a centrally symmetric field	118
V	Radiation	123
VI	Scattering of Light	125
VII	The Scattering Matrix	127
VIII	Invariant Perturbation Theory	129
IX	Interaction of Electrons	131
X	Interaction of Electrons with Photons	133
XI	Exact Propagators and Vertex Parts	135
XII	Radiative Corrections	137
XIII	Asymptotic Formulae of Quantum Electrodynamics	139
XIV	Electrodynamics of Hadrons	141

Introduction

§ 1. Uncertainty Relations in the Relativistic Domain

The quantum theory presented in Vol. III of this course is non-relativistic in its very substance, and it fails for phenomena in which motion occurs at speeds no longer small next to the speed of light. One might at first imagine that a relativistic theory could be reached by generalizing the formalism of non-relativistic quantum mechanics in a more or less straightforward manner. A closer look shows, however, that a logically complete relativistic theory cannot be built without invoking fresh physical principles.

We begin by recalling several physical notions on which non-relativistic quantum mechanics rests (Vol. III, § 1). A fundamental part was played there, as we saw, by the concept of measurement — the process in which a quantum system interacts with a “classical object” (an “apparatus”) and thereby comes to possess definite values of particular dynamical variables (coordinates, velocities, and so forth). We saw, too, that quantum mechanics places severe limits on the extent to which different dynamical variables of an electron² can exist together. Thus the uncertainties Δq and Δp with which coordinate and momentum can coexist obey the relation $\Delta q \Delta p \sim \hbar$;³ the more precisely either of these quantities is measured, the less precisely the other can be known at the same moment.

What matters, however, is that any single dynamical variable of the electron, on its own, admitted measurement of unlimited precision, and within a time interval that could be taken as short as desired. On this circumstance the whole of non-relativistic quantum mechanics depends in a fundamental way. It alone makes it possible to introduce the wave function, the central object of the theory’s formalism. The physical content of the wave function $\psi(q)$ is, after all, that its squared modulus gives the probability that a measurement performed at a given instant will yield one or another value of the electron’s coordinate. Evidently the very notion of such a probability presupposes that a coordinate measurement of unlimited precision and speed can be realized in principle; failing that, the notion would be empty of content and would forfeit its physical meaning.

The existence of a limiting speed (the speed of light c) leads to new limitations, of a fundamental kind, on the possibilities of measuring various physical quantities (L. D. Landau, R. Peierls, 1930).

In Vol. III, § 44 the relation

$$(v' - v) \Delta p \Delta t \sim \hbar \tag{1.1}$$

was obtained; it relates the uncertainty Δp of a momentum measurement on the electron to the length Δt of the measuring process itself, v and v' denoting the electron’s velocity before and after that process. It follows that momentum can be measured accurately within a short time (small Δp together with small Δt) only if the measurement itself

²As in Vol. III, § 1, we say “electron” for brevity, understanding by it any quantum system.

³Ordinary units are used in this section.

alters the velocity strongly enough. In the non-relativistic theory this expresses the fact that a momentum measurement repeated after a short interval will not reproduce its result, but it places no restriction of principle on how accurate a single measurement of the momentum may be, the difference $v' - v$ being capable of unlimited increase.

The presence of a limiting speed, however, changes the state of affairs radically. The difference $v' - v$, like the velocities themselves, can now not exceed c (more precisely, $2c$). Replacing $v' - v$ in (1.1) by c , we obtain the relation

$$\Delta p \Delta t \sim \hbar/c, \quad (1.2)$$

which determines the best accuracy of momentum measurement attainable in principle for a given duration Δt of the measurement. Thus in the relativistic theory an arbitrarily accurate and arbitrarily rapid measurement of the momentum turns out to be impossible in principle. An exact measurement of the momentum ($\Delta p \rightarrow 0$) is possible only in the limit of an infinitely long duration of the measurement.

One has reason to expect that the measurability of the electron's coordinate as such is likewise affected. Within the mathematical formalism of the theory this appears as a conflict between an exact coordinate measurement and the statement that a free particle's energy is positive. As we shall see, the full set of eigenfunctions of a free particle's relativistic wave equation contains, besides the solutions whose time dependence is the "proper" one, solutions of "negative frequency" as well. These will in general appear also in the expansion of a wave packet describing an electron confined to a small region of space.

As will be shown, the "negative-frequency" wave functions are tied to the existence of antiparticles — the positrons. Their presence in the expansion of the wave packet reflects the creation of electron–positron pairs, unavoidable in the general case, during a measurement of the electron's coordinates. Since the emergence of these new particles cannot be controlled by the measuring process itself, coordinate measurement for the electron is drained of meaning.

In the rest frame of the electron the minimum error in a measurement of its coordinates is

$$\Delta q \sim \hbar/mc. \quad (1.3)$$

This value (the only one that dimensional considerations would permit) answers to a momentum uncertainty $\Delta p \sim mc$, which corresponds in turn to the minimum threshold energy needed to create a pair.

In a reference frame where the electron moves with energy ε , (1.3) is replaced by

$$\Delta q \sim c\hbar/\varepsilon. \quad (1.4)$$

In the extreme ultra-relativistic limit, in particular, energy and momentum are related by $\varepsilon \approx cp$, so that

$$\Delta q \sim \hbar/p, \quad (1.5)$$

the error Δq then coinciding with the particle's de Broglie wavelength.⁴

⁴What is meant here are measurements any outcome of which permits an inference about the state of

For photons the ultra-relativistic case always holds, so that expression (1.5) is valid. This means that it makes sense to speak of the coordinates of a photon only in those cases where the characteristic dimensions are large compared with the wavelength. But this is nothing other than the “classical” limiting case, corresponding to geometrical optics, in which one may speak of the propagation of light along definite trajectories — rays. In the quantum case, however, when the wavelength cannot be regarded as small, the concept of the coordinates of a photon becomes vacuous. We shall see later (see § 4) that in the mathematical formalism of the theory the non-measurability of the photon’s coordinates already shows itself in the impossibility of forming from its wave function a quantity that could play the part of a probability density satisfying the requirements which relativistic invariance demands.

On the basis of all that has been said, it is natural to suppose that the future theory will renounce altogether the consideration of the time development of the processes of interaction of particles. It will show that in these processes no exactly definable characteristics exist (even within the limits of the usual quantum-mechanical accuracy), so that the description of a process in time will turn out to be just as illusory as classical trajectories proved to be in non-relativistic quantum mechanics. The only observable quantities will be the characteristics (momenta, polarizations) of free particles — the initial particles that enter into the interaction and the final particles that have arisen as a result of the process (*L. D. Landau, R. Peierls, 1930*).

The characteristic formulation of a problem in relativistic quantum theory consists in the determination of the probability amplitudes of transitions connecting given initial and final states (that is, at $t \rightarrow \mp\infty$) of a system of particles. The totality of the transition amplitudes between all possible states constitutes the *scattering matrix*, or *S-matrix*. This matrix will be the carrier of all the information about the processes of interaction of particles that possesses an observable physical meaning (*W. Heisenberg, 1938*).

At the present time a complete, logically closed relativistic quantum theory does not yet exist. We shall see that the existing theory introduces new physical aspects into the manner of describing the states of particles, which acquires certain features of field theory (see § 10). The theory is constructed, however, to a considerable degree on the model of, and with the help of the concepts of, ordinary quantum mechanics. This way of building the theory has led to success in the domain of quantum electrodynamics. The absence of complete logical closure shows itself in this theory in the existence of divergent expressions when its mathematical apparatus is applied directly; but for the removal of these divergences there exist entirely unambiguous procedures. Nevertheless these procedures retain to a considerable degree the character of semi-empirical recipes, and our confidence in the correctness of the results obtained in this way rests, in the end, on their excellent agreement with experiment, and not on the internal consistency and logical coherence of the fundamental principles of the theory.

the electron; we thus set aside coordinate measurements by collision, in which over the time of observation the result occurs with a probability short of unity. In such a case a deflection of the particle does allow the electron’s position to be inferred, but from the absence of a deflection nothing at all can be concluded.

Photon

§2. Quantization of the Free Electromagnetic Field

In order to treat the electromagnetic field as a quantum object, it is convenient to start from a classical description in which the field is characterized by a discrete, albeit infinite, set of variables; such a description will allow one to apply the standard formalism of quantum mechanics directly. A description by means of potentials specified at every point of space is, in essence, a description in terms of a continuous set of variables.

Let $\mathbf{A}(\mathbf{r}, t)$ be the vector potential of the free electromagnetic field, satisfying the “transversality condition”

$$\nabla \cdot \mathbf{A} = 0. \quad (2.1)$$

The scalar potential then vanishes, $\Phi = 0$, and the fields $\mathbf{E}$ and $\mathbf{H}$ are

$$\mathbf{E} = -\dot{\mathbf{A}}, \quad \mathbf{H} = \nabla \times \mathbf{A}. \quad (2.2)$$

Maxwell’s equations reduce to the wave equation for $\mathbf{A}$:

$$\Delta \mathbf{A} - \frac{\partial^2 \mathbf{A}}{\partial t^2} = 0. \quad (2.3)$$

As is well known (see Vol. II, § 52), in classical electrodynamics the passage to a description in terms of a discrete set of variables is effected by considering the field inside a large but finite volume V .¹ Let us recall, omitting the details of the calculation, how this is done.

A field in a finite volume can be expanded in travelling plane waves, so that its potential takes the form of a series

$$\mathbf{A} = \sum_{\mathbf{k}} (\mathbf{a}_{\mathbf{k}} e^{i\mathbf{k}\mathbf{r}} + \mathbf{a}_{\mathbf{k}}^* e^{-i\mathbf{k}\mathbf{r}}), \quad (2.4)$$

where the coefficients $\mathbf{a}_{\mathbf{k}}$ depend on time according to

$$\mathbf{a}_{\mathbf{k}} \sim e^{-i\omega t}, \quad \omega = |\mathbf{k}|. \quad (2.5)$$

By virtue of condition (2.1) the complex vectors $\mathbf{a}_{\mathbf{k}}$ are orthogonal to the corresponding wave vectors: $\mathbf{a}_{\mathbf{k}} \mathbf{k} = 0$.

The summation in (2.4) runs over an infinite discrete set of values of the wave vector (its three components k_x, k_y, k_z). The transition to an integration over a continuous distribution is made with the aid of the expression

$$\frac{d^3 k}{(2\pi)^3}$$

¹To avoid burdening formulae with superfluous factors, we set $V = 1$.

for the number of allowed values of $\mathbf{k}$ falling within a volume element $d^3k = dk_x dk_y dk_z$ of $\mathbf{k}$ -space.

Once the vectors $\mathbf{a}_\mathbf{k}$ are given, the field inside the volume in question is completely determined. One may therefore view these quantities as a discrete set of classical “field variables”. In order to find the passage to a quantum theory, however, a further transformation of these quantities is needed, one that brings the field equations into a form analogous to the canonical (Hamiltonian) equations of classical mechanics. The canonical variables of the field are introduced by means of

$$\mathbf{Q}_\mathbf{k} = \frac{1}{\sqrt{4\pi}}(\mathbf{a}_\mathbf{k} + \mathbf{a}_\mathbf{k}^*), \quad \mathbf{P}_\mathbf{k} = -\frac{i\omega}{\sqrt{4\pi}}(\mathbf{a}_\mathbf{k} - \mathbf{a}_\mathbf{k}^*) = \dot{\mathbf{Q}}_\mathbf{k} \quad (2.6)$$

(they are evidently real). The vector potential is expressed through the canonical variables as

$$\mathbf{A} = \sqrt{4\pi} \sum_{\mathbf{k}} \left(\mathbf{Q}_\mathbf{k} \cos \mathbf{k}\mathbf{r} - \frac{1}{\omega} \mathbf{P}_\mathbf{k} \sin \mathbf{k}\mathbf{r} \right). \quad (2.7)$$

To find the Hamiltonian H one must evaluate the total field energy

$$\frac{1}{8\pi} \int (\mathbf{E}^2 + \mathbf{H}^2) d^3x,$$

expressed in terms of the quantities $\mathbf{Q}_\mathbf{k}$, $\mathbf{P}_\mathbf{k}$. Substituting the expansion (2.7) for $\mathbf{A}$, computing $\mathbf{E}$ and $\mathbf{H}$ by means of (2.2), and integrating, one obtains

$$H = \frac{1}{2} \sum_{\mathbf{k}} (\mathbf{P}_\mathbf{k}^2 + \omega^2 \mathbf{Q}_\mathbf{k}^2).$$

Each of the vectors $\mathbf{P}_\mathbf{k}$ and $\mathbf{Q}_\mathbf{k}$ is perpendicular to the wave vector $\mathbf{k}$ and therefore has two independent components. The direction of these vectors fixes the polarization direction of the corresponding wave. Denoting the two components of $\mathbf{Q}_\mathbf{k}$, $\mathbf{P}_\mathbf{k}$ (in the plane perpendicular to $\mathbf{k}$) by $Q_{\mathbf{k}\alpha}$, $P_{\mathbf{k}\alpha}$ ($\alpha = 1, 2$), we rewrite the Hamiltonian as

$$H = \sum_{\mathbf{k}\alpha} \frac{1}{2} (P_{\mathbf{k}\alpha}^2 + \omega^2 Q_{\mathbf{k}\alpha}^2). \quad (2.8)$$

The Hamiltonian has thus broken up into a sum of independent terms, each involving just one pair $Q_{\mathbf{k}\alpha}$, $P_{\mathbf{k}\alpha}$. Every term answers to a travelling wave of a given wave vector and polarization and reproduces the structure of a single harmonic-oscillator Hamiltonian. One accordingly speaks of this decomposition as the expansion of the field in *oscillators*.

We now turn to the quantization of the free electromagnetic field. The method of classical description just set out makes the route to quantum theory obvious. The canonical variables — the generalized coordinates $Q_{\mathbf{k}\alpha}$ and generalized momenta $P_{\mathbf{k}\alpha}$ — are now to be treated as operators obeying the commutation rule

$$\hat{P}_{\mathbf{k}\alpha} \hat{Q}_{\mathbf{k}\alpha} - \hat{Q}_{\mathbf{k}\alpha} \hat{P}_{\mathbf{k}\alpha} = -i \quad (2.9)$$

(operators with different $\mathbf{k}\alpha$ all commute with one another). Together with them the potential $\mathbf{A}$ and, by (2.2), the field strengths $\mathbf{E}$ and $\mathbf{H}$ also become (Hermitian) operators.

A consistent determination of the field Hamiltonian requires evaluation of the integral

$$\hat{H} = \frac{1}{8\pi} \int (\hat{\mathbf{E}}^2 + \hat{\mathbf{H}}^2) d^3x, \quad (2.10)$$

in which $\hat{\mathbf{E}}$ and $\hat{\mathbf{H}}$ are expressed through the operators $\hat{P}_{\mathbf{k}\alpha}$, $\hat{Q}_{\mathbf{k}\alpha}$. In practice, however, the non-commutativity of the latter does not make itself felt here, since the products $Q_{\mathbf{k}\alpha}P_{\mathbf{k}\alpha}$ enter with the factor $\cos \mathbf{k}\mathbf{r} \cdot \sin \mathbf{k}\mathbf{r}$, which vanishes upon integration over the full volume. Consequently the Hamiltonian takes the form

$$\hat{H} = \sum_{\mathbf{k}\alpha} \frac{1}{2} (\hat{P}_{\mathbf{k}\alpha}^2 + \omega^2 \hat{Q}_{\mathbf{k}\alpha}^2), \quad (2.11)$$

exactly reproducing the classical Hamiltonian, as one would naturally expect.

Finding the eigenvalues of this Hamiltonian requires no special computation, since the problem reduces to the familiar one of the energy levels of linear oscillators (see Vol. III, § 23). We can therefore write down at once the energy levels of the field:

$$E = \sum_{\mathbf{k}\alpha} \left(N_{\mathbf{k}\alpha} + \frac{1}{2} \right) \omega, \quad (2.12)$$

where $N_{\mathbf{k}\alpha}$ are non-negative integers.

Discussion of this formula will be taken up in the next section; here we write out the matrix elements of the quantities $Q_{\mathbf{k}\alpha}$, which can be obtained directly from the known formulae for the matrix elements of the oscillator coordinate (see Vol. III, § 23). The non-vanishing matrix elements are

$$\langle N_{\mathbf{k}\alpha} | Q_{\mathbf{k}\alpha} | N_{\mathbf{k}\alpha} - 1 \rangle = \sqrt{\frac{N_{\mathbf{k}\alpha}}{2\omega}}. \quad (2.13)$$

The matrix elements of $P_{\mathbf{k}\alpha} = \dot{Q}_{\mathbf{k}\alpha}$ differ from those of $Q_{\mathbf{k}\alpha}$ only by a factor $\pm i\omega$.

For subsequent calculations it will, however, be more convenient to work not with $Q_{\mathbf{k}\alpha}$, $P_{\mathbf{k}\alpha}$ themselves but with their linear combinations $\omega Q_{\mathbf{k}\alpha} \pm iP_{\mathbf{k}\alpha}$, which have matrix elements only for the transitions $N_{\mathbf{k}\alpha} \rightarrow N_{\mathbf{k}\alpha} \pm 1$. Accordingly we introduce the operators

$$\hat{c}_{\mathbf{k}\alpha} = \frac{1}{\sqrt{2\omega}} (\omega \hat{Q}_{\mathbf{k}\alpha} + i \hat{P}_{\mathbf{k}\alpha}), \quad \hat{c}_{\mathbf{k}\alpha}^+ = \frac{1}{\sqrt{2\omega}} (\omega \hat{Q}_{\mathbf{k}\alpha} - i \hat{P}_{\mathbf{k}\alpha}) \quad (2.14)$$

(the classical quantities $c_{\mathbf{k}\alpha}$, $c_{\mathbf{k}\alpha}^*$ coincide, up to a factor $\sqrt{2\pi/\omega}$, with the coefficients $a_{\mathbf{k}\alpha}$, $a_{\mathbf{k}\alpha}^*$ in the expansion (2.4)).

The matrix elements of these operators are

$$\langle N_{\mathbf{k}\alpha} - 1 | \hat{c}_{\mathbf{k}\alpha} | N_{\mathbf{k}\alpha} \rangle = \langle N_{\mathbf{k}\alpha} | \hat{c}_{\mathbf{k}\alpha}^+ | N_{\mathbf{k}\alpha} - 1 \rangle = \sqrt{N_{\mathbf{k}\alpha}}. \quad (2.15)$$

The commutation rule between $\hat{c}_{\mathbf{k}\alpha}$ and $\hat{c}_{\mathbf{k}\alpha}^+$ follows from the definition (2.14) and the rule (2.9):

$$\hat{c}_{\mathbf{k}\alpha} \hat{c}_{\mathbf{k}\alpha}^+ - \hat{c}_{\mathbf{k}\alpha}^+ \hat{c}_{\mathbf{k}\alpha} = 1. \quad (2.16)$$

Returning to the vector potential, we adopt an expansion of the same type as (2.4), but with the coefficients now promoted to operators:

$$\hat{\mathbf{A}} = \sum_{\mathbf{k}\alpha} (\hat{c}_{\mathbf{k}\alpha} \mathbf{A}_{\mathbf{k}\alpha} + \hat{c}_{\mathbf{k}\alpha}^+ \mathbf{A}_{\mathbf{k}\alpha}^*), \quad (2.17)$$

where

$$\mathbf{A}_{\mathbf{k}\alpha} = \sqrt{4\pi} \frac{\mathbf{e}^{(\alpha)}}{\sqrt{2\omega}} e^{i\mathbf{k}\mathbf{r}}. \quad (2.18)$$

Here $\mathbf{e}^{(\alpha)}$ denotes a unit vector specifying the polarization direction of the oscillator; the vectors $\mathbf{e}^{(\alpha)}$ are perpendicular to the wave vector $\mathbf{k}$, and for each $\mathbf{k}$ there are two independent polarizations.

Analogously, for the operators $\hat{\mathbf{E}}$ and $\hat{\mathbf{H}}$ we write

$$\hat{\mathbf{E}} = \sum_{\mathbf{k}\alpha} (\hat{c}_{\mathbf{k}\alpha} \mathbf{E}_{\mathbf{k}\alpha} + \hat{c}_{\mathbf{k}\alpha}^+ \mathbf{E}_{\mathbf{k}\alpha}^*), \quad \hat{\mathbf{H}} = \sum_{\mathbf{k}\alpha} (\hat{c}_{\mathbf{k}\alpha} \mathbf{H}_{\mathbf{k}\alpha} + \hat{c}_{\mathbf{k}\alpha}^+ \mathbf{H}_{\mathbf{k}\alpha}^*), \quad (2.19)$$

with

$$\mathbf{E}_{\mathbf{k}\alpha} = i\omega \mathbf{A}_{\mathbf{k}\alpha}, \quad \mathbf{H}_{\mathbf{k}\alpha} = \mathbf{n} \times \mathbf{E}_{\mathbf{k}\alpha} \quad (\mathbf{n} = \mathbf{k}/\omega). \quad (2.20)$$

The vectors $\mathbf{A}_{\mathbf{k}\alpha}$ are mutually orthogonal in the sense that

$$\int \mathbf{A}_{\mathbf{k}\alpha} \mathbf{A}_{\mathbf{k}'\alpha'}^* d^3x = \frac{2\pi}{\omega} \delta_{\alpha\alpha'} \delta_{\mathbf{k}\mathbf{k}'}. \quad (2.21)$$

To see this, note that when $\mathbf{A}_{\mathbf{k}\alpha}$ and $\mathbf{A}_{\mathbf{k}'\alpha'}^*$ carry different wave vectors their product involves $e^{i(\mathbf{k}-\mathbf{k}')\mathbf{r}}$, which integrates to zero over the volume; when only the polarizations differ, the orthogonality $\mathbf{e}^{(\alpha)} \mathbf{e}^{(\alpha')*} = 0$ of the two independent polarization directions provides the required vanishing. Analogous relations hold for the vectors $\mathbf{E}_{\mathbf{k}\alpha}$, $\mathbf{H}_{\mathbf{k}\alpha}$. Their normalization is conveniently written as

$$\frac{1}{4\pi} \int (\mathbf{E}_{\mathbf{k}\alpha} \mathbf{E}_{\mathbf{k}'\alpha'}^* + \mathbf{H}_{\mathbf{k}\alpha} \mathbf{H}_{\mathbf{k}'\alpha'}^*) d^3x = \omega \delta_{\mathbf{k}\mathbf{k}'} \delta_{\alpha\alpha'}. \quad (2.22)$$

Substituting the operators (2.19) into (2.10) and integrating with the help of (2.22), we get the field Hamiltonian expressed through the operators $\hat{c}$, $\hat{c}^+$:

$$\hat{H} = \sum_{\mathbf{k}\alpha} \frac{1}{2} \omega (\hat{c}_{\mathbf{k}\alpha} \hat{c}_{\mathbf{k}\alpha}^+ + \hat{c}_{\mathbf{k}\alpha}^+ \hat{c}_{\mathbf{k}\alpha}). \quad (2.23)$$

In the representation under discussion (with the matrix elements of $\hat{c}$, $\hat{c}^+$ given by (2.15)) this operator is diagonal, and its eigenvalues coincide, naturally, with (2.12).

In classical theory the field momentum is defined by the integral

$$\mathbf{P} = \frac{1}{4\pi} \int \mathbf{E} \times \mathbf{H} d^3x.$$

On passing to the quantum theory by replacing $\mathbf{E}$ and $\mathbf{H}$ with the operators (2.19), one readily finds

$$\hat{\mathbf{P}} = \sum_{\mathbf{k}\alpha} \frac{1}{2} (\hat{P}_{\mathbf{k}\alpha}^2 + \omega^2 \hat{Q}_{\mathbf{k}\alpha}^2) \mathbf{n} \quad (2.24)$$

— which mirrors the familiar classical link between energy and momentum of plane waves. This operator has eigenvalues

$$\mathbf{P} = \sum_{\mathbf{k}\alpha} \mathbf{k} \left(N_{\mathbf{k}\alpha} + \frac{1}{2} \right). \quad (2.25)$$

The operator representation realized by the matrix elements (2.15) is called the “occupation-number representation”: it amounts to characterizing the state of the system (the field) through the quantum numbers $N_{\mathbf{k}\alpha}$ (*occupation numbers*). Within this representation the field operators (2.19) (and hence the Hamiltonian (2.11)) act on the system’s wave function, written as a function of the numbers $N_{\mathbf{k}\alpha}$; we denote it $\Phi(N_{\mathbf{k}\alpha}, t)$. The field operators (2.19) do not depend on time explicitly. This corresponds to the Schrödinger representation of operators, familiar from non-relativistic quantum mechanics. What does depend on time is the state of the system, $\Phi(N_{\mathbf{k}\alpha}, t)$, its time dependence being governed by the Schrödinger equation

$$i \frac{\partial \Phi}{\partial t} = \hat{H} \Phi.$$

This description of the field is relativistically invariant in essence, since it rests on the invariant Maxwell equations. Yet the invariance is not exhibited explicitly — above all because the spatial coordinates and the time enter the description in a highly asymmetric way.

Within a relativistic framework it is desirable to give the description a more overtly invariant appearance. For this purpose one uses the so-called Heisenberg representation, wherein the explicit dependence on time is shifted onto the operators (see Vol. III, § 13). Time and spatial coordinates then appear symmetrically in the field-operator expressions, while the system state Φ depends only on the occupation numbers.

For the operator $\hat{\mathbf{A}}$ the passage to the Heisenberg representation amounts to replacing, in each term of the sum in (2.17), (2.18), the factor $e^{i\mathbf{k}\mathbf{r}}$ by $e^{i(\mathbf{k}\mathbf{r} - \omega t)}$; that is, one is to understand by $\mathbf{A}_{\mathbf{k}\alpha}$ the time-dependent functions

$$\mathbf{A}_{\mathbf{k}\alpha} = \sqrt{4\pi} \frac{\mathbf{e}^{(\alpha)}}{\sqrt{2\omega}} e^{-i(\omega t - \mathbf{k}\mathbf{r})}. \quad (2.26)$$

This is easily verified by noting that the matrix element of a Heisenberg operator for the transition $i \rightarrow f$ must contain the factor $\exp\{-i(E_i - E_f)t\}$, where E_i and E_f are the energies of the initial and final states (see Vol. III, § 13). For a transition in which $N_{\mathbf{k}}$ decreases or increases by one, this factor reduces to $e^{-i\omega t}$ or $e^{i\omega t}$, respectively. The indicated replacement satisfies this requirement.

Henceforth (in treating both the electromagnetic field and the fields of particles) we shall always understand the Heisenberg representation of operators.

§3. Photons

Let us now turn to a discussion of the quantization formulae derived above.

To begin with, formula (2.12) for the energy of the field exposes the following difficulty. At the lowest energy level, all oscillator quantum numbers $N_{\mathbf{k}\alpha}$ vanish (this

state is known as the *electromagnetic field vacuum*). Yet even in this state every oscillator retains a nonzero “zero-point energy” $\omega/2$. Summing over the entire infinite set of oscillators then produces an infinite result. One is thus confronted with one of the “divergences” stemming from the absence of full logical self-consistency in the existing theory.

As long as one is concerned only with the eigenvalues of the field energy, this difficulty can be eliminated by simply subtracting away the zero-point oscillation energy, i.e. by writing for the energy and momentum of the field²

$$E = \sum_{\mathbf{k}\alpha} N_{\mathbf{k}\alpha} \omega, \quad \mathbf{P} = \sum_{\mathbf{k}\alpha} N_{\mathbf{k}\alpha} \mathbf{k}. \quad (3.1)$$

These formulae make it possible to introduce the concept fundamental to all of quantum electrodynamics: that of *light quanta*, or *photons*³. Namely, the free electromagnetic field can be regarded as a collection of particles, each carrying energy $\omega (= \hbar\omega)$ and momentum $\mathbf{k} (= \hbar\omega/c)$. The relationship between a photon’s energy and momentum is precisely what relativistic mechanics requires for particles of zero rest mass that propagate at the speed of light. The occupation numbers $N_{\mathbf{k}\alpha}$ acquire the meaning of the number of photons with given momenta $\mathbf{k}$ and polarizations $\mathbf{e}^{(\alpha)}$. The polarization property of a photon is analogous to the notion of spin for other particles (the specific features of the photon in this regard will be examined below, in § 6).

It is easy to see that the mathematical formalism developed in the preceding section is fully consistent with the picture of the electromagnetic field as an assembly of photons; it is nothing other than the apparatus of so-called second quantization as applied to a system of photons⁴. In this method (see Vol. III, § 64) the occupation numbers serve as the independent variables, and the operators act on functions of those numbers. The central role is played by the “annihilation” and “creation” operators for particles, which respectively decrease or increase an occupation number by one. Exactly such operators are $\hat{c}_{\mathbf{k}\alpha}$ and $\hat{c}_{\mathbf{k}\alpha}^+$: the operator $\hat{c}_{\mathbf{k}\alpha}$ destroys a photon in the state $\mathbf{k}\alpha$, while $\hat{c}_{\mathbf{k}\alpha}^+$ creates one in that state.

The commutation rule (2.16) corresponds to particles obeying Bose statistics. Photons are therefore bosons, as one might have anticipated: the number of photons occupying any given state can be arbitrary (we shall return to the significance of this fact in § 5).

The plane waves $\mathbf{A}_{\mathbf{k}\alpha}$ (2.26), which appear in the operator $\hat{\mathbf{A}}$ (2.17) as coefficients multiplying the photon annihilation operators, can be interpreted as the wave functions of photons possessing definite momenta $\mathbf{k}$ and polarizations $\mathbf{e}^{(\alpha)}$. This interpretation corresponds to expanding the ψ -operator as a series in the stationary-state wave functions of the particle within the non-relativistic second-quantization framework (unlike the

²This subtraction can be carried out in a formally consistent manner by agreeing to interpret the operator products in (2.10) as “normal-ordered”, i.e. arranged so that the operators $\hat{c}^+$ always stand to the left of the operators $\hat{c}$. Formula (2.23) then takes the form

$$\hat{H} = \sum_{\mathbf{k}\alpha} (\omega \hat{c}_{\mathbf{k}\alpha}^+ \hat{c}_{\mathbf{k}\alpha}).$$

³The photon concept was first put forward by *Einstein* (A. Einstein, 1905).

⁴The method of second quantization as applied to radiation theory was developed by *Dirac* (P. A. M. Dirac, 1927).

non-relativistic case, however, the expansion (2.17) involves both annihilation and creation operators; the significance of this distinction will become clear later, see § 12).

The wave function (2.26) is normalized by the condition

$$\int \frac{1}{4\pi} (|\mathbf{E}_{\mathbf{k}\alpha}|^2 + |\mathbf{H}_{\mathbf{k}\alpha}|^2) d^3x = \omega \quad (3.2)$$

This amounts to normalizing to one photon in a volume $V = 1$. Indeed, the integral on the left-hand side of the equation represents the quantum-mechanical expectation value of the photon's energy in a state described by the given wave function⁵. On the right-hand side of (3.2) stands the energy of a single photon.

The role of the “Schrödinger equation” for the photon is played by Maxwell's equations. In the present case (for the potential $\mathbf{A}(\mathbf{r}, t)$ satisfying condition (2.1)) this is the wave equation

$$\frac{\partial^2 \mathbf{A}}{\partial t^2} - \Delta \mathbf{A} = 0.$$

In the general case of arbitrary stationary states, the photon “wave functions” are complex solutions of this equation whose time dependence enters through the factor $e^{-i\omega t}$.

Speaking of the photon wave function, let us emphasize once more that it cannot by any means be treated as a probability amplitude for the spatial localization of a photon — in contrast to the basic meaning of the wave function in non-relativistic quantum mechanics. This is linked to the fact that (as noted in § 1) the notion of photon coordinates has no physical meaning at all. We shall revisit the mathematical side of this issue at the end of the next section.

The Fourier-transform components of $\mathbf{A}(\mathbf{r}, t)$ with respect to the coordinates constitute the photon wave function in the momentum representation; we denote it $\mathbf{A}(\mathbf{k}, t) = \mathbf{A}(\mathbf{k})e^{-i\omega t}$. Thus, for a state with a definite momentum $\mathbf{k}$ and polarization $\mathbf{e}^{(\alpha)}$ the momentum-representation wave function is given simply by the coefficient of the exponential factor in (2.26):

$$\mathbf{A}_{\mathbf{k}\alpha}(\mathbf{k}', \alpha') = \sqrt{4\pi} \frac{\mathbf{e}^{(\alpha)}}{\sqrt{2\omega}} \delta_{\mathbf{k}'\mathbf{k}} \delta_{\alpha\alpha'}. \quad (3.3)$$

In keeping with the measurability of the momentum of a free particle, the momentum-representation wave function carries deeper physical content than its coordinate-space counterpart: it provides a way to compute the probabilities $w_{\mathbf{k}\alpha}$ of the various values of momentum and polarization for a photon in a given state. According to the general rules of quantum mechanics, $w_{\mathbf{k}\alpha}$ is given by the squared modulus of the expansion coefficients of $\mathbf{A}(\mathbf{k}')$ in the wave functions of states with definite $\mathbf{k}$ and $\mathbf{e}^{(\alpha)}$:

$$w_{\mathbf{k}\alpha} \propto \left| \sum_{\mathbf{k}'\alpha'} \mathbf{A}_{\mathbf{k}\alpha}^*(\mathbf{k}', \alpha') \mathbf{A}(\mathbf{k}') \right|^2$$

⁵Note that the coefficient $1/(4\pi)$ in the integral (3.2) is twice the usual coefficient $1/(8\pi)$ in (2.10). This difference is ultimately connected with the complex character of the vectors $\mathbf{E}_{\mathbf{k}\alpha}$, $\mathbf{H}_{\mathbf{k}\alpha}$, as opposed to the Hermitian field operators $\hat{\mathbf{E}}$, $\hat{\mathbf{H}}$.

(the proportionality coefficient depends on the choice of normalization for the functions). Substituting (3.3), we obtain

$$w_{\mathbf{k}\alpha} \propto |\mathbf{e}^{(\alpha)} \mathbf{A}(\mathbf{k})|^2. \quad (3.4)$$

Summing over the two polarizations, we find the probability that a photon has momentum $\mathbf{k}$:

$$w_{\mathbf{k}} \propto |\mathbf{A}(\mathbf{k})|^2. \quad (3.5)$$

§4. Gauge invariance

As is well known, the choice of field potentials in classical electrodynamics is not unique: the components of the 4-potential A_μ may be subjected to an arbitrary *gauge* (or *gradient*) transformation of the form

$$A_\mu \rightarrow A_\mu + \partial_\mu \chi, \quad (4.1)$$

where χ is an arbitrary function of the coordinates and time (see Vol. II, § 18). In the case of a plane wave, restricting oneself to transformations that preserve the functional form of the potential (its proportionality to $\exp(-ik_\mu x^\mu)$), the non-uniqueness amounts to the freedom of adding to the wave amplitude an arbitrary 4-vector proportional to k^μ .

The ambiguity of the potential persists, of course, in the quantum theory as well — as applied to the field operators or to the photon wave functions. Without prejudging the choice of potentials, one should write in place of (2.17) an analogous expansion for the operator 4-potential

$$\hat{A}^\mu = \sum_{\mathbf{k}\alpha} (\hat{c}_{\mathbf{k}\alpha} A_{\mathbf{k}\alpha}^\mu + \hat{c}_{\mathbf{k}\alpha}^+ A_{\mathbf{k}\alpha}^{\mu*}), \quad (4.2)$$

where the wave functions $A_{\mathbf{k}\alpha}^\mu$ are 4-vectors of the form

$$A_{\mathbf{k}}^\mu = \sqrt{4\pi} \frac{e^\mu}{\sqrt{2\omega}} e^{-ik_\nu x^\nu}, \quad e_\mu e^{\mu*} = -1,$$

or, in a compact notation that suppresses the four-dimensional vector indices:

$$A_{\mathbf{k}} = \sqrt{4\pi} \frac{e}{\sqrt{2\omega}} e^{-ikx}, \quad ee^* = -1. \quad (4.3)$$

Here the 4-momentum is $k_\mu = (\omega, \mathbf{k})$ (so that $kx = \omega t - \mathbf{k}\mathbf{r}$), and e is the unit polarization 4-vector⁶.

If one restricts oneself to gauge transformations that do not alter how the function (4.3) depends on the coordinates and time, they amount to the replacement

$$e_\mu \rightarrow e_\mu + \chi k_\mu, \quad (4.4)$$

⁶Expression (4.3) does not have an entirely relativistically covariant (4-vector) form, which is related to the non-invariant character of our normalization to a finite volume $V = 1$. This is, however, of no fundamental significance and is fully compensated by the convenience of such a normalization scheme. We shall see later that it ensures simple and automatic derivation of real physical quantities in the proper invariant form.

where $\chi = \chi(k^\mu)$ is an arbitrary function. The transversality of the polarization means that one can always choose a gauge in which the 4-vector e takes the form

$$e^\mu = (0, \mathbf{e}), \quad \mathbf{ek} = 0 \quad (4.5)$$

(we shall call such a gauge *three-dimensionally transverse*). In an invariant four-dimensional notation this requirement is written as the condition of *four-dimensional transversality*

$$ek = 0. \quad (4.6)$$

Note that this condition (as well as the normalization condition $ee^* = -1$) is not violated by the transformation (4.4), by virtue of $k^2 = 0$. On the other hand, the vanishing of the 4-momentum squared of a particle signifies that its mass is zero. A direct link between gauge invariance and the vanishing of the photon mass is thereby revealed (further aspects of this connection will be discussed in § 14).

No measurable physical quantity should change under a gauge transformation of the wave functions of the photons taking part in a given process. This requirement of *gauge invariance* plays an even greater role in quantum electrodynamics than in the classical theory. As we shall see from numerous examples, it serves here, alongside the requirement of relativistic invariance, as a powerful heuristic principle.

In turn, the gauge invariance of the theory is closely bound up with the law of conservation of electric charge; we shall dwell on this aspect in § 43.

We already remarked in the preceding section that the coordinate-space wave function of a photon cannot be interpreted as a probability amplitude for its spatial localization. From a mathematical standpoint this circumstance manifests itself in the impossibility of constructing, out of the wave function, a quantity whose formal properties alone would qualify it to serve as a probability density. Such a quantity would have to be an essentially positive bilinear combination of the wave function A_μ and its complex conjugate. Moreover, it would need to satisfy definite requirements of relativistic covariance — namely, to be the time component of a 4-vector (the point being that the continuity equation expressing conservation of particle number takes the four-dimensional form of a vanishing 4-divergence of the current 4-vector; the time component of this current is, in the present context, the probability density for the localization of the particle; see Vol. II, § 29). On the other hand, gauge invariance demands that A_μ enter the current only through the antisymmetric tensor $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu = -i(k_\mu A_\nu - k_\nu A_\mu)$. The current 4-vector would therefore have to be assembled bilinearly from $F_{\mu\nu}$ and $F_{\mu\nu}^*$ (together with components of the 4-vector k_μ). But no such 4-vector can be constructed at all: every expression satisfying the stated conditions (for instance, $k^\lambda F_{\mu\nu}^* F_{\lambda}{}^\nu$) vanishes by the transversality condition ($k^\lambda F_{\nu\lambda} = 0$), quite apart from the fact that it would not be essentially positive (since it contains odd powers of the components k_μ).

§ 5. The electromagnetic field in quantum theory

Treating the field as a collection of photons is the sole description that fully captures the physical meaning of the electromagnetic field within quantum theory. It replaces the classical characterization through field strengths. The field strengths themselves enter the mathematical apparatus of the photon picture as operators of second quantization.

It is well known that a quantum system behaves classically when the quantum numbers characterizing its stationary states become large. For the free electromagnetic field (within a given volume) this requires that the oscillator quantum numbers — that is, the photon occupation numbers $N_{\mathbf{k}\alpha}$ — be large. The fact that photons obey Bose statistics is of profound importance in this context. In the mathematical formalism, the link between Bose statistics and classical field behavior is encoded in the commutation rules for $\hat{c}_{\mathbf{k}\alpha}$ and $\hat{c}_{\mathbf{k}\alpha}^\dagger$. When $N_{\mathbf{k}\alpha}$ is large, so that the matrix elements of these operators are also large, the unity appearing on the right-hand side of the commutation relation (2.16) may be dropped, giving

$$\hat{c}_{\mathbf{k}\alpha}\hat{c}_{\mathbf{k}\alpha}^\dagger \simeq \hat{c}_{\mathbf{k}\alpha}^\dagger\hat{c}_{\mathbf{k}\alpha},$$

i.e. the operators go over into mutually commuting classical quantities $c_{\mathbf{k}\alpha}$, $c_{\mathbf{k}\alpha}^*$ that determine the classical field strengths.

The quasi-classicality condition for the field requires, however, a further refinement. The point is that if all the numbers $N_{\mathbf{k}\alpha}$ are large, then upon summing over every state $\mathbf{k}\alpha$ the field energy is in any case infinite, rendering the condition meaningless.

A physically sensible formulation of the quasi-classicality criterion rests on examining values of the field averaged over some small time interval Δt . If one writes the classical electric field $\bar{\mathbf{E}}$ (or the magnetic field $\bar{\mathbf{H}}$) as a Fourier integral over time, then upon time-averaging over an interval Δt only those Fourier components whose frequencies satisfy $\omega\Delta t \lesssim 1$ contribute appreciably to the mean value of $\bar{\mathbf{E}}$; in the opposite limit the oscillating factor $e^{-i\omega t}$ nearly averages to zero. Consequently, in establishing the quasi-classicality condition for the time-averaged field, one need consider only those quantum oscillators whose frequencies satisfy $\omega \lesssim 1/\Delta t$. It suffices to require that the quantum numbers of these oscillators be large. The number of oscillators with frequencies between zero and $\omega \sim 1/\Delta t$ (referred to the volume $V = 1$) is of order⁷

$$\left(\frac{\omega}{c}\right)^3 \sim \frac{1}{(c\Delta t)^3}. \quad (5.1)$$

The total energy of the field per unit volume is of order $\bar{\mathbf{E}}^2$. Dividing this quantity by the number of oscillators and by a characteristic energy of one photon ($\sim \hbar\omega$), one obtains an estimate for the photon occupation numbers

$$N_{\mathbf{k}} \sim \frac{\bar{\mathbf{E}}^2 c^3}{\hbar\omega^4}.$$

Demanding that this number be large, we arrive at the inequality

$$|\mathbf{E}| \gg \frac{\sqrt{\hbar c}}{(c\Delta t)^2}. \quad (5.2)$$

This is the sought-after condition permitting a classical treatment of the time-averaged (over intervals Δt) field. We see that the field must be sufficiently strong — all the stronger, the shorter the averaging interval Δt . For time-varying fields this interval must not, of course, exceed the characteristic periods over which the field changes appreciably.

⁷In this section we use ordinary units.

Consequently, sufficiently weak time-varying fields can under no circumstances be quasi-classical. Only for static (time-independent) fields may one set $\Delta t \rightarrow \infty$, so that the right-hand side of the inequality (5.2) vanishes. This means that a static field is always classical.

It has already been noted that the classical expressions for the electromagnetic field in the form of a superposition of plane waves must be treated as operator-valued in the quantum theory. The physical meaning of these operators is, however, rather limited. Indeed, a physically meaningful field operator ought to yield vanishing field values in the photon-vacuum state. Yet the expectation value of the squared-field operator $\hat{\mathbf{E}}^2$ in the ground state, which coincides up to a factor with the zero-point energy of the field, turns out to be infinite (by “expectation value” we mean the quantum-mechanical average, i.e. the corresponding diagonal matrix element of the operator). This divergence cannot be circumvented even by any formal subtraction procedure (as can be done for the field energy), since in the present case we would need to carry out the subtraction through some sensible modification of the operators $\hat{\mathbf{E}}$, $\hat{\mathbf{H}}$ themselves (rather than their squares), which proves to be impossible.

§ 6. Angular momentum and parity of the photon

As with any particle, the photon can possess a definite angular momentum. In order to elucidate the properties of this quantity for the photon, let us begin by recalling how the angular momentum of a particle is connected with its wave function in the formalism of quantum mechanics.

The total angular momentum $\mathbf{j}$ of a particle consists of the orbital part $\mathbf{l}$ and the intrinsic part — the spin $\mathbf{s}$. A particle of spin s is described by a symmetric spinor of rank $2s$, i.e. a set of $2s + 1$ components that transform into one another under coordinate-system rotations according to a specific law. The orbital angular momentum is instead linked to how the wave functions depend on coordinates: states with orbital angular momentum l have wave-function components that are expressed (linearly) in terms of spherical harmonics of order l .

The ability to distinguish consistently between spin and orbital angular momentum therefore demands that the “spin” and “coordinate” properties of the wave functions be independent: the coordinate dependence of each spinor component (at a given instant of time) should not be constrained by any supplementary conditions. Passing to the momentum representation, coordinate dependence becomes dependence on $\mathbf{k}$. In the three-dimensionally transverse gauge the photon wave function is the vector $\mathbf{A}(\mathbf{k})$. Because a vector is equivalent to a rank-2 spinor, one could formally ascribe spin 1 to the photon. This vector wave function, however, must satisfy the transversality requirement $\mathbf{kA}(\mathbf{k}) = 0$, an extra constraint on the way $\mathbf{A}(\mathbf{k})$ depends on momentum. It follows that the dependence on $\mathbf{k}$ can no longer be chosen freely for every vector component at once, making it impossible to disentangle orbital angular momentum and spin.

We note also that for the photon one cannot define spin as the angular momentum of the particle at rest, since for a photon traveling at the speed of light no rest frame exists at all.

Thus, for a photon one can speak only of its total angular momentum. It is clear

from the outset that the total angular momentum can assume only integer values. This is already evident from the fact that among the quantities characterizing the photon there are no spinors of odd rank.

As for any particle, the state of a photon is further characterized by its parity, associated with the behavior of the wave function under spatial inversion (see Vol. III, § 30). In the momentum representation, a reversal of the sign of the coordinates corresponds to reversing the sign of every component of $\mathbf{k}$. The action of the inversion operator $\hat{P}$ on a scalar function $\varphi(\mathbf{k})$ amounts simply to this sign change: $\hat{P}\varphi(\mathbf{k}) = \varphi(-\mathbf{k})$. When acting on the vector function $\mathbf{A}(\mathbf{k})$, one must additionally take into account that reversing the axes also reverses the sign of every component of the vector; hence⁸

$$\hat{P}\mathbf{A}(\mathbf{k}) = -\mathbf{A}(-\mathbf{k}). \quad (6.1)$$

Although the decomposition of the photon's angular momentum into orbital and spin parts lacks physical meaning, it is nevertheless convenient to introduce "spin" s and "orbital" angular momentum l in a formal sense as auxiliary concepts describing the transformation properties of the wave function with respect to rotations: the value $s = 1$ reflects the vectorial character of the wave function, while l is the order of the spherical harmonics entering it. Here we have in mind the wave functions of states with definite values of the photon's angular momentum, which for a free particle take the form of spherical waves. The number l determines, in particular, the parity of the photon state, equal to

$$P = (-1)^{l+1} \quad (6.2)$$

In the same formal sense one can write the angular-momentum operator $\hat{\mathbf{j}}$ as the sum $\hat{\mathbf{s}} + \hat{\mathbf{l}}$. As is well known, the angular-momentum operator is associated with infinitesimally small rotations of the coordinate system; here, with the action of such a rotation on a vector field. Within the decomposition $\hat{\mathbf{s}} + \hat{\mathbf{l}}$, the operator $\hat{\mathbf{s}}$ acts on the vector index and intermixes the various components of the vector. The operator $\hat{\mathbf{l}}$ acts on these same components viewed as functions of momentum (or of coordinates). Let us count the number of states (at a given energy) that are possible for a given value j of the photon's angular momentum (setting aside the trivial $(2j + 1)$ -fold degeneracy with respect to the direction of the angular momentum).

For independent $\mathbf{l}$ and $\mathbf{s}$ this counting is carried out simply by enumerating the ways in which, according to the rules of the vector model, the angular momenta $\hat{\mathbf{l}}$ and $\mathbf{s}$ can be composed so as to yield the required value of j . For a particle with spin $s = 1$ we would find in this way (for a given nonzero value of j) three states with the following values of l and parity:

$$l = j, \quad P = (-1)^{l+1} = (-1)^{j+1}; \quad l = j \pm 1, \quad P = (-1)^{l+1} = (-1)^j.$$

⁸We adopt the convention of defining the parity of a state by the action of the inversion operator on the polar vector $\mathbf{A}$ (or, equivalently, the corresponding electric vector $\mathbf{E} = i\omega\mathbf{A}$). This differs in sign from the action on the axial vector $\mathbf{H} = i[\mathbf{k}\mathbf{A}]$, since inversion does not reverse the direction of such a vector:

$$\hat{P}\mathbf{H}(\mathbf{k}) = \mathbf{H}(-\mathbf{k}).$$

If $j = 0$, there is only a single state (with $l = 1$) having parity $P = +1$.

This counting, however, has not accounted for the transversality condition on $\mathbf{A}$; all three components were regarded as independent. From the result obtained one must therefore subtract the number of states associated with a longitudinal vector. A longitudinal vector can be expressed as $\mathbf{k}\varphi(\mathbf{k})$, and it is apparent that, owing to its transformation behavior (with respect to rotations), its three components reduce to a single scalar φ ⁹. The extra state that is excluded by transversality would therefore belong to a particle whose wave function is a scalar (a spinor of rank zero), that is, a particle with “spin 0”¹⁰. For this state the angular momentum j equals the order of the spherical harmonics appearing in φ . Its parity, viewed as a photon state, follows from how the inversion operator acts on the vector function $\mathbf{k}\varphi$:

$$\hat{P}(\mathbf{k}\varphi) = -(-\mathbf{k})\varphi(-\mathbf{k}) = (-1)^j \mathbf{k}\varphi(\mathbf{k}),$$

i.e. it equals $(-1)^j$. Thus, from the above count of states with parity $(-1)^j$ (two for $j \neq 0$ and one for $j = 0$) one must subtract one.

We arrive finally at the result that for nonzero photon angular momentum j there exist one even-parity and one odd-parity state. For $j = 0$ we obtain no states whatsoever. This means that the photon can never have zero angular momentum, so that j takes only the values $1, 2, 3, \dots$. The impossibility of $j = 0$ is, moreover, obvious: the wave function of a state with vanishing angular momentum would have to be spherically symmetric, which is manifestly impossible for a transverse wave.

There is a standard nomenclature for the various photon states. When a photon carries angular momentum j with parity $(-1)^j$ it is called an *electric 2^j -pole* (or Ej) photon; when its parity is $(-1)^{j+1}$ it is called a *magnetic 2^j -pole* (or Mj) photon. In particular, the electric dipole photon has odd parity and $j = 1$; the electric quadrupole photon has even parity and $j = 2$; the magnetic dipole photon has even parity and $j = 1$ ¹¹.

§ 7. Spherical waves of photons

Having established the admissible values of the photon's angular momentum, our task is now to find the wave functions that correspond to them¹².

Let us first consider a formal problem: to identify the vector functions that serve as eigenfunctions of the operators $\hat{\mathbf{j}}^2$ and $\hat{j}_z$; in doing so we do not stipulate in advance which particular functions enter the photon wave functions of interest, nor do we impose the transversality constraint.

⁹Indeed, when one speaks of the transformation character of a quantity under rotation, one means a transformation at a given point, i.e. at a given $\mathbf{k}$. Under such a transformation $\mathbf{k}\varphi(\mathbf{k})$ does not change at all, i.e. it behaves as a scalar.

¹⁰Let us emphasize once more that what is meant here is not the state of any real particle. The counting carried out here is purely formal and amounts, mathematically, to sorting the complete collection of quantities that mix under rotations into irreducible representations of the rotation group.

¹¹This nomenclature mirrors the usage in the classical theory of radiation: as will be shown later (§§ 46–47), the production of electric- and magnetic-type photons is controlled by the respective electric and magnetic multipole moments of the system of charges.

¹²This problem was first studied by Heitler (W. Heitler, 1936). The form of the solution presented here is due to V. B. Berestetskii (1947).

The functions will be sought in the momentum representation, where the coordinate operator takes the form $\hat{\mathbf{r}} = i\partial/\partial\mathbf{k}$ (see III, (15.12)). For the orbital angular momentum operator one then obtains

$$\hat{\mathbf{l}} = [\hat{\mathbf{r}}\mathbf{k}] = -i \left[\mathbf{k} \frac{\partial}{\partial\mathbf{k}} \right],$$

i.e. it differs from the angular momentum operator in coordinate space only by the replacement of $\mathbf{r}$ with $\mathbf{k}$. Accordingly, the formal solution of the problem is identical in both representations.

Let us denote the desired eigenfunctions by $\mathbf{Y}_{jm}$ and call them *spherical vectors*. They must satisfy the equations

$$\hat{\mathbf{j}}^2 = j(j+1)\mathbf{Y}_{jm}, \quad \hat{j}_z \mathbf{Y}_{jm} = m\mathbf{Y}_{jm} \quad (7.1)$$

(the z axis being a fixed direction in space). We shall demonstrate that any function of the form $\mathbf{a}Y_{jm}$ possesses this property, where $\mathbf{a}$ is a vector constructed from the unit vector $\mathbf{n} = \mathbf{k}/\omega$ and Y_{jm} are the ordinary (scalar) spherical harmonics. The latter will everywhere be defined in accordance with III, § 28:

$$Y_{lm}(\mathbf{n}) = (-1)^{\frac{m+|m|}{2}} i^l \sqrt{\frac{(2l+1)(l-|m|)!}{4\pi(l+|m|)!}} P_l^{|m|}(\cos\theta) e^{im\varphi} \quad (7.2)$$

(θ, φ being the spherical angles specifying the direction of $\mathbf{n}$)¹³.

To show this, recall the commutation rule III, (29.4):

$$\{\hat{l}_i, a_k\}_- = ie_{ikl}a_l.$$

The right-hand side can be written as $(-\hat{s}_i a_k)$, where $\hat{\mathbf{s}}$ is the spin-1 operator (its action on a vector function is precisely $\hat{s}_i a_k = -ie_{ikl}a_l$; see III, § 57, problem 2). Consequently,

$$\hat{l}_i a_k - a_k \hat{l}_i = -\hat{s}_i a_k.$$

Making use of this relation, we find

$$\hat{j}_i a_k = (\hat{l}_i + \hat{s}_i) a_k = a_k \hat{l}_i.$$

It follows that

$$\hat{\mathbf{j}}^2(\mathbf{a}Y_{jm}) = \mathbf{a}\hat{\mathbf{l}}^2 Y_{jm}, \quad \hat{j}_z(\mathbf{a}Y_{jm}) = \mathbf{a}\hat{l}_z Y_{jm}.$$

Since the spherical harmonic Y_{jm} is an eigenfunction of $\hat{l}^2$ and $\hat{l}_z$ with eigenvalues $j(j+1)$ and m respectively, we arrive at equations (7.1).

¹³For future reference, note the value of these functions at $\theta = 0$ ($\mathbf{n}$ along the z axis):

$$Y_{lm}(\mathbf{n}_z) = i^l \sqrt{\frac{2l+1}{4\pi}} \delta_{m0}. \quad (7.2a)$$

Three essentially distinct types of spherical vectors are obtained by choosing the vector $\mathbf{a}$ to be one of the following¹⁴:

$$\frac{\nabla_{\mathbf{n}}}{\sqrt{j(j+1)}}, \quad \frac{[\mathbf{n}\nabla_{\mathbf{n}}]}{\sqrt{j(j+1)}}, \quad \mathbf{n}. \quad (7.3)$$

The spherical vectors are thus defined as follows:

$$\begin{aligned} \mathbf{Y}_{jm}^{(e)} &= \frac{1}{\sqrt{j(j+1)}} \nabla_{\mathbf{n}} Y_{jm}, & P &= (-1)^j; \\ \mathbf{Y}_{jm}^{(m)} &= [\mathbf{n}\mathbf{Y}_{jm}^{(e)}], & P &= (-1)^{j+1}; \\ \mathbf{Y}_{jm}^{(l)} &= \mathbf{n}Y_{jm}, & P &= (-1)^j. \end{aligned} \quad (7.4)$$

Next to each vector we have indicated its parity P . The spherical vectors of the three types are mutually orthogonal; $\mathbf{Y}_{jm}^{(l)}$ is longitudinal, while $\mathbf{Y}_{jm}^{(e)}$ and $\mathbf{Y}_{jm}^{(m)}$ are transverse with respect to $\mathbf{n}$.

The spherical vectors can be expressed through scalar spherical harmonics. In this expansion, $\mathbf{Y}_{jm}^{(m)}$ involves harmonics of a single order $l = j$ only, whereas $\mathbf{Y}_{jm}^{(e)}$ and $\mathbf{Y}_{jm}^{(l)}$ involve harmonics of orders $l = j \pm 1$. This is self-evident: one need only compare the parities listed in (7.4) with the parity $(-1)^{l+1}$ of a vector field, expressed via the order of the spherical harmonics it contains.

The spherical vectors within each type are mutually orthogonal and normalized according to

$$\int \mathbf{Y}_{jm} \mathbf{Y}_{j'm'}^* d\Omega = \delta_{jj'} \delta_{mm'}. \quad (7.5)$$

For the vectors $\mathbf{Y}_{jm}^{(l)}$ this follows immediately from the normalization of the spherical harmonics Y_{jm} . For $\mathbf{Y}_{jm}^{(e)}$ the normalization integral becomes

$$\frac{1}{j(j+1)} \int \nabla_{\mathbf{n}} Y_{jm} \nabla_{\mathbf{n}} Y_{j'm'}^* d\Omega = -\frac{1}{j(j+1)} \int Y_{j'm'}^* \Delta_{\mathbf{n}} Y_{jm} d\Omega,$$

and since $\Delta_{\mathbf{n}} Y_{jm} = -j(j+1)Y_{jm}$, we recover (7.5). The normalization of the vectors $\mathbf{Y}_{jm}^{(m)}$ reduces to an identical integral.

We remark that the spherical vectors (7.4) could also have been obtained without the direct verification of equations (7.1) carried out above — simply on the basis of general transformation-property arguments. Such reasoning led us in the preceding section to conclude that a vector function of the form $\mathbf{n}\varphi$ corresponds to a value of j equal to

¹⁴The operator $\nabla_{\mathbf{n}} \equiv |\mathbf{k}| \nabla_{\mathbf{k}}$ acts on functions that depend only on the direction of $\mathbf{n}$. In spherical coordinates it has just two components:

$$\nabla_{\mathbf{n}} = \left(\frac{\partial}{\partial \theta}, \frac{1}{\sin \theta} \frac{\partial}{\partial \varphi} \right).$$

The operator denoted below by $\Delta_{\mathbf{n}}$ is the angular part of the Laplacian:

$$\Delta_{\mathbf{n}} = \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \sin \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2}.$$

the order of the spherical harmonics appearing in φ ; setting $\varphi = Y_{jm}$ further fixes the projection m . In this way one immediately arrives at the spherical vectors $\mathbf{Y}_{jm}^{(1)}$. The transformation-property arguments of § 6, however, remain valid if the factor $\mathbf{n}$ in the product $\mathbf{n}\varphi$ is replaced by $\nabla_{\mathbf{n}}$ or by $[\mathbf{n}\nabla_{\mathbf{n}}]$, yielding in this manner the spherical vectors of the other two types.

Let us return to the photon wave functions. For an electric-type photon (Ej) the vector $\mathbf{A}(\mathbf{k})$ must have parity $(-1)^j$. Among the spherical vectors, $\mathbf{Y}_{jm}^{(e)}$ and $\mathbf{Y}_{jm}^{(l)}$ possess this parity; of the two, however, only the former satisfies the transversality condition. For a magnetic-type photon (Mj) the vector $\mathbf{A}(\mathbf{k})$ must have parity $(-1)^{j+1}$; only the spherical vector $\mathbf{Y}_{jm}^{(m)}$ has this parity. The wave functions of a photon with definite angular momentum j , projection m , and energy ω are therefore

$$\mathbf{A}_{\omega jm}(\mathbf{k}) = \frac{4\pi^2}{\omega^{3/2}} \delta(|\mathbf{k}| - \omega) \mathbf{Y}_{jm}(\mathbf{n}), \quad (7.6)$$

where one should write $\mathbf{Y}_{jm}^{(e)}$ or $\mathbf{Y}_{jm}^{(m)}$ for $\mathbf{Y}_{jm}$ according as the photon is of electric or magnetic type; the specified energy is incorporated through the factor $\delta(|\mathbf{k}| - \omega)$. The functions (7.6) are normalized by the condition

$$\frac{1}{(2\pi)^4} \int \omega \omega' \mathbf{A}_{\omega' j' m'}^*(\mathbf{k}) \mathbf{A}_{\omega jm}(\mathbf{k}) d^3 k = \omega \delta(\omega' - \omega) \delta_{jj'} \delta_{mm'}. \quad (7.7)$$

For the coordinate-representation wave functions, condition (7.7) is equivalent to¹⁵

$$\frac{1}{4\pi} \int \left\{ \mathbf{E}_{\omega' j' m'}^*(\mathbf{r}) \mathbf{E}_{\omega jm}(\mathbf{r}) + \mathbf{H}_{\omega' j' m'}^*(\mathbf{r}) \mathbf{H}_{\omega jm}(\mathbf{r}) \right\} d^3 x = \omega \delta(\omega' - \omega) \delta_{jj'} \delta_{mm'}. \quad (7.8)$$

Indeed, when written in terms of potentials, the integral on the left-hand side takes the form

$$\frac{1}{2\pi} \int \mathbf{A}_{\omega' j' m'}^*(\mathbf{r}) \mathbf{A}_{\omega jm}(\mathbf{r}) \omega' \omega d^3 x.$$

One must substitute here the Fourier expansions

$$\begin{aligned} \mathbf{A}_{\omega jm}(\mathbf{r}) &= \int \mathbf{A}_{\omega jm}(\mathbf{k}) e^{i\mathbf{k}\mathbf{r}} \frac{d^3 k}{(2\pi)^3}, \\ \mathbf{A}_{\omega' j' m'}^*(\mathbf{r}) &= \int \mathbf{A}_{\omega' j' m'}^*(\mathbf{k}') e^{-i\mathbf{k}'\mathbf{r}} \frac{d^3 k'}{(2\pi)^3}. \end{aligned} \quad (7.9)$$

After that, integration over $d^3 x$ produces the delta function $(2\pi)^3 \delta(\mathbf{k}' - \mathbf{k})$, which is removed by the integration over $d^3 k'$, bringing the integral to the form (7.7).

Up to this point we have assumed the transverse gauge for the potentials, in which the scalar potential $\Phi = 0$. In various applications, however, other gauge choices for a spherical wave may prove more convenient. An admissible gauge transformation of the potentials in the momentum representation consists of the substitution

$$\mathbf{A} \rightarrow \mathbf{A} + \mathbf{n}f(\mathbf{k}), \quad \Phi \rightarrow \Phi + f(\mathbf{k}),$$

¹⁵This condition is of the same type as (2.22). The appearance of the factor $\delta(\omega' - \omega)$ on the right-hand side is due to the fact that the field (a spherical wave) is being considered in all of infinite space rather than in a finite volume $V = 1$.

with $f(\mathbf{k})$ an arbitrary function. We choose it here so that the new potentials are still expressed through the same spherical harmonics and continue to have definite parity. For an electric-type photon these requirements restrict the potentials to the following form:

$$\begin{aligned} \mathbf{A}_{\omega jm}^{(e)}(\mathbf{k}) &= \frac{4\pi^2}{\omega^{3/2}} \delta(|\mathbf{k}| - \omega) (\mathbf{Y}_{jm}^{(e)} + C \mathbf{n} Y_{jm}), \\ \Phi_{\omega jm}^{(e)}(\mathbf{k}) &= \frac{4\pi^2}{\omega^{3/2}} \delta(|\mathbf{k}| - \omega) C Y_{jm}, \end{aligned} \quad (7.10)$$

where C is an arbitrary constant. For a magnetic-type photon, on the other hand, such an addition to $\mathbf{A}^{(m)}(\mathbf{k})$ would destroy its definite parity, so under the same conditions the choice (7.6) is uniquely fixed.

The probability that a photon of given angular momentum and parity is detected traveling in the direction $\mathbf{n}$ within the solid-angle element do equals, by (3.5) and (7.6),

$$w(\mathbf{n})do = |\mathbf{Y}_{jm}^{(e)}|^2 do. \quad (7.11)$$

We have written the expression for an E -type photon. Since, however, $|\mathbf{Y}_{jm}^{(m)}|^2 = |\mathbf{Y}_{jm}^{(e)}|^2$, the probability distributions $w(\mathbf{n})$ for photons of both types are identical.

The squared modulus $|\mathbf{Y}_{jm}^{(e)}|^2$ is independent of the azimuthal angle φ (the factors $e^{\pm im\varphi}$ in the spherical harmonics cancel). The probability distribution $w(\mathbf{n})$ is therefore axially symmetric about the z axis. Furthermore, since each spherical vector has definite parity, the squared moduli are even under inversion, i.e. under the replacement $\theta \rightarrow \pi - \theta$; this means that the function $w(\theta)$, when expanded in Legendre polynomials, contains only polynomials of even order. The determination of the expansion coefficients reduces to computing integrals of products of three spherical harmonics followed by summation over components. Both steps are carried out with the aid of formulas derived in III, §§ 107–108, and lead to the following result:

$$\begin{aligned} w(\theta) &= (-1)^{m+1} \frac{(2j+1)^2}{4\pi} \sum_{n=0}^{\infty} (4n+1) \begin{pmatrix} j & j & 2n \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} j & j & 2n \\ m & -m & 0 \end{pmatrix} \times \\ &\quad \times \left\{ \begin{matrix} j & j & 1 \\ j & j & 2n \end{matrix} \right\} P_{2n}(\cos \theta). \end{aligned} \quad (7.12)$$

Finally, we present expressions for the components of the spherical vectors as expansions in spherical harmonics. We employ the “spherical components” of a vector $\mathbf{f}$, defined in accordance with III, § 107; the components f_λ of the vector $\mathbf{f}$ are

$$f_0 = if_z, \quad f_{+1} = -\frac{i}{\sqrt{2}}(f_x + if_y), \quad f_{-1} = \frac{i}{\sqrt{2}}(f_x - if_y). \quad (7.13)$$

Introducing “circular basis vectors”:

$$\mathbf{e}^{(0)} = i\mathbf{e}^{(z)}, \quad \mathbf{e}^{(+1)} = -\frac{i}{\sqrt{2}}(\mathbf{e}^{(x)} + i\mathbf{e}^{(y)}), \quad \mathbf{e}^{(-1)} = \frac{i}{\sqrt{2}}(\mathbf{e}^{(x)} - i\mathbf{e}^{(y)}) \quad (7.14)$$

($\mathbf{e}^{(x,y,z)}$ being the unit vectors along the x, y, z axes), we have

$$\mathbf{f} = \sum_{\lambda} (-1)^{1-\lambda} f_{-\lambda} \mathbf{e}^{(\lambda)}, \quad f_{\lambda} = (-1)^{1-\lambda} \mathbf{f} \mathbf{e}^{(-\lambda)*} = \mathbf{f} \mathbf{e}^{(\lambda)}. \quad (7.15)$$

The spherical components of the spherical vectors are expressed through $3j$ -symbols and spherical harmonics by the following formulas:

$$\begin{aligned} (-1)^{j+m+\lambda+1} (\mathbf{Y}_{jm}^{(e)})_{\lambda} &= -\sqrt{j} \begin{pmatrix} j+1 & 1 & j \\ m+\lambda & -\lambda & -m \end{pmatrix} Y_{j+1,m+\lambda} \\ &\quad + \sqrt{j+1} \begin{pmatrix} j-1 & 1 & j \\ m+\lambda & -\lambda & -m \end{pmatrix} Y_{j-1,m+\lambda}, \\ (-1)^{j+m+\lambda+1} (\mathbf{Y}_{jm}^{(m)})_{\lambda} &= -\sqrt{2j+1} \begin{pmatrix} j & 1 & j \\ m+\lambda & -\lambda & -m \end{pmatrix} Y_{j,m+\lambda}, \\ (-1)^{j+m+\lambda+1} (\mathbf{Y}_{jm}^{(l)})_{\lambda} &= \sqrt{j+1} \begin{pmatrix} j+1 & 1 & j \\ m+\lambda & -\lambda & -m \end{pmatrix} Y_{j+1,m+\lambda} \\ &\quad + \sqrt{j} \begin{pmatrix} j-1 & 1 & j \\ m+\lambda & -\lambda & -m \end{pmatrix} Y_{j-1,m+\lambda}. \end{aligned} \quad (7.16)$$

These formulas are derived as follows. Each of the three spherical vectors has the form $\mathbf{Y}_{jm} = \mathbf{a} Y_{jm}$, where $\mathbf{a}$ is one of the three vectors (7.3). Therefore

$$\mathbf{Y}_{jm} = \sum_{lm'} \langle lm' | \mathbf{a} | jm \rangle Y_{lm'},$$

so that one must determine the matrix elements of the vectors $\mathbf{a}$ between eigenstates of the orbital angular momentum. From formula (107.6) of III it follows that

$$\langle lm' | a_{\lambda} | jm \rangle = i(-1)^{j_{\max}-m'} \begin{pmatrix} l & 1 & j \\ -m' & \lambda & m \end{pmatrix} \langle l || a || j \rangle,$$

where $j_{\max}$ is the larger of the numbers l and j . It therefore suffices to know the nonvanishing reduced matrix elements $\langle l || a || j \rangle$. The relevant formulas are:

$$\begin{aligned} \langle l-1 || n || l \rangle &= \langle l || n || l-1 \rangle^* = i\sqrt{l}, \\ \langle l || \nabla_{\mathbf{n}} || l-1 \rangle &= i(l-1)\sqrt{l}, \\ \langle l-1 || \nabla_{\mathbf{n}} || l \rangle &= i(l+1)\sqrt{l}, \\ \langle l || [\mathbf{n} \nabla_{\mathbf{n}}] || l \rangle &= i\sqrt{l(l+1)(2l+1)}. \end{aligned} \quad (7.17)$$

§8. Polarization of the photon

The polarization vector $\mathbf{e}$ plays for the photon the role of the “spin part” of the wave function (with the reservations expressed in § 6 regarding the notion of photon spin).

The various cases arising for the polarization of a photon are in no way distinct from the possible types of polarization of a classical electromagnetic wave (see II, § 48).

An arbitrary polarization $\mathbf{e}$ can be represented as a superposition of two mutually orthogonal polarizations $\mathbf{e}^{(1)}$ and $\mathbf{e}^{(2)}$ ($\mathbf{e}^{(1)}\mathbf{e}^{(2)*} = 0$), chosen in some definite manner. In the decomposition

$$\mathbf{e} = e_1\mathbf{e}^{(1)} + e_2\mathbf{e}^{(2)} \quad (8.1)$$

the squared moduli of the coefficients e_1 and e_2 give the probability that the photon has polarization $\mathbf{e}^{(1)}$ or $\mathbf{e}^{(2)}$.

As these two basis polarizations one may choose two mutually perpendicular linear polarizations. Alternatively, any polarization can be decomposed into two circular ones with opposite senses of rotation. We shall denote the right and left circular polarization vectors by $\mathbf{e}^{(+1)}$ and $\mathbf{e}^{(-1)}$ respectively; in a coordinate system $\xi\eta\zeta$ with the ζ axis along the photon's direction $\mathbf{n} = \mathbf{k}/\omega$,

$$\mathbf{e}^{(+1)} = -\frac{i}{\sqrt{2}}(\mathbf{e}^{(\xi)} + i\mathbf{e}^{(\eta)}), \quad \mathbf{e}^{(-1)} = \frac{i}{\sqrt{2}}(\mathbf{e}^{(\xi)} - i\mathbf{e}^{(\eta)}). \quad (8.2)$$

The existence of two distinct polarizations of the photon (at a given momentum) means, in other words, that every eigenvalue of the momentum is doubly degenerate. This circumstance is intimately connected with the fact that the photon mass equals zero.

For a freely moving particle of nonvanishing mass a rest frame always exists. It is clear that the intrinsic symmetry properties of the particle as such manifest themselves precisely in this frame. The symmetry to be considered is with respect to all possible rotations about the center, that is, with respect to the full spherical symmetry group. The quantity characterizing the symmetry properties of the particle under this group is its spin s , which determines the multiplicity of the degeneracy (the number $2s + 1$ of distinct wave functions that transform into one another). In particular, a particle whose wave function is a vector (three components) has spin 1.

For a particle of zero mass, however, no rest frame exists — in every reference frame it travels at the speed of light. For such a particle there is always a preferred spatial direction — the direction of the momentum vector $\mathbf{k}$ (the ζ axis). In this situation the full three-dimensional rotation symmetry is evidently absent, and one may speak only of axial symmetry about the distinguished axis.

Under axial symmetry only the *helicity* of the particle is conserved — the projection of angular momentum onto the ζ axis; we denote it by λ ¹⁶. If one further demands symmetry under reflections in planes passing through the ζ axis, then states differing in the sign of λ are mutually degenerate; for $\lambda \neq 0$ there is therefore a twofold degeneracy¹⁷. The state of a photon with a definite momentum belongs to precisely this type of twofold-degenerate state. It is described by a “spin” wave function that is a vector $\mathbf{e}$ in the $\xi\eta$ plane; the two components of this vector transform into each other under all rotations about the ζ axis and under reflections in planes containing that axis.

The various polarization cases of a photon are in a definite correspondence with the possible values of its helicity. This correspondence can be established using formulas III,

¹⁶In contrast to the projection m of the angular momentum onto a fixed direction (the z axis) in space, discussed in the preceding section.

¹⁷We note that the electronic terms of a diatomic molecule are classified in exactly the same way (see III, § 78).

(57.9), which relate the components of a vector wave function to the components of the equivalent rank-2 spinor¹⁸. The projections $\lambda = +1$ and -1 correspond to vectors $\mathbf{e}$ that have only the component $e_\xi - ie_\eta$ or $e_\xi + ie_\eta$ nonzero, i.e. $\mathbf{e} = \mathbf{e}^{(+1)}$ or $\mathbf{e} = \mathbf{e}^{(-1)}$ respectively. In other words, $\lambda = +1$ and -1 correspond to the right and left circular polarizations of the photon (in § 16 the same result will be derived by directly computing the eigenfunctions of the spin-projection operator). Thus the projection of the photon's angular momentum onto the direction of its motion can take only the two values (± 1) ; the value 0 is not possible.

A photon state with definite momentum and polarization is a pure state (in the sense explained in III, § 14); it is described by a wave function and constitutes a complete quantum-mechanical specification of the particle (photon) state. There also exist “mixed” photon states corresponding to a less complete description, carried out not by a wave function but only by a density matrix.

Let us examine a photon state that is mixed in its polarization but possesses a definite momentum $\mathbf{k}$. Such a state (called a state of *partial polarization*) admits a “coordinate” wave function.

The polarization density matrix of a photon is a second-rank tensor $\rho_{\alpha\beta}$ defined in the plane orthogonal to $\mathbf{n}$ (the $\xi\eta$ plane; the indices α, β assume only two values). This tensor is Hermitian:

$$\rho_{\alpha\beta} = \rho_{\beta\alpha}^*, \quad (8.3)$$

and normalized by the condition

$$\rho_{\alpha\alpha} = \rho_{11} + \rho_{22} = 1. \quad (8.4)$$

By virtue of (8.3) the diagonal components ρ_{11} and ρ_{22} are real, and one determines the other through condition (8.4). The component ρ_{12} is complex, while $\rho_{21} = \rho_{12}^*$. Altogether, then, the density matrix is characterized by three real parameters.

Once the polarization density matrix is known, one can find the probability that the photon has any given definite polarization $\mathbf{e}$. This probability is obtained by “projecting” the tensor $\rho_{\alpha\beta}$ onto the direction of the vector $\mathbf{e}$, i.e. it equals the quantity

$$\rho_{\alpha\beta} e_\alpha^* e_\beta. \quad (8.5)$$

Thus the components ρ_{11} and ρ_{22} are the probabilities of linear polarizations along the ξ and η axes. Projection onto the vectors (8.2) yields the probabilities of the two circular polarizations:

$$\frac{1}{2} [1 \pm i(\rho_{12} - \rho_{21})]. \quad (8.6)$$

In both form and substance the properties of the tensor $\rho_{\alpha\beta}$ coincide with those of the tensor $J_{\alpha\beta}$ that describes partially polarized light in the classical theory (see II, § 50). Let us recall some of these properties here.

For a pure state of definite polarization $\mathbf{e}$, the tensor $\rho_{\alpha\beta}$ reduces to products of the components of the vector $\mathbf{e}$:

$$\rho_{\alpha\beta} = e_\alpha e_\beta^*. \quad (8.7)$$

¹⁸Recall that the wave-function components, viewed as probability amplitudes for the various values of the angular-momentum projection of the particle (which is what is meant here), correspond to the contravariant components of the spinor.

In this case the determinant $|\rho_{\alpha\beta}| = 0$. In the opposite case of an unpolarized photon, all polarization directions are equally probable, i.e.

$$\rho_{\alpha\beta} = \delta_{\alpha\beta}/2, \quad (8.8)$$

and then $|\rho_{\alpha\beta}| = 1/4$.

For the general case of partial polarization it is convenient to introduce three real Stokes parameters ξ_1, ξ_2, ξ_3 ¹⁹; the density matrix is then expressible as

$$\rho_{\alpha\beta} = \frac{1}{2} \begin{pmatrix} 1 + \xi_3 & \xi_1 - i\xi_2 \\ \xi_1 + i\xi_2 & 1 - \xi_3 \end{pmatrix}. \quad (8.9)$$

All three parameters range between -1 and $+1$. In the unpolarized state $\xi_1 = \xi_2 = \xi_3 = 0$; for a completely polarized photon $\xi_1^2 + \xi_2^2 + \xi_3^2 = 1$.

The parameter ξ_3 characterizes linear polarization along the ξ or η axis; the probability of linear polarization along these axes is $(1 + \xi_3)/2$ and $(1 - \xi_3)/2$ respectively. The values $\xi_3 = +1$ or -1 therefore correspond to complete polarization in these directions.

The parameter ξ_1 characterizes linear polarization along directions making an angle $\varphi = \pi/4$ or $\varphi = -\pi/4$ with the ξ axis. The probability of linear polarization in these directions is $(1 + \xi_1)/2$ or $(1 - \xi_1)/2$; this is easily verified by projecting the tensor $\rho_{\alpha\beta}$ onto the directions $\mathbf{e} = (1, \pm 1)/\sqrt{2}$.

Lastly, the parameter ξ_2 represents the degree of circular polarization; by (8.6), the probability of right or left circular polarization is $(1 + \xi_2)/2$ or $(1 - \xi_2)/2$ respectively. Because these two polarizations answer to helicities $\lambda = \pm 1$, it is evident that in general ξ_2 coincides with the mean helicity of the photon. Note further that for a pure state with polarization $\mathbf{e}$,

$$\xi_2 = i[\mathbf{e}\mathbf{e}^*]\mathbf{n}. \quad (8.10)$$

Recall (see II, § 50) that the quantities ξ_2 and $\sqrt{\xi_1^2 + \xi_3^2}$ are invariant under Lorentz transformations.

In what follows we shall also need to know how the Stokes parameters transform under time reversal. One readily verifies that they are invariant under this operation. Since the result clearly does not depend on whether the polarization state is pure or mixed, it is enough to check invariance for a pure state. In quantum mechanics, time reversal replaces the wave function with its complex conjugate (see III, § 18). For a plane-polarized wave the corresponding substitution is²⁰

$$\mathbf{k} \rightarrow -\mathbf{k}, \quad \mathbf{e} \rightarrow -\mathbf{e}^*. \quad (8.11)$$

Under this transformation the symmetric part of the density matrix, $(1/2)(e_\alpha e_\beta^* + e_\beta e_\alpha^*)$, and hence the parameters ξ_1 and ξ_3 remain unchanged. The invariance of ξ_2 under the

¹⁹The notation for the parameters should not be confused with that for the ξ axis!

²⁰The extra sign reversal of $\mathbf{e}$ is due to the fact that time reversal changes the sign of the electromagnetic vector potential. The scalar potential is unaffected; consequently, for the 4-vector e time reversal takes the form

$$(e_0, \mathbf{e}) \rightarrow (e_0^*, -\mathbf{e}^*). \quad (8.11a)$$

same transformation is apparent from (8.10); it is also obvious already from the physical meaning of ξ_2 as the mean helicity. Indeed, helicity is the projection of the angular momentum $\mathbf{j}$ onto the direction $\mathbf{n}$, that is, the product $\mathbf{j}\mathbf{n}$; time reversal reverses the sign of both these vectors.

For the calculations that follow, the photon density matrix must be cast in four-dimensional form, that is, expressed as a certain 4-tensor $\rho_{\mu\nu}$. When the photon is polarized and described by a 4-vector e_μ , a natural definition of this tensor is

$$\rho_{\mu\nu} = e_\mu e_\nu^*. \quad (8.12)$$

In the three-dimensionally transverse gauge, $e = (0, \mathbf{e})$, if one of the spatial coordinate axes is chosen along $\mathbf{n}$, the nonvanishing components of this 4-tensor coincide with (8.7).

For an unpolarized photon the three-dimensionally transverse gauge yields a tensor $\rho_{\mu\nu}$ with components

$$\rho_{ik} = \frac{1}{2}(\delta_{ik} - n_i n_k), \quad \rho_{0i} = \rho_{i0} = \rho_{00} = 0 \quad (8.13)$$

(if one of the axes coincides with the direction of $\mathbf{n}$, we return to (8.8)). Using the tensor $\rho_{\mu\nu}$ directly in this three-dimensional form is, however, inconvenient. We may instead perform a gauge transformation; for the density matrix this is a transformation of the type

$$\rho_{\mu\nu} \rightarrow \rho_{\mu\nu} + \chi_\mu k_\nu + \chi_\nu k_\mu, \quad (8.14)$$

where χ_μ are arbitrary functions. Setting

$$\chi_0 = -1/(4\omega), \quad \chi_i = k_i/(4\omega^2)$$

we obtain in place of (8.13) the simple four-dimensional expression

$$\rho_{\mu\nu} = -g_{\mu\nu}/2. \quad (8.15)$$

The four-dimensional representation of the density matrix of a partially polarized photon is readily obtained by first rewriting the two-dimensional tensor (8.9) in three-dimensional form. In the three-dimensionally transverse gauge: $e^{(1)} = (0, \mathbf{e}^{(1)})$, $e^{(2)} = (0, \mathbf{e}^{(2)})$. The four-dimensional photon density matrix is therefore

$$\begin{aligned} \rho_{\mu\nu} = & (1/2)(e_\mu^{(1)} e_\nu^{(1)} + e_\mu^{(2)} e_\nu^{(2)}) + (\xi_1/2)(e_\mu^{(1)} e_\nu^{(2)} + e_\mu^{(2)} e_\nu^{(1)}) - \\ & - (i\xi_2/2)(e_\mu^{(1)} e_\nu^{(2)} - e_\mu^{(2)} e_\nu^{(1)}) + (\xi_3/2)(e_\mu^{(1)} e_\nu^{(1)} - e_\mu^{(2)} e_\nu^{(2)}). \end{aligned} \quad (8.17)$$

The convenience of any particular choice of the 4-vectors $e^{(1)}$, $e^{(2)}$ depends on the specific conditions of the problem at hand.

It should be borne in mind that the conditions (8.16) do not fix the choice of $e^{(1)}$ and $e^{(2)}$ uniquely. If any 4-vector e_μ satisfies these conditions, so does any 4-vector of the form $e_\mu + \chi k_\mu$ (since $k^2 = 0$). This ambiguity is tied to the gauge freedom of the density matrix.

The first term in (8.17) corresponds to the unpolarized state. It could therefore be replaced, by (8.15), with $-g_{\mu\nu}/2$. Such a replacement is again equivalent to a gauge

transformation. When working with 4-tensors of the form (8.17), decomposed in two independent 4-vectors, it is useful to adopt the following formal device. Writing (8.17) as

$$\rho_{\mu\nu} = \sum_{a,b=1}^2 \rho^{(ab)} e_{\mu}^{(a)} e_{\nu}^{(b)},$$

we represent the coefficients $\rho^{(ab)}$ as a 2×2 matrix

$$\rho = \begin{pmatrix} \rho^{(11)} & \rho^{(12)} \\ \rho^{(21)} & \rho^{(22)} \end{pmatrix}.$$

Any Hermitian 2×2 matrix can be expanded over four linearly independent 2×2 matrices — the Pauli matrices $\sigma_x, \sigma_y, \sigma_z$ and the identity matrix 1:

$$\rho = (1/2)(1 + \boldsymbol{\xi} \boldsymbol{\sigma}), \quad \boldsymbol{\xi} = (\xi_1, \xi_2, \xi_3), \quad (8.18)$$

as is easily verified by direct comparison with (8.17) using the well-known explicit forms of the Pauli matrices (18.5) (grouping the three quantities ξ_1, ξ_2, ξ_3 into a “vector” $\boldsymbol{\xi}$ is of course purely formal and serves only to shorten the notation).

The requisite generalization is achieved by replacing these 3-vectors by spacelike real 4-vectors $e^{(1)}, e^{(2)}$, orthogonal to each other and to the photon 4-momentum k :

$$e^{(1)2} = e^{(2)2} = -1, \quad e^{(1)} e^{(2)} = 0, \quad e^{(1)} k = e^{(2)} k = 0. \quad (8.16)$$

Problem

Write the photon density matrix in the representation whose “coordinate axes” are the circular basis vectors (8.2).

Solution. The components of the tensor $\rho'_{\alpha\beta}$ in the new axes ($\alpha, \beta = \pm 1$) are obtained by projecting the tensor (8.9) onto the basis vectors (8.2):

$$\rho'_{11} = \rho_{\alpha\beta} e_{\alpha}^{(+1)*} e_{\beta}^{(+1)}, \quad \rho'_{1,-1} = \rho_{\alpha\beta} e_{\alpha}^{(+1)*} e_{\beta}^{(-1)}, \quad \dots,$$

$$\rho' = \frac{1}{2} \begin{pmatrix} 1 + \xi_2 & -\xi_3 + i\xi_1 \\ -\xi_3 - i\xi_1 & 1 - \xi_2 \end{pmatrix}.$$

§9. The two-photon system

Arguments analogous to those developed in § 6 make it possible to count the number of allowed states in the more involved case of a system of two photons as well (*L. Landau*, 1948).

Let us work in the center-of-inertia frame of the two photons, where $\mathbf{k}_1 = -\mathbf{k}_2 \equiv \mathbf{k}$ ²¹. In the momentum representation the wave function of the pair can be cast as a three-dimensional rank-2 tensor $A_{ik}(\mathbf{n})$ built bilinearly from the components of the individual

²¹ A center-of-inertia frame can be found in every case except when the two photons propagate in the same direction. For such a pair the total momentum $\mathbf{k}_1 + \mathbf{k}_2$ and total energy $\omega_1 + \omega_2$ are connected by exactly the same relation as for one photon, so no frame with $\mathbf{k}_1 + \mathbf{k}_2 = 0$ exists.

photon vector wave functions; each of its two indices refers to one photon ($\mathbf{n}$ denotes the unit vector in the direction of $\mathbf{k}$). The transversality requirement for each photon is encoded in the orthogonality of A_{ik} to $\mathbf{n}$:

$$A_{il}n_l = 0, \quad A_{lk}n_l = 0. \quad (9.1)$$

Interchanging the two photons amounts to transposing the indices of the tensor A_{ik} together with a simultaneous reversal of the sign of $\mathbf{n}$. Because photons obey Bose statistics,

$$A_{ik}(-\mathbf{n}) = A_{ki}(\mathbf{n}). \quad (9.2)$$

The tensor A_{ik} is, in general, not symmetric in its indices. We decompose it into its symmetric part (s_{ik}) and antisymmetric part (a_{ik}): $A_{ik} = s_{ik} + a_{ik}$. Each of these parts must individually satisfy the relation (9.2) (as well as the orthogonality conditions (9.1)). This gives

$$s_{ik}(-\mathbf{n}) = s_{ik}(\mathbf{n}), \quad (9.3)$$

$$a_{ik}(-\mathbf{n}) = -a_{ik}(\mathbf{n}). \quad (9.4)$$

Spatial inversion does not by itself reverse the sign of the components of a second-rank tensor, but it does reverse the sign of $\mathbf{n}$. From (9.3) it is therefore apparent that the wave function s_{ik} is symmetric under inversion, i.e. it corresponds to even-parity states of the photon system; the wave function a_{ik} , on the other hand, corresponds to odd-parity states.

An antisymmetric second-rank tensor is equivalent (dual) to an axial vector $\mathbf{a}$ whose components are expressed via the tensor components as $a_i = \frac{1}{2}e_{ikl}a_{kl}$, where e_{ikl} is the antisymmetric unit tensor (see II, § 6). The orthogonality of the tensor a_{kl} to the vector $\mathbf{n}$ implies that the vectors $\mathbf{a}$ and $\mathbf{n}$ are parallel²². One may therefore write $\mathbf{a} = \mathbf{n}\varphi(\mathbf{n})$, where φ is a scalar; by (9.4) we must have $\mathbf{a}(-\mathbf{n}) = -\mathbf{a}(\mathbf{n})$, and hence

$$\varphi(-\mathbf{n}) = \varphi(\mathbf{n}).$$

This relation means that the scalar φ can be built linearly only from spherical harmonics of even order L (including zero).

We see that the antisymmetric tensor a_{ik} , by its transformation properties (under rotations), is equivalent to a single scalar (cf. the footnote on p. 34). Assigning to the latter a “spin” of 0, we find that the angular momentum of the state is $J = L$. Thus the tensor a_{ik} corresponds to odd-parity states of the photon system with even total angular momentum J .

Let us turn to the symmetric tensor s_{ik} . Since it is even under reversal of the sign of $\mathbf{n}$, it describes even-parity states of the photon system. It follows likewise that all components of s_{ik} are expressed through spherical harmonics of even order L (including $L = 0$). An arbitrary symmetric second-rank tensor can be decomposed, as is well known, into a scalar (s_{ii}) and a traceless symmetric tensor (s'_{ik}) with $s'_{ii} = 0$.

²²We have: $a_{ik} = e_{ikl}a_l$, and the orthogonality condition yields

$$a_{ik}n_k = e_{ikl}a_l n_k = [\mathbf{n}\mathbf{a}]_i = 0.$$

The scalar s_{ii} is assigned “spin” 0, so the angular momentum of the corresponding states is $J = L$, which is even. The tensor s'_{ik} corresponds to “spin” 2 (see III, § 57). Composing this “spin” with the even “orbital angular momentum” L according to the addition rule for angular momenta, we find that for a given even $J \neq 0$ there are three states (with $L = J \pm 2, J$), while for odd $J \neq 1$ there are two states (with $L = J \pm 1$). The exceptions are $J = 0$ with a single state ($L = 2$) and $J = 1$ with a single state ($L = 2$).

In this counting, however, we have not yet accounted for the orthogonality of the tensor s_{ik} to the vector $\mathbf{n}$. One must therefore subtract from the number of states obtained those corresponding to a symmetric second-rank tensor that is “parallel” to the vector $\mathbf{n}$. Such a tensor (denote it s''_{ik}) can be written as

$$s''_{ik} = n_i b_k + n_k b_i,$$

where $\mathbf{b}$ is some vector. By (9.3) this vector must satisfy $\mathbf{b}(-\mathbf{n}) = -\mathbf{b}(\mathbf{n})$. The tensor s''_{ik} responsible for the “extra” states is thus equivalent to an odd-parity vector. This vector must consequently be expressed through spherical harmonics of odd orders L only. Noting further that a vector corresponds to “spin” 1, we find that for each even $J \neq 0$ there are two states (with $L = J \pm 1$), and for each odd J a single state (with $L = J$); the special case $J = 0$ has one state (with $L = 1$).

Collecting all the results obtained, we arrive at the following table giving the number of possible even- and odd-parity states of a system of two photons (with vanishing total momentum) for the various values of the total angular momentum J :

J	Even	Odd
0	1	1
1	—	—
$2k$	2	1
$2k + 1$	1	—

(9.5)

(k being a positive integer other than zero). We observe that for odd J no odd-parity states exist, while the value $J = 1$ is altogether excluded²³.

The wave function of the two-photon system A_{ik} determines the correlation between their polarizations. The probability that both photons simultaneously possess definite polarizations $\mathbf{e}_1$ and $\mathbf{e}_2$ is proportional to

$$A_{ik} e_{1i}^* e_{2k}^*.$$

In other words, if the polarization $\mathbf{e}_1$ of one photon is specified, then the polarization of the second photon, $\mathbf{e}_2$, is proportional to

$$e_{2k} \propto A_{ik} e_{1i}^*. \quad (9.6)$$

In odd-parity states of the system, A_{ik} coincides with the antisymmetric tensor a_{ik} . In that case

$$\mathbf{e}_2 \mathbf{e}_1^* \propto a_{ik} e_{1i}^* e_{1k}^* = 0,$$

²³For an alternative derivation of these results, see problem 1 to § 69.

i.e. the polarizations of the two photons are mutually orthogonal. For linear polarization this signifies that their directions are perpendicular, while for circular polarization it means opposite senses of rotation.

The even-parity state with $J = 0$ is described by a symmetric tensor reducing to a scalar,

$$s_{ik} = \text{const} \cdot (\delta_{ik} - n_i n_k).$$

From (9.6) we then obtain $\mathbf{e}_1 = \mathbf{e}_2^*$. For linear polarization this means that their directions are parallel, whereas for circular polarization it means opposite senses of rotation. The latter fact is self-evident: when $J = 0$ the sum of the angular-momentum projections of the two photons onto any common direction $\mathbf{k}$ must in all cases vanish (the projections onto the opposite directions $\mathbf{k}_1$ and $\mathbf{k}_2$, i.e. the helicities, are accordingly equal).

Bosons

§ 10. Wave equation for particles with spin 0

Chapter I demonstrated the construction of a quantum description for the free electromagnetic field, proceeding from the known classical-limit properties of the field and relying on the apparatus of ordinary quantum mechanics. The scheme of field description as a photon system obtained in this way retains many features that carry over to the relativistic quantum-theoretical treatment of particles as well.

The electromagnetic field is a system with an infinite number of degrees of freedom. For it there is no conservation law for the number of particles (photons), and among its possible states are those with any number of particles¹. However, the same property must, generally speaking, also hold in relativistic theory for systems of any particles. The conservation of particle number in non-relativistic theory is tied to the conservation of mass: the sum of the masses (rest masses) of the particles does not change during their interaction; conservation of the total mass in a system of electrons, say, implies the constancy of their number as well. In relativistic mechanics, however, no law of mass conservation exists; only the total energy of the system (which includes the rest energies of the particles) must be conserved. For this reason the particle number need no longer be conserved, and consequently every relativistic particle theory must be a theory of systems possessing an infinite number of degrees of freedom. In other words, such a particle theory acquires the character of a field theory.

The adequate mathematical apparatus for treating systems whose particle number varies is that of second quantization (see III, §§ 64, 65). In the quantum treatment of the electromagnetic field the 4-potential $\hat{A}$ serves as the second-quantization operator. It is built from the (coordinate-space) wave functions of the individual particles (photons) together with their creation and annihilation operators. A completely analogous role in the description of a system of particles belongs to the operator of the quantized wave function. Its construction requires, first and foremost, knowledge of the wave-function form for a single free particle and of the equation governing this function.

It should be stressed that the notion of free-particle fields is auxiliary in character. Real particles interact, and the task of the theory lies in studying these interactions. But every interaction reduces to a collision, before and after which the system can be regarded as a collection of free particles. In § 1 it was noted that these are the only directly measurable objects. We therefore use the fields of free particles as a means for describing initial and final states.

We begin the relativistic description of free particles with the case of spin-0 particles. The mathematical simplicity of this case makes it possible to bring out most clearly the basic ideas and characteristic features of such a description.

¹In practice, of course, the photon number changes only through various interaction processes.

The state of a free particle (without spin) can be completely specified by giving its momentum $\mathbf{p}$ alone. The energy ε of the particle² satisfies $\varepsilon^2 = \mathbf{p}^2 + m^2$ (where m is the particle mass), or in four-dimensional form:

$$p^2 = m^2. \quad (10.1)$$

It is well known that the conservation of momentum and energy is tied to the homogeneity of space and time, that is, to symmetry with respect to any parallel translation of the 4-coordinate system. Within the quantum description, this symmetry requirement implies that under such a coordinate transformation the wave function of a particle possessing a definite 4-momentum may only acquire a phase factor (whose modulus equals unity). This requirement is met solely by an exponential with an exponent that depends linearly on the 4-coordinates. Put differently, the wave function of a free particle in a state of definite 4-momentum $p^\mu = (\varepsilon, \mathbf{p})$ must have the form of a plane wave:

$$\text{const} \cdot e^{-ipx}, \quad px = \varepsilon t - \mathbf{p}\mathbf{r} \quad (10.2)$$

(the choice of sign in the exponent is in itself a convention within relativistic theory; it is made so as to agree with the non-relativistic case).

The wave equation must possess the functions (10.2) as particular solutions for every 4-vector p obeying the condition (10.1). It must be linear—this being a manifestation of the superposition principle: every linear combination of functions (10.2) likewise describes a possible particle state and hence must also satisfy the equation. Lastly, the equation should have the lowest attainable order; raising the order would bring in extraneous solutions.

Spin is the angular momentum that a particle possesses in the frame where it is at rest. When the particle's spin equals s , the wave function in the rest frame takes the form of a three-dimensional spinor of rank $2s$. For describing the particle in an arbitrary reference frame one must cast the wave function in terms of four-dimensional objects.

A particle of spin 0 is described in its rest frame by a three-dimensional scalar. This scalar may, however, have a different four-dimensional “ancestry”: it can be a four-scalar ψ , but it can equally well be the temporal component of a timelike 4-vector ψ_μ whose only component that does not vanish in the rest frame is ψ_0 ³.

For a free particle the only operator that can enter the wave equation is the 4-momentum operator $\hat{p}$. Its components are the differentiation operators with respect to the coordinates and time:

$$\hat{p}^\mu = i\partial^\mu = \left(i\frac{\partial}{\partial t}, -i\nabla\right). \quad (10.3)$$

The wave equation must represent a differential relation between the quantities ψ and ψ_μ , effected by means of the operator $\hat{p}$. This relation must, of course, be expressed through relativistically invariant equations. Such equations are

$$m\psi_\mu = \hat{p}_\mu\psi, \quad \hat{p}^\mu\psi_\mu = m\psi, \quad (10.4)$$

²We denote the energy of an individual particle by ε , as distinct from the energy E of the system of particles.

³Or, by the same token, the temporal component of a higher-rank 4-tensor; that possibility, however, would give rise to equations of higher order.

where m is a dimensional constant characterizing the particle⁴.

Substituting ψ_μ from the first equation into the second, we obtain

$$(\hat{p}^2 - m^2)\psi = 0 \quad (10.5)$$

(*O. Klein, V. A. Fock, 1926; W. Gordon, 1927*). Written out explicitly, this equation reads

$$-\partial_\mu \partial^\mu \psi \equiv \left(-\frac{\partial^2}{\partial t^2} + \Delta \right) \psi = m^2 \psi. \quad (10.6)$$

If one inserts a plane wave (10.2) for ψ , the result is $p^2 = m^2$, showing that m is indeed the particle mass. It is worth noting that the structure of equation (10.5) could, of course, have been anticipated, since $\hat{p}^2$ is the sole scalar operator one can form from $\hat{p}$ (and for this reason every component of the wave function of a particle of arbitrary spin obeys the same equation—something we shall encounter many times below).

Thus a spin-0 particle is described essentially by a single (four-dimensional) scalar ψ obeying the second-order equation (10.5). In the first-order equations (10.4) the role of the wave function is played by the aggregate of the quantities ψ and ψ_μ , the 4-vector ψ_μ reducing to the 4-gradient of the scalar ψ . In the rest frame the particle's wave function does not depend on the (spatial) coordinates, so that the spatial components of the 4-vector ψ_μ vanish, as they should.

For carrying out second quantization it proves useful to write the particle's energy and momentum as spatial integrals of certain bilinear (in ψ and ψ^*) combinations playing the role of spatial densities of these quantities. This amounts to finding the energy-momentum tensor $T_{\mu\nu}$ that corresponds to equation (10.5). With the aid of this tensor, the energy-momentum conservation law takes the form

$$\partial_\mu T_\nu^\mu = 0. \quad (10.7)$$

Following the general rules of field theory (see II, § 32), let us write down the variational principle whose consequence would be equation (10.5). Such a principle must consist in requiring the minimality of the “action integral”

$$S = \int L d^4x \quad (10.8)$$

with respect to a certain real 4-scalar L —the Lagrangian density of the field⁵. From the scalar ψ (and the operator ∂^μ) one can construct a real scalar bilinear expression of the form

$$L = \partial_\mu \psi^* \cdot \partial^\mu \psi - m^2 \psi^* \psi, \quad (10.9)$$

where m is a dimensional constant. Treating ψ and ψ^* as independent variables describing the field (“generalized coordinates” of the field q), it is easy to verify that the Lagrange

⁴The constants m are introduced in (10.4) in such a way that ψ_μ and ψ have the same dimensionality. It would be pointless to introduce different constants m_1 and m_2 in these two equations, since they could always be made equal by redefining ψ or ψ_μ .

⁵The corresponding second-quantized operator $\hat{L}$ is called the field *Lagrangian*. For brevity of terminology we shall use this term both for the “quantized” and the “unquantized” Lagrangian density.

equations

$$\frac{\partial}{\partial x^\mu} \frac{\partial L}{\partial q_{,\mu}} = \frac{\partial L}{\partial q} \quad (10.10)$$

$(q_{,\mu} \equiv \partial_\mu q)$ do indeed coincide with the equations (10.5) for ψ and ψ^* , with m being the particle mass. Note also that the expression (10.9) is written with an overall sign such that the square of the time derivative, $|\partial\psi/\partial t|^2$, enters L with a plus sign; in the opposite case the action could not have a minimum (cf. II, § 27). The choice of the overall numerical coefficient of L is a matter of convention (and is reflected only in the normalization coefficient in ψ).

The energy-momentum tensor is now computed by the formula

$$T_\mu^\nu = \sum q_{,\mu} \frac{\partial L}{\partial q_{,\nu}} - L \delta_\mu^\nu \quad (10.11)$$

(summation over all q). Substituting (10.9), we obtain

$$T_{\mu\nu} = \partial_\mu \psi^* \cdot \partial_\nu \psi + \partial_\nu \psi^* \cdot \partial_\mu \psi - L g_{\mu\nu} \quad (10.12)$$

(these quantities are, as they should be, real, which is ensured by the reality of L). In particular,

$$T_{00} = 2 \frac{\partial \psi^*}{\partial t} \frac{\partial \psi}{\partial t} - L = \frac{\partial \psi^*}{\partial t} \frac{\partial \psi}{\partial t} + \nabla \psi^* \cdot \nabla \psi + m^2 \psi^* \psi, \quad (10.13)$$

$$T_{i0} = \frac{\partial \psi^*}{\partial t} \frac{\partial \psi}{\partial x^i} + \frac{\partial \psi^*}{\partial x^i} \frac{\partial \psi}{\partial t}. \quad (10.14)$$

The 4-momentum of the field is given by the integral

$$P_\mu = \int T_{\mu 0} d^3x, \quad (10.15)$$

i.e. T_{00} and T_{i0} play the role of the energy and momentum densities. We note that the quantity T_{00} is essentially positive.

Formula (10.13) can be used to normalize the wave function. A plane wave normalized to one particle in a volume $V = 1$ takes the form

$$\psi_p = \frac{1}{\sqrt{2\varepsilon}} e^{-ipx}. \quad (10.16)$$

For this function one has $T_{00} = \varepsilon$, whence the total energy within the volume $V = 1$ coincides with the energy of one particle.

The angular momentum, whose conservation follows from the isotropy of space, likewise admits a representation as a spatial integral; we shall, however, have no occasion to use such a representation later on.

Finally, besides the conservation laws linked directly with space-time symmetry, equations (10.4) admit one more conservation law. It is easy to verify that, by virtue of (10.4) (and the analogous equations for ψ^*), the following equation holds:

$$\partial_\mu j^\mu = 0, \quad (10.17)$$

where

$$j_\mu = m(\psi^* \psi_\mu + \psi_\mu^* \psi) = i[\psi^* \partial_\mu \psi - (\partial_\mu \psi^*) \psi]. \quad (10.18)$$

It is apparent from this that j^μ plays the role of the 4-vector current density. Equation (10.17) is then the continuity equation expressing the conservation of the quantity

$$Q = \int j_0 d^3x, \quad (10.19)$$

where

$$j_0 = j^0 = i \left(\psi^* \frac{\partial \psi}{\partial t} - \frac{\partial \psi^*}{\partial t} \psi \right). \quad (10.20)$$

Let us call attention to the fact that j_0 is not a positive-definite quantity. This circumstance alone shows that in the general case it certainly cannot be interpreted as the probability density of the spatial localization of the particle. The meaning of the conservation law expressed by equation (10.17) will be clarified in the next section.

§ 11. Particles and antiparticles

In keeping with the general procedure of second quantization, one must expand an arbitrary wave function over the eigenfunctions spanning the full set of possible free-particle states, for example over the plane waves ψ_p :

$$\psi = \sum_{\mathbf{p}} a_{\mathbf{p}} \psi_p, \quad \psi^* = \sum_{\mathbf{p}} a_{\mathbf{p}}^* \psi_p^*.$$

After that, the coefficients $a_{\mathbf{p}}$, $a_{\mathbf{p}}^*$ would have to be promoted to operators $\hat{a}_{\mathbf{p}}$, $\hat{a}_{\mathbf{p}}^*$ for annihilating and creating particles in the corresponding states⁶.

At this point, however, we immediately encounter the following fundamentally new circumstance (compared with the non-relativistic theory). In a plane wave that solves equation (10.5), the energy ε must satisfy (for a given momentum $\mathbf{p}$) only the condition $\varepsilon^2 = \mathbf{p}^2 + m^2$, so that it can take two values: $\pm\sqrt{\mathbf{p}^2 + m^2}$. Physically meaningful values of the free-particle energy can, however, be only positive ones. Yet simply discarding the negative values is inadmissible: the general solution of the wave equation is formed only by a superposition of all its independent particular solutions. This points to the need for a certain re-interpretation of the expansion coefficients of ψ and ψ^* upon second quantization.

Let us write this expansion in the form

$$\psi = \sum_{\mathbf{p}} \frac{1}{\sqrt{2\varepsilon}} a_{\mathbf{p}}^{(+)} e^{i(\mathbf{p}\mathbf{r} - \varepsilon t)} + \sum_{\mathbf{p}} \frac{1}{\sqrt{2\varepsilon}} a_{\mathbf{p}}^{(-)} e^{i(\mathbf{p}\mathbf{r} + \varepsilon t)}, \quad (11.1)$$

where the first sum contains plane waves normalized according to (10.16) with positive “frequencies,” and the second with negative ones; everywhere ε stands for the positive

⁶We label the ψ -functions by the 4-momentum index p ; functions with “negative frequency” will henceforth be denoted ψ_{-p} . The operators $\hat{a}$, $\hat{a}^*$, on the other hand, carry the three-momentum index $\mathbf{p}$, which completely specifies the state of a real particle.

quantity: $\varepsilon = +\sqrt{\mathbf{p}^2 + m^2}$. In performing second quantization, the coefficients $a_{\mathbf{p}}^{(+)}$ in the first sum are replaced in the usual way by particle annihilation operators $\hat{a}_{\mathbf{p}}$. In the second sum we observe that upon subsequent evaluation of matrix elements the time dependence of its terms will correspond not to annihilation but to creation of particles; the factor $e^{i\varepsilon t} = (e^{-i\varepsilon t})^*$ corresponds to one extra particle with energy ε in the final state (cf. the end of § 2). Accordingly, the coefficients $a_{\mathbf{p}}^{(-)}$ are replaced by creation operators $\hat{b}_{-\mathbf{p}}^+$ for certain other particles. Changing the summation variable in the second sum of (11.1) from $\mathbf{p}$ to $-\mathbf{p}$ (so that the exponential factor takes the form $e^{-i(\mathbf{p}\mathbf{r} - \varepsilon t)}$), we arrive at the ψ -operators

$$\hat{\psi} = \sum_{\mathbf{p}} \frac{1}{\sqrt{2\varepsilon}} (\hat{a}_{\mathbf{p}} e^{-i\mathbf{p}\mathbf{x}} + \hat{b}_{\mathbf{p}}^+ e^{i\mathbf{p}\mathbf{x}}), \quad \hat{\psi}^+ = \sum_{\mathbf{p}} \frac{1}{\sqrt{2\varepsilon}} (\hat{a}_{\mathbf{p}}^+ e^{i\mathbf{p}\mathbf{x}} + \hat{b}_{\mathbf{p}} e^{-i\mathbf{p}\mathbf{x}}). \quad (11.2)$$

As a result, every operator $\hat{a}_{\mathbf{p}}$ or $\hat{b}_{\mathbf{p}}$ appears multiplied by a function carrying the “correct” time factor ($\sim e^{-i\varepsilon t}$), whereas each $\hat{a}_{\mathbf{p}}^+$ or $\hat{b}_{\mathbf{p}}^+$ is accompanied by the complex conjugate of that function. This permits the identification, following the general rules, of $\hat{a}_{\mathbf{p}}$ and $\hat{b}_{\mathbf{p}}$ as annihilation operators and of $\hat{a}_{\mathbf{p}}^+$ and $\hat{b}_{\mathbf{p}}^+$ as creation operators for particles possessing momenta $\mathbf{p}$ and energies ε .

We thus arrive at the picture of particles of two kinds, appearing together and on an equal footing. They are referred to as *particles* and *antiparticles* (the reason for this name will become clear below). One kind corresponds, within the second-quantization formalism, to the operators $\hat{a}_{\mathbf{p}}$, $\hat{a}_{\mathbf{p}}^+$, and the other to $\hat{b}_{\mathbf{p}}$, $\hat{b}_{\mathbf{p}}^+$. Both species of particles, whose operators enter the same ψ -operator, thereby possess identical masses.

These conclusions can also be reached by starting directly from the demands of relativistic invariance.

From a mathematical viewpoint, Lorentz transformations are rotations of the four-dimensional coordinate system that change the orientation of the time axis (combined with purely spatial rotations, which leave the time axis unchanged, they constitute the transformation group known as the *Lorentz group*⁷). All these transformations share the property that they do not carry the t axis out of the corresponding half of the light cone, which expresses the physical principle of the existence of a limiting signal-propagation speed.

From a purely mathematical standpoint, however, a simultaneous reversal of the sign of all four coordinates is likewise a rotation (*four-dimensional inversion*): the determinant of this transformation equals +1, just as for every other rotational transformation. Under it the time axis passes from one half of the light cone to the other. Although this circumstance signifies the physical unrealizability of such a transformation (as a change of reference frame), mathematically the distinction reduces merely to the fact that (because of the pseudo-Euclidean character of the metric) such a rotation cannot be carried out without admitting a complex transformation of the coordinates in the process.

It is natural to expect that this distinction should be immaterial when one speaks of four-dimensional invariance. Then every expression that is invariant under Lorentz

⁷Note that the totality of all three-dimensional (spatial) rotations constitutes by itself a group, entering the Lorentz group as a subgroup. The totality of Lorentz transformations does not by itself form a group: the result of successive Lorentz transformations may reduce to a purely spatial rotation.

transformations should also be invariant under 4-inversion. The precise formulation of this requirement as applied to the scalar ψ -operator will be given in § 13. But let us immediately note that in any case it entails the simultaneous presence in the ψ -operators of terms with both signs in front of ε in the exponents, since the replacement $t \rightarrow -t$ reverses precisely this sign.

Returning to expressions (11.2), let us now determine the commutation relations among the operators $\hat{a}_{\mathbf{p}}, \hat{a}_{\mathbf{p}}^+$ (and $\hat{b}_{\mathbf{p}}, \hat{b}_{\mathbf{p}}^+$). In the photon case this was done (for $\hat{c}_{\mathbf{p}}, \hat{c}_{\mathbf{p}}^+$) by drawing on the oscillator analogy, that is, ultimately from the classical-limit behavior of the electromagnetic field. Here no comparable analogy exists. In order to decide whether the operators obey Bose or Fermi commutation rules, one can appeal only to the structure of the Hamiltonian built from these operators.

The Hamiltonian is obtained (see III, § 64) by substituting $\hat{\psi}$ and $\hat{\psi}^+$ for ψ and ψ^* in the integral $\int T_{00} d^3x$ ⁸. In this way we find

$$\hat{H} = \sum_{\mathbf{p}} \varepsilon(\hat{a}_{\mathbf{p}}^+ \hat{a}_{\mathbf{p}} + \hat{b}_{\mathbf{p}} \hat{b}_{\mathbf{p}}^+). \quad (11.3)$$

It is easy to see that a sensible result for the eigenvalues of this Hamiltonian is obtained only if the operators satisfy Bose commutation rules:

$$\{\hat{a}_{\mathbf{p}}, \hat{a}_{\mathbf{p}}^+\}_- = \{\hat{b}_{\mathbf{p}}, \hat{b}_{\mathbf{p}}^+\}_- \quad (11.4)$$

(all other pairs of operators commute; in particular, all particle operators $\hat{a}_{\mathbf{p}}, \hat{a}_{\mathbf{p}}^+$ commute with all antiparticle operators $\hat{b}_{\mathbf{p}}, \hat{b}_{\mathbf{p}}^+$). Indeed, in that case

$$\hat{H} = \sum_{\mathbf{p}} \varepsilon(\hat{a}_{\mathbf{p}}^+ \hat{a}_{\mathbf{p}} + \hat{b}_{\mathbf{p}} \hat{b}_{\mathbf{p}}^+ + 1).$$

The eigenvalues of the products $\hat{a}_{\mathbf{p}}^+ \hat{a}_{\mathbf{p}}$ and $\hat{b}_{\mathbf{p}} \hat{b}_{\mathbf{p}}^+$ are non-negative integers $N_{\mathbf{p}}$ and $\bar{N}_{\mathbf{p}}$ —the numbers of particles and antiparticles. The infinite additive constant $\sum \varepsilon$ (“vacuum energy”) can once again simply be dropped:

$$E = \sum_{\mathbf{p}} \varepsilon(N_{\mathbf{p}} + \bar{N}_{\mathbf{p}}) \quad (11.5)$$

(cf. formula (3.1) and the remark following it). This expression is manifestly positive and accords with the picture of two species of actually existing particles. In an analogous fashion, for the total momentum of the particle system we obtain

$$\mathbf{P} = \sum_{\mathbf{p}} \mathbf{p}(N_{\mathbf{p}} + \bar{N}_{\mathbf{p}}). \quad (11.6)$$

Had we adopted, in place of (11.4), the Fermi commutation relations (anticommutators instead of commutators), we would have obtained

$$\hat{H} = \sum_{\mathbf{p}} \varepsilon(\hat{a}_{\mathbf{p}}^+ \hat{a}_{\mathbf{p}} - \hat{b}_{\mathbf{p}}^+ \hat{b}_{\mathbf{p}} + 1)$$

⁸In non-relativistic theory one customarily places the conjugate operator $\hat{\psi}^+$ to the left of ψ . Here, however, the ordering is immaterial, since interchanging $\hat{\psi}^+$ and $\hat{\psi}$ would merely interchange the equivalent operators $\hat{a}_{\mathbf{p}}$ and $\hat{b}_{\mathbf{p}}$. It is, however, necessary, once a particular ordering has been chosen, always to adhere to the same convention.

and, instead of formula (11.5), the physically meaningless expression $\sum \varepsilon(N_{\mathbf{p}} - \bar{N}_{\mathbf{p}})$. This expression is not positive-definite and therefore cannot represent the energy of a system of free particles.

Thus spin-0 particles are bosons.

Next, consider the integral Q (10.19). Replacing the functions ψ and ψ^* in j^0 by the operators $\hat{\psi}$ and $\hat{\psi}^+$ and carrying out the integration, we obtain

$$\hat{Q} = \sum_{\mathbf{p}} (\hat{a}_{\mathbf{p}}^+ \hat{a}_{\mathbf{p}} - \hat{b}_{\mathbf{p}} \hat{b}_{\mathbf{p}}^+) = \sum_{\mathbf{p}} (\hat{a}_{\mathbf{p}}^+ \hat{a}_{\mathbf{p}} - \hat{b}_{\mathbf{p}}^+ \hat{b}_{\mathbf{p}} - 1). \quad (11.7)$$

The eigenvalues of this operator (apart from the inessential additive constant $\sum 1$) are

$$Q = \sum_{\mathbf{p}} (N_{\mathbf{p}} - \bar{N}_{\mathbf{p}}), \quad (11.8)$$

i.e. they equal the difference between the total numbers of particles and antiparticles.

As long as we deal with free particles, abstracting from all interaction among them, the content of the conservation law for Q (as well as the conservation of total energy and momentum (11.5, 11.6)) remains, of course, largely formal: what is actually conserved is not merely this sum, but each of the individual numbers $N_{\mathbf{p}}$, $\bar{N}_{\mathbf{p}}$ separately. Whether the quantity Q will remain conserved as a result of the interaction depends on the nature of that interaction. If Q is conserved (i.e. if the operator $\hat{Q}$ commutes with the interaction Hamiltonian), then expression (11.8) shows what restriction this imposes on the possible changes in particle number: only pairs “particle + antiparticle” can appear or disappear.

If a particle carries an electric charge, its antiparticle must bear a charge of the opposite sign: were both to carry identical charges, the creation or annihilation of a pair would violate the strict law of nature—conservation of the total electric charge. We shall see below (§ 32) how this opposition of charges arises automatically within the theory (in the interaction of particles with the electromagnetic field).

The quantity Q is sometimes called the *charge* of the field of the given particles. For electrically charged particles, Q determines, in particular, the total electric charge of the system (in units of the elementary charge e). Let us stress, however, that particles and antiparticles may well be electrically neutral.

We see in this way how the character of the relativistic energy–momentum relation (the two-valuedness of the root of the equation $\varepsilon^2 = \mathbf{p}^2 + m^2$), combined with the requirements of relativistic invariance, leads in quantum theory to the emergence of a new classification principle for particles—the possibility of existence of pairs of distinct particles (“particle + antiparticle”), standing in the correspondence described above. This remarkable prediction was first made (for spin-1/2 particles) by *Dirac* in 1930, even before the actual discovery of the first antiparticle—the positron⁹.

§ 12. Strictly neutral particles

When carrying out the second quantization of the ψ -function (11.1), the coefficients $a_{\mathbf{p}}^{(+)}$ and $a_{\mathbf{p}}^{(-)}$ were treated as operators pertaining to different particles. This is, however,

⁹The notion of antiparticles was extended to bosons by *Weisskopf* and *Pauli* (V. Weisskopf, W. Pauli, 1934).

not obligatory: as a special case, the annihilation and creation operators appearing in $\hat{\psi}$ may refer to one and the same type of particle (as was the case for photons—cf. (2.17)). Denoting these operators $\hat{c}_{\mathbf{p}}$ and $\hat{c}_{\mathbf{p}}^+$ in this case, we write the ψ -operator as

$$\hat{\psi} = \sum_{\mathbf{p}} \frac{1}{\sqrt{2\varepsilon}} (\hat{c}_{\mathbf{p}} e^{-ipx} + \hat{c}_{\mathbf{p}}^+ e^{ipx}). \quad (12.1)$$

The field described by such an operator corresponds to a system of identical particles that can be said to “coincide with their own antiparticles.”

The operator (12.1) is Hermitian ($\hat{\psi}^+ = \hat{\psi}$); in this sense such a field has half as many “degrees of freedom” as a complex field, for which the operators $\hat{\psi}$ and $\hat{\psi}^+$ do not coincide.

In connection with this, the Lagrangian of the field, when expressed through the Hermitian operator $\hat{\psi}$, must contain an extra factor of 1/2 (compared with (10.9))¹⁰

$$\hat{L} = (1/2)(\partial_{\mu}\hat{\psi} \cdot \partial^{\mu}\hat{\psi} - m^2\hat{\psi}^2). \quad (12.2)$$

The corresponding energy–momentum tensor is

$$\hat{T}_{\mu\nu} = \partial_{\mu}\hat{\psi} \cdot \partial_{\nu}\hat{\psi} - \hat{L}g_{\mu\nu}, \quad (12.3)$$

so that the energy-density operator reads

$$\hat{T}_{00} = \left(\frac{\partial\hat{\psi}}{\partial t}\right)^2 - \hat{L} = \frac{1}{2} \left[\left(\frac{\partial\hat{\psi}}{\partial t}\right)^2 + (\nabla\hat{\psi})^2 + m^2\hat{\psi}^2 \right]. \quad (12.4)$$

Inserting (12.1) into the integral $\int \hat{T}_{00} d^3x$, we obtain the field Hamiltonian

$$\hat{H} = \frac{1}{2} \sum_{\mathbf{p}} \varepsilon (\hat{c}_{\mathbf{p}}^+ \hat{c}_{\mathbf{p}} + \hat{c}_{\mathbf{p}} \hat{c}_{\mathbf{p}}^+). \quad (12.5)$$

This again reveals the necessity of Bose quantization:

$$\{\hat{c}_{\mathbf{p}}, \hat{c}_{\mathbf{p}}^+\}_{-} = 1, \quad (12.6)$$

and the energy eigenvalues (once more dropping the additive constant) are

$$E = \sum_{\mathbf{p}} \varepsilon_{\mathbf{p}} N_{\mathbf{p}}. \quad (12.7)$$

Fermi quantization, on the other hand, would yield a meaningless result—a value of E independent of $N_{\mathbf{p}}$.

The “charge” Q of the field under consideration vanishes. This is already clear from the fact that the charge Q must change sign when particles are replaced by antiparticles,

¹⁰This is analogous to the extra factor 1/2 in the operator (2.10) for the electromagnetic field energy density (expressed via the Hermitian operators $\hat{\mathbf{E}}$ and $\hat{\mathbf{H}}$), as compared to the photon energy density (3.2) expressed through the photon’s complex wave function; cf. the remark on p. 26.

and here the two coincide. Correspondingly, no 4-vector current density exists either. Indeed, the expression

$$\hat{j}_\mu = i[\hat{\psi}^+ \partial_\mu \hat{\psi} - (\partial_\mu \hat{\psi}^+) \hat{\psi}] \quad (12.8)$$

for the operator of the conserved 4-current $\hat{j}$ vanishes when $\hat{\psi} = \hat{\psi}^+$ (and the vector $\hat{\psi} \partial_\mu \hat{\psi}$ is not by itself conserved). This in turn signifies the absence of any special conservation law that would restrict the possible changes of the particle number. It is evident that such particles are, in any case, electrically neutral.

Particles of this kind are called *strictly neutral*, as distinct from electrically neutral particles that possess an antiparticle. While the latter can annihilate (converting into photons) only in pairs, strictly neutral particles can annihilate singly.

The structure of the ψ -operator (12.1) is the same as that of the operators (2.17)–(2.20) of the electromagnetic field. In this sense one can say that photons themselves are strictly neutral particles. In the case of the electromagnetic field, the Hermiticity of the operators was linked to the reality of the field strengths as measurable (in the classical limit) physical quantities. For the ψ -operators of particles, however, no such link exists, since they do not correspond to any directly measurable quantities at all.

The absence of a conserved 4-vector current is a general property of strictly neutral particles and is not connected with zero spin (it holds for photons as well). Physically it expresses the absence of the corresponding prohibitions on changes in particle number. From a formal standpoint there is a direct connection between the absence of a conserved current and the reality of the field—the Hermiticity of the operator $\hat{\psi}$.

The Lagrangian of a complex field

$$\hat{L} = \partial_\mu \hat{\psi}^+ \cdot \partial^\mu \hat{\psi} - m^2 \hat{\psi}^+ \hat{\psi} \quad (12.9)$$

is invariant under multiplication of the ψ -operator by an arbitrary phase factor, i.e. under transformations of the form

$$\hat{\psi} \rightarrow e^{i\alpha} \hat{\psi}, \quad \hat{\psi}^+ \rightarrow e^{-i\alpha} \hat{\psi}^+ \quad (12.10)$$

(these are termed *gauge* transformations). In particular, the Lagrangian remains unchanged under an infinitesimally small gauge transformation

$$\hat{\psi} \rightarrow \hat{\psi} + i\delta\alpha \cdot \hat{\psi}, \quad \hat{\psi}^+ \rightarrow \hat{\psi}^+ - i\delta\alpha \cdot \hat{\psi}^+. \quad (12.11)$$

Under an infinitesimally small variation of the “generalized coordinates” q , the Lagrangian undergoes the change

$$\begin{aligned} \delta\hat{L} &= \sum \left(\frac{\partial\hat{L}}{\partial q} \delta q + \frac{\partial\hat{L}}{\partial q_{,\mu}} \delta q_{,\mu} \right) = \\ &= \sum \left(\frac{\partial\hat{L}}{\partial q} - \frac{\partial}{\partial x^\mu} \frac{\partial\hat{L}}{\partial q_{,\mu}} \right) \delta q + \sum \frac{\partial}{\partial x^\mu} \left(\frac{\partial\hat{L}}{\partial q_{,\mu}} \delta q \right) \end{aligned}$$

(summation over all q). The first term vanishes by virtue of the “equations of motion” (Lagrange’s equations). Taking $\hat{\psi}$ and $\hat{\psi}^+$ as the “coordinates” q and setting

$$\delta\hat{\psi} = i\delta\alpha \cdot \hat{\psi}, \quad \delta\hat{\psi}^+ = -i\delta\alpha \cdot \hat{\psi}^+,$$

we get

$$\delta \hat{L} = i\delta\alpha \frac{\partial}{\partial x^\mu} \left(\hat{\psi} \frac{\partial \hat{L}}{\partial \hat{\psi}_{,\mu}} - \hat{\psi}^+ \frac{\partial \hat{L}}{\partial \hat{\psi}_{,\mu}^+} \right).$$

It is apparent that the condition of invariance of the Lagrangian ($\delta \hat{L} = 0$) is equivalent to the continuity equation ($\partial_\mu \hat{j}^\mu = 0$) for the 4-vector

$$\hat{j}^\mu = i \left(\hat{\psi}^+ \frac{\partial \hat{L}}{\partial \hat{\psi}_{,\mu}^+} - \hat{\psi} \frac{\partial \hat{L}}{\partial \hat{\psi}_{,\mu}} \right). \quad (12.12)$$

It is readily verified that for the Lagrangian (12.9) this formula yields the current (12.8).

We see, therefore, that in the mathematical framework of the theory the existence of a conserved current is tied to gauge invariance of the Lagrangian (W. Pauli, 1941). The Lagrangian (12.2) of a strictly neutral field, however, lacks this symmetry.

§ 13. The transformations C, P, T

Unlike four-dimensional inversion, a three-dimensional (spatial) inversion cannot be reduced to any rotation of the 4-coordinate system: the determinant of this transformation equals not +1 but -1. The symmetry properties of particles with respect to inversion (P -transformation) are therefore not dictated by the requirements of relativistic invariance¹¹.

Applied to a scalar wave function, the inversion operation amounts to the transformation

$$\hat{P}\psi(t, \mathbf{r}) = \pm\psi(t, -\mathbf{r}), \quad (13.1)$$

where the sign “+” or “-” on the right-hand side corresponds, respectively, to a true scalar or a pseudoscalar.

It is apparent from this that one must distinguish two aspects of the wave function's behavior under inversion. One of them is related to the dependence of the wave function on the coordinates. In nonrelativistic quantum mechanics only this aspect was considered—it leads to the notion of the parity of a state (which we shall henceforth call the *orbital parity*), characterizing the symmetry properties of the particle's motion. If a state possesses a definite orbital parity (+1 or -1), this means that

$$\psi(t, -\mathbf{r}) = \pm\psi(t, \mathbf{r}).$$

The second aspect is the behavior of the wave function at a given point (which one may conveniently take to be the coordinate origin) when the coordinate axes are inverted. This behavior gives rise to the concept of the *intrinsic parity of the particle*. For a spin-0 particle the two signs in definition (13.1) correspond to intrinsic-parity values of +1 and -1. The overall parity of a particle system equals the product of their individual intrinsic parities and the orbital parity associated with the relative motion.

The “intrinsic” symmetry properties of different particles manifest themselves, of course, only in the processes of their mutual transformations. The analogue of intrinsic

¹¹The Lorentz group supplemented by spatial inversion is called the *extended Lorentz group* (as opposed to the original group, which does not contain P and which in this context is termed *proper*). The extended group includes all transformations that do not carry the t axis out of the corresponding half of the light cone.

parity in nonrelativistic quantum mechanics is the parity of a bound state of a composite system (for instance, a nucleus). From the standpoint of relativistic theory, which draws no fundamental distinction between composite and elementary particles, this kind of intrinsic parity is no different from the intrinsic parity of the particles that figure as elementary ones in nonrelativistic theory. In the nonrelativistic domain, where the latter behave as immutable objects, their intrinsic symmetry properties are unobservable, and consequently any consideration of them would be devoid of physical content.

Within the second-quantization formalism, the intrinsic parity is expressed through the behavior of the ψ -operators under inversion. For scalar and pseudoscalar fields the transformation laws read

$$P : \hat{\psi}(t, \mathbf{r}) \rightarrow \pm \hat{\psi}(t, -\mathbf{r}). \quad (13.2)$$

The very meaning of the action of inversion on the ψ -operator must, however, be formulated as a definite transformation of the particle annihilation and creation operators—one such that the change (13.2) results from it. It is easy to see that the required transformation is

$$P : \hat{a}_{\mathbf{p}} \rightarrow \pm \hat{a}_{-\mathbf{p}}, \quad \hat{b}_{\mathbf{p}} \rightarrow \pm \hat{b}_{-\mathbf{p}} \quad (13.3)$$

(and likewise for the conjugate operators). Indeed, carrying out this substitution in the operator

$$\hat{\psi}(t, \mathbf{r}) = \sum_{\mathbf{p}} \frac{1}{\sqrt{2\varepsilon}} \left(\hat{a}_{\mathbf{p}} e^{-i\omega t + i\mathbf{p}\mathbf{r}} + \hat{b}_{\mathbf{p}}^{\dagger} e^{i\omega t - i\mathbf{p}\mathbf{r}} \right) \quad (13.4)$$

and then relabeling the summation variable ($\mathbf{p} \rightarrow -\mathbf{p}$), we bring it to the form $\pm \hat{\psi}(t, -\mathbf{r})$. Thus, denoting by $\hat{\psi}^P(t, \mathbf{r})$ the operator in which transformation (13.3) has been performed, one may write

$$\hat{\psi}^P(t, \mathbf{r}) = \pm \hat{\psi}(t, -\mathbf{r}). \quad (13.5)$$

Note that transformation (13.3) has a perfectly natural form: inversion reverses the sign of the polar vector $\mathbf{p}$, so that particles with momentum $\mathbf{p}$ are replaced by particles with momentum $-\mathbf{p}$.

In (13.3) the operators $\hat{a}_{\mathbf{p}}$ and $\hat{b}_{\mathbf{p}}$ both transform with either the upper or the lower signs. In the second-quantization formalism this expresses the equality of the intrinsic parities of particle and antiparticle (with spin 0). This equality is in itself already obvious from the fact that particles and antiparticles (of spin 0) are described by the very same (scalar or pseudoscalar) wave functions.

In relativistic theory there also arises a symmetry with respect to a transformation that has no counterpart in nonrelativistic theory; it is called *charge conjugation* (C -transformation). If all operators $\hat{a}_{\mathbf{p}}$ and $\hat{b}_{\mathbf{p}}$ are mutually interchanged:

$$C : \hat{a}_{\mathbf{p}} \rightarrow \hat{b}_{\mathbf{p}}, \quad \hat{b}_{\mathbf{p}} \rightarrow \hat{a}_{\mathbf{p}} \quad (13.6)$$

(i.e. particles and antiparticles are interchanged), then $\hat{\psi}$ goes over into the “charge-conjugate” operator $\hat{\psi}^C$, with

$$\hat{\psi}^C(t, \mathbf{r}) = \hat{\psi}^{\dagger}(t, \mathbf{r}) \quad (13.7)$$

This equality expresses the symmetry with which the notions of particles and antiparticles enter the theory.

Let us note that the definition of the charge-conjugation transformation contains a certain inessential formal arbitrariness. The content of the transformation is unaltered if one introduces into definition (13.6) an arbitrary phase factor:

$$\hat{a}_{\mathbf{p}} \rightarrow e^{i\alpha} \hat{b}_{\mathbf{p}}, \quad \hat{b}_{\mathbf{p}} \rightarrow e^{-i\alpha} \hat{a}_{\mathbf{p}}.$$

Then one would have

$$\hat{\psi} \rightarrow e^{i\alpha} \hat{\psi}^+, \quad \hat{\psi}^+ \rightarrow e^{-i\alpha} \hat{\psi},$$

and a twofold application of this transformation would still yield the identity ($\hat{\psi} \rightarrow \hat{\psi}$). All such definitions are, however, equivalent to one another. Since the properties of the ψ -operators are unaffected by multiplication by a phase factor (cf. the end of the preceding section), one may simply relabel $\hat{\psi}$ as $\hat{\psi} e^{i\alpha/2}$ and then return to the definition of charge conjugation in the form (13.6), (13.7).

Because charge conjugation replaces a particle by a non-identical antiparticle, it does not in the general case give rise to any new characteristic of a particle or a system of particles as such.

Systems containing equal numbers of particles and antiparticles constitute an exception. The operator $\hat{C}$ carries such a system into itself, so that eigenstates exist with eigenvalues $C = \pm 1$ (which follow from $\hat{C}^2 = 1$). When describing charge symmetry one may treat particle and antiparticle as two distinct “charge states” of a single particle, distinguished by the charge quantum number $Q = \pm 1$. The system’s wave function then factorizes into an orbital part and a “charge” part, and must be symmetric with respect to the simultaneous interchange of all variables (spatial and charge) belonging to any pair of particles. The symmetry character of the “charge” factor fixes the charge parity of the system (see the Problem)¹².

The concept of charge parity, which arises naturally for “strictly neutral” systems, should apply also to strictly neutral “elementary” particles. In the second-quantization formalism this concept is described by the relation

$$\hat{\psi}^C = \pm \hat{\psi}; \quad (13.8)$$

the signs “+” and “−” correspond to charge-even and charge-odd particles.

In § 11 it was pointed out that relativistic invariance must also entail invariance under 4-inversion. For the operator of a scalar (in the sense of 4-rotations) field this means that under such a transformation one must have

$$\hat{\psi}(t, \mathbf{r}) = \hat{\psi}(-t, -\mathbf{r})$$

always with the same sign “+” on the right-hand side. In terms of the operators $\hat{a}_{\mathbf{p}}, \hat{b}_{\mathbf{p}}$, the conversion of $\hat{\psi}(t, \mathbf{r})$ into $\hat{\psi}(-t, -\mathbf{r})$ is accomplished by interchanging the coefficients of e^{-ipx} and e^{ipx} in (13.4), i.e. by the replacement

$$\hat{a}_{\mathbf{p}} \rightarrow \hat{b}_{\mathbf{p}}^+, \quad \hat{b}_{\mathbf{p}} \rightarrow \hat{a}_{\mathbf{p}}^+ \quad (13.9)$$

¹²In these arguments we have in mind particles of spin 0. The method of analysis described here generalizes directly to other cases—see, for example, the Problem in § 27.

Since it replaces the a -operators by b -operators, this transformation incorporates the mutual interchange of particles and antiparticles. We see that in relativistic theory there naturally arises a requirement of invariance under a transformation in which spatial inversion (P) and time reversal (T) are accompanied by charge conjugation (C); this statement is known as the *CPT theorem*¹³.

In this connection it is, however, appropriate to stress that although the arguments presented here and in §§ 11, 12 appear to be a natural development of the concepts of ordinary quantum mechanics and classical relativity, the results obtained in this way go beyond that framework both in form (ψ -operators that simultaneously contain creation and annihilation operators for particles) and in substance (particles and antiparticles). These results cannot therefore be regarded as a matter of pure logical necessity. They embody new physical principles whose correctness can be judged only by experiment.

If we denote by $\hat{\psi}^{CPT}(t, \mathbf{r})$ the operator (13.4) in which the transformation (13.9) has been carried out, one may write

$$\hat{\psi}^{CPT}(t, \mathbf{r}) = \hat{\psi}(-t, -\mathbf{r}). \quad (13.10)$$

Having thus formulated 4-inversion as the transformation (13.9), we thereby establish for the ψ -operator also the formulation of the time-reversal transformation: together with the CP transformation (called the *combined inversion*) it must yield (13.9). Taking into account the definitions (13.3) and (13.6), we therefore find

$$T : \hat{a}_{\mathbf{p}} \rightarrow \pm \hat{a}_{-\mathbf{p}}^+, \quad \hat{b}_{\mathbf{p}} \rightarrow \pm \hat{b}_{-\mathbf{p}}^+ \quad (13.11)$$

(the signs “ $\pm$ ” match those in (13.3)). The meaning of this transformation is perfectly natural: time reversal not only converts motion with momentum $\mathbf{p}$ into motion with momentum $-\mathbf{p}$, but also interchanges the initial and final states in matrix elements; the annihilation operators for particles with momenta $\mathbf{p}$ are therefore replaced by creation operators for particles with momenta $-\mathbf{p}$. Performing the substitution (13.11) in (13.4) and relabeling the summation variable ($\mathbf{p} \rightarrow -\mathbf{p}$), we find that¹⁴

$$\hat{\psi}^T(t, \mathbf{r}) = \pm \hat{\psi}^+(-t, \mathbf{r}). \quad (13.12)$$

This result mirrors the standard time-reversal prescription of quantum mechanics: when a state has wave function $\psi(t, \mathbf{r})$, the “temporally reversed” state is represented by $\psi^*(-t, \mathbf{r})$; complex conjugation is needed in order to recover the “proper” character of the time dependence, which the sign flip of t would otherwise spoil (*E. P. Wigner*, 1932).

Since the transformation T (and with it CPT) interchange initial and final states, the concepts of eigenstates and eigenvalues have no meaning for them. They therefore do not lead to new characteristics of particles as such. The consequences to which they lead when applied to scattering processes will be discussed in §§ 69, 71.

Let us examine how the operator 4-vector of the current $\hat{j}^\mu$ (12.8) transforms under C , P , and T . Transformation (13.2) together with the replacement $(\partial_0, \partial_i) \rightarrow (\partial_0, -\partial_i)$

¹³It was formulated by *Lüders* (G. Lüders, 1954) and *Pauli* (W. Pauli, 1955).

¹⁴If one were to define the operation T independently of the other transformations, the same arbitrariness in the choice of a phase factor would arise as for the operation C . The requirement of CPT symmetry, however, leaves the phase-factor choice free in only one of the two transformations, C or T .

gives

$$P : (\hat{j}^0, \hat{\mathbf{j}})_{t,\mathbf{r}} \rightarrow (\hat{j}^0, -\hat{\mathbf{j}})_{t,-\mathbf{r}}, \quad (13.13)$$

as it must for a true 4-vector. Transformation (13.7) would simply give

$$C : (\hat{j}^0, \hat{\mathbf{j}})_{t,\mathbf{r}} \rightarrow (-\hat{j}^0, -\hat{\mathbf{j}})_{t,\mathbf{r}}, \quad (13.14)$$

were the operators $\hat{\psi}$ and $\hat{\psi}^+$ commutative. The noncommutativity of these operators arises, however, solely from the noncommutativity of $\hat{a}_{\mathbf{p}}$ and $\hat{a}_{\mathbf{p}}^+$ (or of $\hat{b}_{\mathbf{p}}$ and $\hat{b}_{\mathbf{p}}^+$) at the same $\mathbf{p}$; but by virtue of the commutation rules (11.4) the interchange of these operators merely introduces terms independent of the occupation numbers, i.e. of the field state. Discarding these terms as inessential (just as in (11.5),(11.6)), we recover rule (13.14), which has a natural meaning: by replacing particles with antiparticles, charge conjugation reverses the sign of every component of the 4-current.

Since the time-reversal operation is linked with the transposition of initial and final states, its application to a product of operators reverses the order of the factors. Thus,

$$(\hat{\psi}^+ \partial_\mu \hat{\psi})^T = (\partial_\mu \hat{\psi})^T (\hat{\psi}^+)^T.$$

In the present case, however, this is immaterial: by virtue of the commutativity of the ψ -operators (in the sense indicated above), reverting to the original order of the factors does not affect the result. Noting also that under time reversal $(\partial_0, \partial_i) \rightarrow (-\partial_0, \partial_i)$, we find the transformation rule for the current:

$$T : (\hat{j}^0, \hat{\mathbf{j}})_{t,\mathbf{r}} \rightarrow (\hat{j}^0, -\hat{\mathbf{j}})_{-t,\mathbf{r}}. \quad (13.15)$$

The three-dimensional vector $\mathbf{j}$ changes sign in accordance with the classical meaning of this quantity.

Finally, under the CPT transformation we have

$$CPT : (\hat{j}^0, \hat{\mathbf{j}})_{t,\mathbf{r}} \rightarrow (-\hat{j}^0, -\hat{\mathbf{j}})_{-t,-\mathbf{r}}, \quad (13.16)$$

in accord with the meaning of this operation as a 4-inversion. Let us stress in this connection that, since 4-inversion reduces to a rotation of the 4-coordinate system, with respect to it there is no distinction at all between two types (true and pseudo) of 4-tensors of any rank.

Up to now we have been considering free particles. But the quantum numbers of parity acquire real significance only when one examines interacting particles, where definite selection rules are associated with them, permitting or forbidding particular processes. Such significance, however, can attach only to conserved quantities—eigenvalues of operators that commute with the Hamiltonian of the interacting particles.

Relativistic invariance guarantees that the CPT -transformation operator commutes with the Hamiltonian in every case. As for the individual transformations C and P (together with T), experiment demonstrates that electromagnetic and strong interactions respect these symmetries, so that the associated parity quantum numbers are conserved within those interactions. In weak processes, however, these conservation laws break down ¹⁵.

¹⁵The suggestion that parity conservation might fail in weak interactions was first put forward by Lee and Yang (*T. D. Lee, C. N. Yang*, 1956). Earlier still, the broader idea that P - and T -invariance of physical laws is not mandatory was advanced by Dirac (1949).

Anticipating later developments, we point out that the operator governing the coupling of charged particles to the electromagnetic field is formed as the product of the operator 4-vectors $\hat{A}$ and $\hat{j}$. Because charge conjugation flips the sign of $\hat{j}$, electromagnetic-interaction invariance under this transformation requires that $\hat{A}$ likewise change sign. Put differently, photons are charge-odd particles.

The stated behavior of the operators $\hat{A}$ is in agreement with the properties of the 4-potential in classical theory. Indeed, from the transformations

$$C : (\hat{A}_0, \hat{\mathbf{A}}) \rightarrow (-\hat{A}_0, -\hat{\mathbf{A}})_{t, \mathbf{r}},$$

$$P : (\hat{A}_0, \hat{\mathbf{A}}) \rightarrow (\hat{A}_0, -\hat{\mathbf{A}})_{t, -\mathbf{r}},$$

$$CPT : (\hat{A}_0, \hat{\mathbf{A}}) \rightarrow (-\hat{A}_0, -\hat{\mathbf{A}})_{-t, -\mathbf{r}},$$

it follows that

$$T : (\hat{A}_0, \hat{\mathbf{A}}) \rightarrow (\hat{A}_0, -\hat{\mathbf{A}})_{-t, \mathbf{r}},$$

which corresponds to the classical rule for transforming the electromagnetic-field potentials under time reversal.

The requirement of *CPT* invariance does not impose any restrictions on the properties of particles in themselves. It does, however, lead to a definite connection between the properties of particles and antiparticles. This includes, first and foremost, the equality of their masses—a fact already clear from the connection, set out in § 11, between 4-inversion and the very origin of the particle–antiparticle concept.

Furthermore, *CPT* invariance implies that the proportionality coefficients relating the electric and magnetic moment vectors to the spin vector differ for particle and antiparticle only in sign. Indeed, the magnetic moment changes sign under *C* and *T* transformations and (being an axial vector) is *P*-invariant. Consequently, the *CPT* transformation, while converting a particle into its antiparticle, does not at the same time change the sign of the magnetic moment; the spin vector, however, does change sign. The same applies to the electric moment, which remains unchanged under time reversal and changes sign under *C*-transformation and (by the properties of a polar vector) under spatial inversion.

The requirements of *P*- or *T*-invariance (if they are obeyed), on the other hand, constrain the properties of each individual particle: they forbid the particle from possessing an electric dipole moment. Indeed, the only vector that can be constructed for an elementary particle at rest from its ψ -operators is the vector of its spin operator. This vector is *P*-even and *T*-odd; it can therefore determine only the magnetic, but not the electric moment. Let us emphasize that for this prohibition it suffices to require merely one of the two—*P*- or *T*-invariance.

Problem

Determine the charge parity and spatial parity of a system consisting of a spin-0 particle and its antiparticle, with orbital angular momentum l of the relative motion.

Solution. Permutation of the particle coordinates is equivalent to inversion (about the center of inertia) and therefore multiplies the orbital function by $(-1)^l$; permutation of

the charge variables is equivalent to charge conjugation and multiplies the “charge” factor in the wave function by the sought quantity C . From the condition $C(-1)^l = 1$ we have

$$C = (-1)^l.$$

The spatial parity P of the system is the product of the orbital parity with the intrinsic parities of the two particles. Because particle and antiparticle share the same intrinsic parity, here P simply equals the orbital parity: $P = (-1)^l$.

§ 14. Wave equation for spin-1 particles

In its rest frame a particle of spin 1 is characterized by a wave function with three components—a three-dimensional vector (one therefore often speaks of a *vector* particle). Viewed four-dimensionally, these three components can be either the spatial part of a (spacelike) 4-vector ψ^μ , or the mixed components of an antisymmetric 4-tensor of rank two, $\psi^{\mu\nu}$, for which the time (ψ^0) and purely spatial (ψ^{ik}) components vanish in the rest frame¹⁶.

The wave equation—a differential relation between the quantities ψ^μ and $\psi^{\mu\nu}$ —is established by the relations that we write as

$$i\psi_{\mu\nu} = \hat{p}_\mu\psi_\nu - \hat{p}_\nu\psi_\mu, \quad (14.1)$$

$$im^2\psi_\mu = \hat{p}^\nu\psi_{\mu\nu}, \quad (14.2)$$

where $\hat{p} = i\partial$ (A. Proca, 1936). Applying the operator $\hat{p}^\mu$ to both sides of equation (14.2), we obtain (on account of the antisymmetry of $\psi_{\mu\nu}$)

$$\hat{p}^\mu\psi_\mu = 0. \quad (14.3)$$

From (14.1, 14.2) one can eliminate $\psi_{\mu\nu}$ by inserting the first equation into the second. With the aid of (14.3) this yields

$$(\hat{p}^2 - m^2)\psi_\mu = 0, \quad (14.4)$$

from which it is again apparent (cf. § 10) that m is the particle mass. A free spin-1 particle can therefore be described entirely by a single 4-vector ψ^μ whose components satisfy the second-order equation (14.4) together with the supplementary condition (14.3), which removes from ψ^μ the part belonging to spin 0.

In the rest frame, where ψ_μ is independent of the spatial coordinates, we find that $\hat{p}^0\psi_0 = 0$. Since at the same time $\hat{p}^0\psi_0 = m\psi_0$, it follows that $\psi_0 = 0$ in the rest frame, as required. Together with ψ_0 the quantities ψ_{ik} also vanish.

A spin-1 particle can possess different intrinsic parities, depending on whether ψ^μ is a true vector or a pseudovector. In the first case

$$\hat{P}\psi^\mu = (\psi^0, -\psi^i),$$

¹⁶Looking ahead, let us mention that the set consisting of the 4-vector ψ_μ and the 4-tensor $\psi^{\nu\mu}$ corresponds to the set of four-dimensional second-rank spinors $\xi^{\alpha\beta}$, $\eta_{\dot{\alpha}\dot{\beta}}$, $\xi^{\alpha\dot{\beta}}$, where $\xi^{\alpha\beta}$ and $\eta_{\dot{\alpha}\dot{\beta}}$ are symmetric spinors that transform into each other under inversion (see § 19).

and in the second

$$\hat{P}\psi^\mu = \left(-\psi^0, \psi^i\right).$$

Equations (14.1),(14.2) can be derived from a variational principle with the Lagrangian

$$L = (1/2)\psi_{\mu\nu}\psi^{\mu\nu*} - (1/2)\psi^{\mu\nu*}(\partial_\mu\psi_\nu - \partial_\nu\psi_\mu) - \\ - (1/2)\psi^{\mu\nu}(\partial_\mu\psi_\nu^* - \partial_\nu\psi_\mu^*) + m^2\psi_\mu\psi^{\mu*}. \quad (14.5)$$

The independent generalized coordinates here are $\psi_\mu, \psi_\mu^*, \psi_{\mu\nu}, \psi_{\mu\nu}^*$ ¹⁷.

For obtaining the energy-momentum tensor, formula (10.11) is not entirely convenient in the present case, since it would lead to a non-symmetric tensor requiring additional symmetrization. One may instead use the formula

$$\frac{1}{2}T_{\mu\nu}\sqrt{-g} = -\frac{\partial}{\partial x^\lambda}\frac{\partial\sqrt{-g}L}{\partial g_{,\lambda}^{\mu\nu}} + \frac{\partial\sqrt{-g}L}{\partial g^{\mu\nu}}, \quad (14.6)$$

in which it is assumed that L is written in a form valid for arbitrary curvilinear coordinates (see II, § 94). If L involves only the components of the metric tensor $g_{\mu\nu}$ itself (but not their coordinate derivatives), the formula simplifies to

$$T_{\mu\nu} = \frac{2}{\sqrt{-g}}\frac{\partial\sqrt{-g}L}{\partial g^{\mu\nu}} = 2\frac{\partial L}{\partial g^{\mu\nu}} - g_{\mu\nu}L$$

(recall that $d\ln g = -g_{\mu\nu}dg^{\mu\nu}$).

Since the differentiation in formula (14.6) is not performed with respect to ψ_μ or $\psi_{\mu\nu}$, when applying it there is no need to treat these quantities as independent; one may directly use relation (14.1) and rewrite the Lagrangian (14.5) as

$$L = (-1/2)\psi_{\mu\nu}\psi_{\lambda\rho}^*g^{\mu\lambda}g^{\nu\rho} + m^2\psi_\mu\psi_\nu^*g^{\mu\nu}. \quad (14.7)$$

Then

$$T_{\mu\nu} = -\psi_{\mu\lambda}\psi_\nu^{\lambda*} - \psi_{\mu\lambda}^*\psi_\nu^\lambda + m^2\left(\psi_\mu^*\psi_\nu + \psi_\nu^*\psi_\mu\right) + \\ + g_{\mu\nu}\left((1/2)\psi_{\lambda\rho}\psi^{\lambda\rho*} - m^2\psi_\lambda^*\psi^\lambda\right). \quad (14.8)$$

In particular, the energy density is given by a manifestly positive expression

$$T_{00} = (1/2)\psi_{ik}\psi_{ik}^* + \psi_{0i}\psi_{0i}^* + m^2(\psi_0\psi_0^* + \psi_i\psi_i^*). \quad (14.9)$$

The conserved 4-vector of the current density reads

$$j^\mu = i(\psi^{\mu\nu*}\psi_\nu - \psi^{\mu\nu}\psi_\nu^*). \quad (14.10)$$

¹⁷Had we carried out the variation with respect to ψ_μ alone (assuming from the outset that $\psi_{\mu\nu}$ is expressed through ψ_μ via (14.1)), equation (14.3) would have to be imposed as an auxiliary condition unrelated to the variational principle.

It can be obtained from formula (12.12) by differentiating the Lagrangian (14.5) with respect to the derivatives $\partial_\mu \psi_\nu$. In particular,

$$j^0 = i \left(\psi^{0k*} \psi_k - \psi^{0k} \psi_k^* \right) \quad (14.11)$$

and is not a positive-definite quantity.

A plane wave normalized to one particle in volume $V = 1$:

$$\psi_\mu = \frac{1}{\sqrt{2\varepsilon}} u_\mu e^{-ipx}, \quad u_\mu u^{\mu*} = -1, \quad (14.12)$$

where u_μ is a unit polarization 4-vector obeying (by virtue of (14.3)) the four-dimensional transversality condition

$$u_\mu p^\mu = 0. \quad (14.13)$$

Indeed, substituting the function (14.12) into (14.9) and (14.11) gives

$$T_{00} = -2\varepsilon^2 \psi_\mu \psi^{\mu*} = \varepsilon, \quad j^0 = 1.$$

In contrast to the photon, a vector particle of nonzero mass possesses three independent polarization directions. For the corresponding amplitudes see (16.21).

For partially polarized vector particles the polarization density matrix is introduced in such a way that for a pure state it becomes simply the product

$$\rho_{\mu\nu} = u_\mu u_\nu^*$$

(analogously to expression (8.7) for photons). According to (14.12), (14.13) it satisfies the conditions

$$p^\mu \rho_{\mu\nu} = 0, \quad \rho_\mu^\mu = -1. \quad (14.14)$$

For unpolarized particles the matrix $\rho_{\mu\nu}$ must have the form $a g_{\mu\nu} + b p_\mu p_\nu$. Fixing the coefficients a and b from (14.14), one finds

$$\rho_{\mu\nu} = -\frac{1}{3} \left(g_{\mu\nu} - \frac{p_\mu p_\nu}{m^2} \right). \quad (14.15)$$

Quantization of the vector-particle field proceeds in the same way as for the scalar case, and there is no need to repeat the whole argument anew. The ψ -operators of the vector field take the form

$$\begin{aligned} \hat{\psi}_\mu &= \sum_{\mathbf{p}\alpha} \frac{1}{\sqrt{2\varepsilon}} \left(\hat{a}_{\mathbf{p}\alpha} u_\mu^{(\alpha)} e^{-ipx} + \hat{b}_{\mathbf{p}\alpha}^+ u_\mu^{(\alpha)*} e^{ipx} \right), \\ \hat{\psi}_\mu^+ &= \sum_{\mathbf{p}\alpha} \frac{1}{\sqrt{2\varepsilon}} \left(\hat{a}_{\mathbf{p}\alpha}^+ u_\mu^{(\alpha)*} e^{ipx} + \hat{b}_{\mathbf{p}\alpha} u_\mu^{(\alpha)} e^{-ipx} \right), \end{aligned} \quad (14.16)$$

where the index α labels the three independent polarizations.

The positive definiteness of expression (14.9) for T_{00} and the indefiniteness of j^0 (14.11) lead, just as in the scalar case, to the requirement of Bose quantization.

There exists a close connection between the properties of the strictly neutral vector field and the electromagnetic field. A neutral vector field is described by a Hermitian ψ -operator:

$$\hat{\psi}_\mu = \sum_{\mathbf{p}\alpha} \frac{1}{\sqrt{2\varepsilon}} \left(\hat{c}_{\mathbf{p}\alpha} u_\mu^{(\alpha)} e^{-ipx} + \hat{c}_{\mathbf{p}\alpha}^+ u_\mu^{(\alpha)*} e^{ipx} \right). \quad (14.17)$$

The Lagrangian of this field is

$$\hat{L} = \frac{1}{4} \hat{\psi}_{\mu\nu} \hat{\psi}^{\mu\nu} - \frac{1}{2} \hat{\psi}^{\mu\nu} (\partial_\mu \hat{\psi}_\nu - \partial_\nu \hat{\psi}_\mu) + \frac{1}{2} m^2 \hat{\psi}_\mu \hat{\psi}^\mu. \quad (14.18)$$

The electromagnetic field corresponds to the case $m = 0$. The 4-vector ψ^μ then becomes the 4-potential A^μ , and the 4-tensor $\psi^{\mu\nu}$ becomes the field tensor $F^{\mu\nu}$, related to the potential through definition (14.1). Equation (14.2) turns into $\partial^\nu \psi_{\mu\nu} = 0$, which corresponds to the second pair of Maxwell's equations. Condition (14.3) no longer follows from it and thus ceases to be obligatory. In the absence of the supplementary condition there is no need to treat $\hat{\psi}_\mu$ and $\hat{\psi}_{\mu\nu}$ as independent “coordinates” in the Lagrangian, and (14.18) reduces to

$$\hat{L} = -(1/4) \hat{\psi}_{\mu\nu} \hat{\psi}^{\mu\nu} \quad (14.19)$$

in agreement with the familiar classical expression for the electromagnetic-field Lagrangian. This Lagrangian, together with the tensor $\hat{\psi}_{\mu\nu}$, is invariant under an arbitrary gauge transformation of the “potentials” $\hat{\psi}_\mu$. The connection between this circumstance and the vanishing mass is clearly visible: the Lagrangian (14.18) lacks this property on account of the term $m^2 \hat{\psi}_\mu \hat{\psi}^\mu$.

§ 15. Wave equation for particles with higher integer spins

Since the wave equations (14.3, 14.4) follow directly from specifying the mass and spin of the particle, the practical use of the Lagrangian comes down not so much to deriving these equations as to constructing the expressions for the energy, momentum, and charge of the field.

For this purpose, as already noted, one may use expression (14.7) in place of (14.5), and the former can be further transformed. Using (14.1) and rewriting the Lagrangian (14.7) as

$$\begin{aligned} L &= -(\partial_\mu \psi_\nu^*) (\partial^\mu \psi^\nu) + (\partial_\nu \psi_\mu^*) (\partial^\mu \psi^\nu) + m^2 \psi_\mu \psi^{\mu*} = \\ &= -(\partial_\mu \psi_\nu^*) (\partial^\mu \psi^\nu) + m^2 \psi_\mu^* \psi^\mu + \partial_\nu (\psi_\mu^* \partial^\mu \psi^\nu) - \psi_\mu^* \partial^\mu \partial_\nu \psi^\nu. \end{aligned} \quad (15.1)$$

By virtue of (14.3) the last term vanishes, while the penultimate one is a total derivative. Dropping it, we arrive at the Lagrangian

$$L' = -(\partial_\mu \psi_\nu^*) (\partial^\mu \psi^\nu) + m^2 \psi_\mu^* \psi^\mu.$$

It has the same structure as the Lagrangian (10.9) for a spin-0 particle, differing only in that the scalar ψ is replaced by the 4-vector ψ_μ and the overall sign is reversed. The sign

change is due to the spacelike character of ψ_μ , so that $\psi_\mu \psi^{\mu*} < 0$, whereas for a scalar particle $\psi \psi^* > 0$.

If one constructs the energy–momentum 4-tensor and the current 4-vector with the help of Lagrangian (15.1), one obtains expressions of the same form as expressions (10.12) and (10.18) for the scalar field:

$$T_{\mu\nu} = -\partial_\mu \psi^{\lambda*} \cdot \partial_\nu \psi_\lambda - \partial_\nu \psi^{\lambda*} \cdot \partial_\mu \psi_\lambda - L' g_{\mu\nu}, \quad (15.2)$$

$$j_\mu = -i [\psi_\lambda^* \partial_\mu \psi^\lambda - (\partial_\mu \psi_\lambda^*) \psi^\lambda]. \quad (15.3)$$

Their difference from (14.8) and (14.10) likewise reduces to total derivatives. But the local values of these quantities lack (as was already stressed) any deep physical significance. Only the volume integrals P_μ (10.15) and Q (10.19) matter, and they coincide for either choice of $T_{\mu\nu}$ and j_μ .

This method of description extends straightforwardly to particles of arbitrary (integer) spin. The wave function of a particle with spin s is an irreducible 4-tensor of rank s , that is, a tensor symmetric in all of its indices and vanishing upon contraction over any pair of them:

$$\psi_{\dots\mu\dots\nu\dots} = \psi_{\dots\nu\dots\mu\dots}, \quad \psi_{\dots\mu\dots}{}^\mu{}_{\dots} = 0. \quad (15.4)$$

This tensor must obey the supplementary four-dimensional transversality condition

$$\hat{p}^\mu \psi_{\dots\mu\dots} = 0, \quad (15.5)$$

and each of its components must satisfy the second-order equation

$$(\hat{p}^2 - m^2) \psi_{\dots} = 0. \quad (15.6)$$

In the rest frame, condition (15.5) causes every component of the 4-tensor that carries a 0 among its indices to vanish. In other words, the wave function in the rest frame (i.e. in the nonrelativistic limit) reduces, as expected, to an irreducible 3-tensor of rank s with $2s + 1$ independent components.

The Lagrangian, the energy–momentum tensor, and the current vector for particles of spin s differ from (15.1)–(15.3) solely by the replacement of ψ_λ with $\psi_{\lambda\mu\dots}$.

The normalized plane wave is

$$\psi^{\mu\nu\dots} = \frac{1}{\sqrt{2\varepsilon}} u^{\mu\nu\dots} e^{-ipx}, \quad u_{\mu\nu\dots}^* u^{\mu\nu\dots} = -1, \quad (15.7)$$

with the wave amplitude satisfying the conditions

$$u^{\dots\mu\dots} p_\mu = 0. \quad (15.8)$$

There are $2s + 1$ independent polarization states.

Field quantization proceeds as an obvious generalization of the spin-0 or spin-1 cases.

The scheme presented above is entirely sufficient for the stated purpose—describing the field of free particles. The situation changes, however, if one poses the problem of describing the interaction of particles with the electromagnetic field. Such an interaction

would need to be incorporated into a Lagrangian from which all equations could be obtained without having to impose supplementary conditions separately. In practice, however, it turns out that this approach to describing the interaction works only for electrons—particles of spin $1/2$ (see § 32). For other spins the problem would therefore be of purely methodological interest.

Let us note that for all (integer and half-integer) spins $s > 1$ it proves impossible to formulate a variational principle using only a single function (tensorial or spinorial) whose rank matches the given spin. For this purpose it becomes necessary to introduce, as auxiliary quantities, tensorial or spinorial objects of lower rank as well. The Lagrangian is then chosen so that these auxiliary quantities automatically vanish by virtue of the free-particle field equations that follow from the variational principle¹⁸.

§ 16. Helicity states of a particle¹⁹

Within relativistic theory, the orbital angular momentum $\mathbf{l}$ and the spin $\mathbf{s}$ of a particle in motion lack individual conservation laws—only their sum, the total angular momentum $\mathbf{j} = \mathbf{l} + \mathbf{s}$, is conserved. Consequently, the spin projection onto a given direction (the z axis) is also nonconserved, rendering it unsuitable for labeling the polarization (spin) states of a particle in motion.

What is conserved, however, is the projection of the spin on the direction of the momentum: since $\mathbf{l} = [\mathbf{r}\mathbf{p}]$, the product $\mathbf{s}\mathbf{n}$ coincides with the conserved product $\mathbf{j}\mathbf{n}$ ($\mathbf{n} = \mathbf{p}/|\mathbf{p}|$). This quantity is called the *helicity*²⁰ (we have already considered it for the photon in § 8). Its eigenvalues will be denoted by λ ($\lambda = -s, \dots, +s$), and states of the particle with definite values of λ will be called helicity states.

Let $\psi_{\mathbf{p}\lambda}$ be the wave function (a plane wave) describing a state of the particle with definite $\mathbf{p}$ and λ , and let $u^{(\lambda)}(\mathbf{p})$ be its amplitude; for brevity we suppress the component indices of this function (for integer spin these are 4-tensor indices).

We saw in the preceding sections that a relativistic description of particles with nonzero (integer) spin requires introducing a wave function whose number of components exceeds $2s + 1$. The number of independent components, however, remains equal to $2s + 1$; the “extra” components are eliminated by imposing supplementary conditions, by virtue of which these components vanish in the rest frame (in the next chapter we shall see the same for half-integer s).

According to the transformation rules for angular momentum (see II, § 14), the helicity is invariant under Lorentz transformations that leave unchanged the direction of $\mathbf{p}$ onto which the angular momentum is projected. Therefore the number λ retains its meaning as a quantum number under such transformations, and to study the symmetry properties of helicity states one may use a reference frame in which $|\mathbf{p}| \ll m$ (in the limit, the rest frame). Then $\psi_{\mathbf{p}\lambda}$ reduces to a nonrelativistic $(2s + 1)$ -component wave function. Let us denote its amplitude by $w^{(\lambda)}(\mathbf{n})$, indicating as the argument the direction $\mathbf{n} = \mathbf{p}/|\mathbf{p}|$ along which the angular momentum is quantized. The amplitude $w^{(\lambda)}$ is an eigenfunction

¹⁸See Fierz M., Pauli W.//Proc. Roy. Soc. — 1939. — V. A 173. — P. 211. In that work the program indicated here was carried out for particles of spin $3/2$ and 2 .

¹⁹The content of this section applies to particles of any (integer or half-integer) spin.

²⁰In the Russian literature the term is *spiralnost'*.

of the operator $\mathbf{n}\hat{\mathbf{s}}$:

$$\mathbf{n}\hat{\mathbf{s}} w^{(\lambda)}(\mathbf{n}) = \lambda w^{(\lambda)}(\mathbf{n}). \quad (16.1)$$

In the spinor representation $w^{(\lambda)}$ is a contravariant symmetric spinor of rank $2s$; by the correspondence formulas (57.2) (see III) its components can also be labeled by the values of the spin projection σ on a fixed axis z ²¹.

In the momentum representation the wave functions of the states under consideration essentially coincide with the amplitudes $u^{(\lambda)}(\mathbf{p})$. Specifically:

$$\psi_{\mathbf{p}\lambda}(\mathbf{k}) = u^{(\lambda)}(\mathbf{k})\delta^{(2)}(\mathbf{v} - \mathbf{n}) = u^{(\lambda)}(\mathbf{p})\delta^{(2)}(\mathbf{v} - \mathbf{n}), \quad (16.2)$$

where the momentum as an independent variable is denoted $\mathbf{k}$, in contrast to its eigenvalue $\mathbf{p}$, and $\mathbf{v} = \mathbf{k}/|\mathbf{k}|$, in contrast to $\mathbf{n} = \mathbf{p}/|\mathbf{p}|$ ²². In the nonrelativistic limit

$$\psi_{\mathbf{n}\lambda}(\mathbf{v}) = w_{\sigma}^{(\lambda)}(\mathbf{v})\delta^{(2)}(\mathbf{v} - \mathbf{n}) = w^{(\lambda)}(\mathbf{n})\delta^{(2)}(\mathbf{v} - \mathbf{n}). \quad (16.3)$$

In more detail this expression should be written as

$$\psi_{\mathbf{n}\lambda}(\mathbf{v}, \sigma) = w_{\sigma}^{(\lambda)}(\mathbf{v})\delta^{(2)}(\mathbf{v} - \mathbf{n}),$$

where the discrete independent variable σ is also displayed explicitly.

The helicity operator $\hat{\mathbf{s}}\mathbf{n}$ is commutative with $\hat{j}_z$ and $\hat{\mathbf{j}}^2$. This follows from the fact that the angular-momentum operator corresponds to an infinitesimal coordinate rotation, while the dot product of any two vectors remains unchanged under arbitrary rotations. Consequently, stationary states exist in which the particle has simultaneously well-defined angular momentum j , its projection $j_z = m$, and helicity λ . We shall refer to such states as *spherical helicity states*.

We now find the wave functions of these states in the momentum representation. The construction proceeds by direct analogy with the expressions derived in III, § 103 for the wave functions of a symmetric top. Those results were obtained from the transformation formulas for wave functions under finite rotations (see III, § 58). Since that derivation relies exclusively on rotational-symmetry properties, it applies to momentum-representation functions just as it does to coordinate-space functions.

Along with the space-fixed coordinate system xyz (with respect to which the functions $\psi_{jm\lambda}$ are written), let us also introduce a “body-fixed” frame $\xi\eta\zeta$ with the ζ axis along the direction of $\mathbf{v}$. Without repeating the corresponding reasoning (cf. the derivation of formula (103.8) (see III)), we write

$$\psi_{jm\lambda}(\mathbf{k}) = \psi_{j\lambda}^{(0)} D_{\lambda m}^{(j)}(\mathbf{v}),$$

²¹The reasoning given here (together with the listing of allowed values of λ) pertains to particles whose mass differs from zero. For a particle of zero mass no rest frame exists, and the helicity admits only the two values $\lambda = \pm s$. This restriction traces back to the circumstance noted already in § 8: states of such a particle fall into representations of the axial-symmetry group, which permits at most a twofold degeneracy of levels (from the viewpoint of the wave equation this signifies that upon passing to the limit $m \rightarrow 0$ the system of equations for a spin- s particle separates into independent equations describing massless particles of spins $s, s-1, \dots$). In particular, for the photon $\lambda = \pm 1$, and the three-dimensional vectors $\mathbf{e}^{(\pm 1)}$ (8.2) assume the role of the corresponding amplitudes $w^{(\lambda)}$.

²²Here the delta function $\delta^{(2)}$ is defined so that $\int \delta^{(2)}(\mathbf{v} - \mathbf{n}) d\mathbf{o}_{\mathbf{v}} = 1$. In (16.2) (and in the analogous case below, see (16.4)) the delta function ensuring the given value of the energy has been omitted.

where $\psi_{j\lambda}^{(0)}$ is the wave function in the “body-fixed” frame describing a state with a definite value of the ζ -projection of the angular momentum: $j_\zeta = \lambda$; in the momentum representation this function evidently coincides with the amplitude $u^{(\lambda)}$. The normalized (see below) wave function is

$$\psi_{jm\lambda}(\mathbf{k}) = \sqrt{\frac{2j+1}{4\pi}} D_{\lambda m}^{(j)}(\mathbf{v}) u^{(\lambda)}(\mathbf{k}). \quad (16.4)$$

Here, however, a question arises regarding the choice of phases, connected with the following ambiguity. The rotation of the system xyz relative to $\xi\eta\zeta$ is specified by three Euler angles α, β, γ ; the direction of $\mathbf{v}$, on which alone the wave function of the particle can depend, is determined only by two spherical angles $\alpha \equiv \varphi, \beta \equiv \theta$. One must therefore adopt some convention for the angle γ . We shall set $\gamma = 0$, i.e. define $D_{\lambda m}^{(j)}(\mathbf{v})$ as

$$D_{\lambda m}^{(j)}(\mathbf{v}) = D_{\lambda m}^{(j)}(\varphi, \theta, 0) = e^{im\varphi} d_{\lambda m}^{(j)}(\theta). \quad (16.5)$$

By virtue of (58.21) (see III) the functions (16.5) satisfy the orthogonality and normalization conditions:

$$\int D_{\lambda_1 m_1}^{(j_1)*}(\mathbf{v}) D_{\lambda_2 m_2}^{(j_2)}(\mathbf{v}) \frac{d\mathbf{v}}{4\pi} = \frac{1}{2j+1} \delta_{j_1 j_2} \delta_{m_1 m_2} \quad (16.6)$$

($d\mathbf{v} = \sin\theta d\theta d\varphi$). Orthogonality in the index λ is guaranteed by the amplitude factor $u^{(\lambda)}$. The functions $\psi_{jm\lambda}$ are therefore mutually orthogonal, as required, across all three labels $jm\lambda$, and the prefactor adopted in (16.4) ensures their normalization according to

$$\int |\psi_{jm\lambda}|^2 d\mathbf{v} = 1. \quad (16.7)$$

Here it is assumed that the amplitudes $u^{(\lambda)}$ are normalized to unity: $u^{(\lambda)} u^{(\lambda)*} = 1$.

Let us consider the behavior of helicity-state wave functions under coordinate inversion. The product of the polar vector $\mathbf{v}$ and the axial vector $\mathbf{j}$ is a pseudoscalar. It is therefore clear that upon inversion a state of helicity λ goes over into a state with $-\lambda$; one must, however, determine the phase factors in these transformations.

Under inversion $\mathbf{v} \rightarrow -\mathbf{v}$. The vector $\mathbf{v}$ is specified by two angles φ, θ , and the transformation $\mathbf{v} \rightarrow -\mathbf{v}$ is effected by the replacement $\varphi \rightarrow \varphi + \pi, \theta \rightarrow \pi - \theta$. This determines the axis ζ , but the position of the axes ξ and η , which also depends on the third Euler angle γ , remains unspecified; the transformation of θ and φ alone does not allow one to distinguish between an inversion and a rotation of the axis ζ in this sense. In terms of all three Euler angles the inversion is the transformation

$$\alpha \equiv \varphi \rightarrow \varphi + \pi, \quad \beta \equiv \theta \rightarrow \pi - \theta, \quad \gamma \rightarrow \pi - \gamma. \quad (16.8)$$

Therefore, if $D_{\lambda m}^{(j)}(\mathbf{v})$ is defined according to (16.5) (i.e. with $\gamma = 0$), and the substitution $\mathbf{v} \rightarrow -\mathbf{v}$ is understood as the result of an inversion, then

$$D_{\lambda m}^{(j)}(-\mathbf{v}) = D_{\lambda m}^{(j)}(\varphi + \pi, \pi - \theta, \pi). \quad (16.9)$$

Using formulas (58.9), (58.16), (58.18) (see III) we therefore find

$$D_{\lambda m}^{(j)}(-\mathbf{v}) = e^{i\lambda\pi} d_{\lambda m}^{(j)}(\pi - \theta) e^{im(\varphi + \pi)} =$$

$$= (-1)^{j-\lambda} e^{im\varphi} d_{-\lambda m}^{(j)}(\theta) = (-1)^{j-\lambda} D_{-\lambda m}^{(j)}(\mathbf{v}),$$

or

$$D_{\lambda m}^{(j)}(-\mathbf{v}) = (-1)^{-\lambda} D_{-\lambda m}^{(j)}(\mathbf{v}) \quad (16.10)$$

($j - \lambda$ is an integer).

An analogous formula for the spinor $w^{(\lambda)}$ can be obtained by noting that its components $w_{\sigma}^{(\lambda)}$ coincide, up to a factor, with the functions

$$w_{\sigma}^{(\lambda)}(\mathbf{v}) \propto D_{\lambda \sigma}^{(s)}(\mathbf{v})^*. \quad (16.11)$$

Indeed, applying the transformation formula (58.7) (see III) to the spin eigenfunctions and setting the ζ -projection of the spin to the definite value λ (i.e. replacing $\psi_{jm'}$ on the right-hand side of (58.7) (see III) by $\delta_{m'\lambda}$), we find that $D_{\lambda \sigma}^{(s)}(\mathbf{v})$ are spin wave functions corresponding to definite values of the z - and ζ -projections (σ and λ). The totality of these functions ($\sigma = -s, \dots, +s$) forms (by the correspondence formulas (57.6) (see III)) a covariant spinor of rank $2s$. The components of the contravariant spinor (57.2) (see III), however—corresponding to the components $w_{\sigma}^{(\lambda)}$ —transform as the complex conjugates of the components of a covariant spinor of the same rank. From (16.10), (16.11) we have

$$w^{(\lambda)}(-\mathbf{v}) = (-1)^{s-\lambda} w^{(-\lambda)}(\mathbf{v}) \quad (16.12)$$

($s - \lambda$ is an integer). The inversion operation applied to $w^{(\lambda)}$, however, consists not only in the replacement $\mathbf{v} \rightarrow -\mathbf{v}$, but also in multiplication by an overall phase factor (the “intrinsic parity” of the particle), which we denote by η :

$$\hat{P}w^{(\lambda)}(\mathbf{v}) = \eta w^{(\lambda)}(-\mathbf{v}) = \eta(-1)^{s-\lambda} w^{(-\lambda)}(\mathbf{v}). \quad (16.13)$$

For the relativistic amplitude $u^{(\lambda)}(\mathbf{k})$ this transformation takes the form

$$\hat{P}u^{(\lambda)}(\mathbf{k}) = \eta \beta u^{(\lambda)}(-\mathbf{k}) = \eta(-1)^{s-\lambda} u^{(-\lambda)}(\mathbf{k}), \quad (16.14)$$

where β is a certain matrix that acts as the identity on the components of $u^{(\lambda)}$ surviving in the limit $|\mathbf{p}| \rightarrow 0$. It is important that this matrix is independent of the quantum numbers of the state, so that the difference between (16.13) and (16.14) is in this sense inessential²⁴.

Applying (16.14) to (16.2), we obtain the transformation law for the wave functions of the states $|\mathbf{n}\lambda\rangle$:

$$\hat{P}\psi_{\mathbf{n}\lambda}(\mathbf{v}) = \eta(-1)^{s-\lambda} \psi_{-\mathbf{n}-\lambda}(\mathbf{v}). \quad (16.15)$$

For the spherical helicity states, making use of (16.10) and (16.12), we obtain the transformation law:

$$\hat{P}\psi_{jm\lambda}(\mathbf{v}) = \eta(-1)^{j-s} \psi_{jm-\lambda}(\mathbf{v}). \quad (16.16)$$

²⁴Thus, for $s = 1$ the amplitudes $u^{(\lambda)}$ are 4-vectors (16.22); in this case β is the full identity matrix with respect to 4-vector indices: $\beta_{\mu\nu} = \delta_{\mu\nu}$. For $s = 1/2$, $u^{(\lambda)}$ is a bispinor; the phase factor is then $\eta = i$, and β is the Dirac matrix γ^0 (see (21.10)).

The states ψ_{jm0} transform, according to (16.16), into themselves, i.e. they possess a definite parity. If, however, $\lambda \neq 0$, then only superpositions of states with opposite helicities have a definite parity:

$$\psi_{jm|\lambda|}^{(\pm)} = \frac{1}{\sqrt{2}} (\psi_{jm\lambda} \pm \psi_{jm-\lambda}). \quad (16.17)$$

Under inversion they transform into themselves according to

$$\hat{P}\psi_{jm|\lambda|}^{(\pm)}(\mathbf{v}) = \pm\eta(-1)^{j-s}\psi_{jm|\lambda|}^{(\pm)}(\mathbf{v}). \quad (16.18)$$

Let us draw attention to the fact that in this section we have classified the states of a free particle with given angular momentum by operating solely with conserved quantities, without appealing to the concept of orbital angular momentum (which was used, for instance, in §§ 6, 7 for the classification of photon states).

As an illustration consider the case of spin 1. In the rest frame the amplitudes $u^{(\lambda)}$ (4-vectors) reduce to three-dimensional vectors $\mathbf{e}^{(\lambda)}$, which play here the role of the amplitudes $w^{(\lambda)}$. The action of the spin-1 operator on a vector function $\mathbf{e}$ is given by

$$(\hat{S}_i \mathbf{e})_k = -ie_{ikl}e_l \quad (16.19)$$

(see III, § 57, Problem 2). Equation (16.1) therefore assumes the form

$$i[\mathbf{n}\mathbf{e}^{(\lambda)}] = \lambda\mathbf{e}^{(\lambda)}. \quad (16.20)$$

Its solutions (in a coordinate frame $\xi\eta\zeta$ with the ζ axis along $\mathbf{n}$) coincide with the circular unit vectors (7.14)²⁶:

$$\mathbf{e}^{(0)} = i(0, 0, 1), \quad \mathbf{e}^{(\pm 1)} = \mp \frac{i}{\sqrt{2}}(1, \pm i, 0). \quad (16.21)$$

In a frame where the particle has momentum $\mathbf{p}$, the helicity-state amplitudes are the 4-vectors

$$u^{(0)\mu} = \left(\frac{|\mathbf{p}|}{m}, \frac{\varepsilon}{m} \mathbf{e}^{(0)} \right), \quad u^{(\pm 1)\mu} = (0, \mathbf{e}^{(\pm 1)}). \quad (16.22)$$

If $\mathbf{e}$ is a polar vector, then $\eta = -1$. In that case the functions (16.17) (which for $s = 1$ are three-dimensional vectors) have the following parities:

$$\begin{aligned} \psi_{jm|\lambda|}^{(+)} : \quad P &= (-1)^j, \\ \psi_{jm|\lambda|}^{(-)} : \quad P &= (-1)^{j+1}, \\ \psi_{jm0} : \quad P &= (-1)^j. \end{aligned}$$

Comparing with the definition of vector spherical harmonics (7.4), we see that these functions are identical (up to phase factors) with $\mathbf{Y}_{jm}^{(e)}$, $\mathbf{Y}_{jm}^{(m)}$, $\mathbf{Y}_{jm}^{(l)}$ respectively.

²⁶The choice of phase factors is fixed by the requirement that the matrix elements of the spin operators computed with the eigenfunctions (16.21) conform to the general definitions in III, §§ 27, 107.

Determining the phase factors (say, by comparing the values at $\theta = 0$), we arrive at the equalities

$$\begin{aligned} \mathbf{Y}_{jm}^{(e)} &= i^{j-1} \sqrt{\frac{2j+1}{8\pi}} \left(\mathbf{e}^{(1)} D_{1m}^{(j)}(\nu) + \mathbf{e}^{(-1)} D_{-1m}^{(j)}(\nu) \right), \\ \mathbf{Y}_{jm}^{(m)} &= i^{j-1} \sqrt{\frac{2j+1}{8\pi}} \left(\mathbf{e}^{(1)'} D_{1m}^{(j)}(\nu) + \mathbf{e}^{(-1)'} D_{-1m}^{(j)}(\nu) \right), \\ \mathbf{Y}_{jm}^{(l)} &= i^{j-1} \sqrt{\frac{2j+1}{4\pi}} \mathbf{e}^{(0)} D_{0m}^{(j)} \end{aligned} \quad (16.23)$$

(j is an integer!); $\mathbf{e}^{(\lambda)'} = [\mathbf{n}\mathbf{e}^{(\lambda)}]$ are circular unit vectors in axes $\xi'\eta'\zeta$ that are rotated relative to $\xi\eta\zeta$ by 90° about the ζ axis.

The last of formulas (16.23) is equivalent to the expression (58.23) (see III) for $d_{0m}^{(j)}(\theta)$. From the first (or the second) formula one can obtain a simple expression for the functions $d_{\pm 1m}^{(j)}$. We have

$$i^{j-1} \sqrt{\frac{2j+1}{8\pi}} D_{\pm 1m}^{(j)} = \mathbf{Y}_{jm}^{(e)} \mathbf{e}^{(\pm 1)*} = \frac{1}{\sqrt{j(j+1)}} \mathbf{e}^{(\pm 1)*} \nabla Y_{jm}.$$

We expand the scalar product on the right-hand side in the frame $\xi\eta\zeta$, where

$$\left(\frac{\partial}{\partial \xi}, \frac{\partial}{\partial \eta} \right) \rightarrow \left(\frac{\partial}{\partial \theta}, \frac{1}{\sin \theta} \frac{\partial}{\partial \varphi} \right).$$

Recalling definition (7.2) of the function Y_{jm} and definition (16.5), we obtain as the final result

$$d_{\pm 1m}^{(j)}(\theta) = (-1)^{m+1} \sqrt{\frac{(j-m)!}{(j+m)!j(j+1)}} \left(\pm \frac{\partial}{\partial \theta} + \frac{m}{\sin \theta} \right) P_j^m(\cos \theta), \quad m \geq 0. \quad (16.24)$$

Fermions

§ 17. Four-dimensional spinors

In nonrelativistic theory a particle of arbitrary spin s is described by a $(2s + 1)$ -component quantity—a symmetric spinor of rank $2s$. From the mathematical standpoint these are quantities that furnish the irreducible representations of the spatial rotation group.

In relativistic theory this group appears only as a subgroup of the broader group of four-dimensional rotations—the Lorentz group. In connection with this there arises the need to construct a theory of four-dimensional spinors (4-spinors)—quantities that carry the irreducible representations of the Lorentz group; the exposition of this theory occupies §§ 17–19. In §§ 17, 18 only the proper Lorentz group is considered, which does not include spatial inversion; the latter will be treated in § 19.

The construction of the 4-spinor theory proceeds along the same lines as the three-dimensional spinor theory (*B. L. van der Waerden*, 1929; *G. E. Uhlenbeck*, *O. Laporte*, 1931).

A spinor ξ^α is a two-component quantity ($\alpha = 1, 2$); regarded as components of the wave function of a spin-1/2 particle, ξ^1 and ξ^2 correspond to eigenvalues of the z -projection of the spin equal to $+1/2$ and $-1/2$ respectively. Under any (proper) Lorentz transformation the two quantities ξ^1, ξ^2 transform through each other:

$$\xi^{1'} = \alpha\xi^1 + \beta\xi^2, \quad \xi^{2'} = \gamma\xi^1 + \delta\xi^2. \quad (17.1)$$

The coefficients $\alpha, \beta, \gamma, \delta$ are definite functions of the rotation angles of the 4-coordinate system, subject to the condition

$$\alpha\delta - \beta\gamma = 1, \quad (17.2)$$

i.e. the determinant of the binary transformation (17.1) equals 1, just as the determinants of coordinate transformations in the Lorentz group do.

By virtue of conditions (17.2) the bilinear form $\xi^1\Xi^2 - \xi^2\Xi^1$ (where ξ^α and Ξ^α are two spinors) is invariant under the transformation (17.1) (it corresponds to a spin-0 particle “composed” of two spin-1/2 particles). For a natural way of writing such invariant expressions, alongside the “contravariant” components of the spinor ξ^α one also introduces “covariant” components ξ_α . The passage from one set to the other is carried out with the aid of the “spinor metric” $g_{\alpha\beta}$ ¹:

$$\xi_\alpha = g_{\alpha\beta}\xi^\beta, \quad (17.3)$$

where

$$g = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (17.4)$$

¹ Spinor indices will be denoted by the first letters of the Greek alphabet: $\alpha, \beta, \gamma, \dots$.

so that

$$\xi_1 = \xi^2, \quad \xi_2 = -\xi^1. \quad (17.5)$$

The invariant $\xi^1 \Xi^2 - \xi^2 \Xi^1$ is then written as a scalar product $\xi^\alpha \Xi_\alpha$. Note that $\xi^\alpha \Xi_\alpha = -\xi_\alpha \Xi^\alpha$.

Up to this point the properties listed above have formally coincided with those of three-dimensional spinors. A difference arises, however, when one turns to complex-conjugate spinors.

In nonrelativistic theory the sum

$$\psi^1 \psi^{1*} + \psi^2 \psi^{2*}, \quad (17.6)$$

which governs the probability density of finding particles at a given point in space, had to be a scalar, and for this purpose the components $\psi^{\alpha*}$ had to transform as covariant spinor components; put differently, the transformation (17.1) had to be unitary ($\alpha = \delta^*$, $\beta = -\gamma^*$). In the relativistic case, however, the particle density does not behave as a scalar; it forms the time component of a 4-vector. Owing to this, the requirement just stated ceases to hold, and no further restrictions on the transformation coefficients arise (apart from (17.2)). Four complex quantities $\alpha, \beta, \gamma, \delta$ constrained only by (17.2) amount to $8 - 2 = 6$ real parameters—in agreement with the number of angles specifying a 4-coordinate rotation (rotations in six coordinate planes).

Consequently, complex-conjugate binary transformations prove to be essentially different from one another, and in relativistic theory two varieties of spinors arise. To distinguish between them one adopts special notation: the indices of spinors whose transformation law is the complex conjugate of (17.1) are written as digits with dots placed above them (*dotted indices*). Thus, by definition,

$$\eta^{\dot{\alpha}} \sim \xi^{\alpha*}, \quad (17.7)$$

where the sign $\sim$ stands for “transforms as.” In other words, the transformation formulas for a “dotted” spinor are:

$$\eta^{\dot{1}'} = \alpha^* \eta^{\dot{1}} + \beta^* \eta^{\dot{2}}, \quad \eta^{\dot{2}'} = \gamma^* \eta^{\dot{1}} + \delta^* \eta^{\dot{2}}. \quad (17.8)$$

The operations of lowering and raising dotted indices are performed in the same way as for undotted ones:

$$\eta_{\dot{1}} = \eta^{\dot{2}}, \quad \eta_{\dot{2}} = -\eta^{\dot{1}}. \quad (17.9)$$

With respect to spatial rotations the behavior of 4-spinors coincides with that of 3-spinors. For the latter, as we know, $\psi_\alpha^* \sim \psi^\alpha$. By virtue of definition (17.7) the 4-spinor $\eta_{\dot{\alpha}}$ therefore behaves under rotations as a contravariant 3-spinor ψ^α . The eigenvalues of the spin projection $1/2$ and $-1/2$ accordingly correspond to the covariant components $\eta_{\dot{1}}$ and $\eta_{\dot{2}}$.

Spinors of higher rank are defined as sets of quantities that transform like products of components of several first-rank spinors. The indices of a higher-rank spinor may include both dotted and undotted ones. For instance, three types of second-rank spinors exist:

$$\xi^{\alpha\beta} \sim \xi^\alpha \Xi^\beta, \quad \zeta^{\alpha\dot{\beta}} \sim \xi^\alpha \eta^{\dot{\beta}}, \quad \eta^{\dot{\alpha}\dot{\beta}} \sim \eta^{\dot{\alpha}} H^{\dot{\beta}}.$$

Merely stating the total rank of a spinor is therefore insufficient for an unambiguous definition of the concept; when necessary we shall specify the rank as a pair of numbers (k, l) —the number of undotted and the number of dotted indices.

Since the transformations (17.1) and (17.8) are algebraically independent, there is no need to fix the ordering of dotted and undotted indices (in this sense, for example, the spinors $\zeta^{\alpha\dot{\beta}}$ and $\zeta^{\dot{\beta}\alpha}$ are one and the same).

For an expression involving spinors to have an invariant character, it must carry the same number of undotted and dotted indices on both sides; otherwise the expression will certainly be violated upon passing from one reference frame to another. One should keep in mind here that complex conjugation entails the interchange of dotted and undotted indices. Therefore the relation $\eta^{\dot{\alpha}\dot{\beta}} = (\xi^{\alpha\beta})^*$ between two spinors does have an invariant character.

Contraction of spinors or their products can be performed only over pairs of indices of the same kind—two dotted or two undotted. Summation over a pair of indices of different kinds is not an invariant operation. For this reason, from the spinor

$$\zeta^{\alpha_1 \alpha_2 \dots \alpha_k \dot{\beta}_1 \dot{\beta}_2 \dots \dot{\beta}_l}, \quad (17.10)$$

which is symmetric in all k undotted and in all l dotted indices, it is impossible to form a spinor of lower rank (recall that contracting over a pair of indices in which the spinor is symmetric gives zero). This means that from the quantities (17.10) one cannot assemble any smaller set of their linear combinations that would transform among themselves under every group transformation. Stated differently, symmetric 4-spinors realize the irreducible representations of the proper Lorentz group. Every such representation is labeled by a pair of numbers (k, l) .

Since each spinorial index runs over two values, there are $k + 1$ essentially distinct sets of numbers $\alpha_1 \alpha_2 \dots \alpha_k$ in (17.10) (containing 0, 1, 2, $\dots$, k ones and $k, k - 1, \dots$, 0 twos) and $l + 1$ sets of numbers $\dot{\beta}_1 \dot{\beta}_2 \dots \dot{\beta}_l$. Altogether, therefore, a symmetric spinor of rank (k, l) has $(k + 1)(l + 1)$ independent components; this is the dimensionality of the irreducible representation it carries.

§ 18. Relation of spinors to 4-vectors

A spinor $\zeta^{\alpha\dot{\beta}}$ carrying one dotted and one undotted index possesses $2 \cdot 2 = 4$ independent components—precisely the number of components belonging to a 4-vector. It is therefore evident that both objects furnish one and the same irreducible representation of the proper Lorentz group, and a definite correspondence must hold between their respective components.

To establish this correspondence we turn first of all to the analogous correspondence in the three-dimensional case, bearing in mind that with respect to purely spatial rotations the behavior of 3- and 4-spinors must be identical.

For a three-dimensional spinor $\psi^{\alpha\dot{\beta}}$ the correspondence formulas hold (see III, § 57),

which we write here in the form

$$\begin{aligned} a_x &= \frac{1}{2}(\psi^{22} - \psi^{11}) = \frac{1}{2}(\psi_1^2 + \psi_2^1), \\ a_y &= -\frac{i}{2}(\psi^{22} + \psi^{11}) = \frac{i}{2}(\psi_2^1 - \psi_1^2), \\ a_z &= \frac{1}{2}(\psi^{12} + \psi^{21}) = \frac{1}{2}(\psi_1^1 - \psi_2^2), \end{aligned} \quad (\text{III.1})$$

where a_x, a_y, a_z are the components of a certain three-dimensional vector $\mathbf{a}$. In passing to the four-dimensional case one must replace the components ψ_β^α by $\zeta^{\alpha\beta}$, and interpret a_x, a_y, a_z as the contravariant components a^1, a^2, a^3 of a 4-vector. As for the expression for the fourth component, a^0 , its form is already clear from the circumstance noted in § 17: the quantity (17.6) must transform as a^0 . Hence $a^0 \sim \zeta^{11} + \zeta^{22}$; the proportionality coefficient is fixed by requiring that the scalar $\zeta_{\alpha\beta}\zeta^{\alpha\beta}$ coincide with $2a_\mu a^\mu \equiv 2a^2$.

We thus arrive at the following correspondence formulas:

$$\begin{aligned} a^1 &= \frac{1}{2}(\zeta^{12} + \zeta^{21}), & a^2 &= \frac{i}{2}(\zeta^{12} - \zeta^{21}), \\ a^3 &= \frac{1}{2}(\zeta^{11} - \zeta^{22}), & a^0 &= \frac{1}{2}(\zeta^{11} + \zeta^{22}). \end{aligned} \quad (18.1)$$

The inverse formulas are:

$$\begin{aligned} \zeta^{11} &= \zeta_{22} = a^3 + a^0, & \zeta^{22} &= \zeta_{11} = a^0 - a^3, \\ \zeta^{12} &= -\zeta_{21} = a^1 - ia^2, & \zeta^{21} &= -\zeta_{12} = a^1 + ia^2. \end{aligned} \quad (18.2)$$

Here

$$\zeta_{\alpha\beta}\zeta^{\alpha\beta} = 2a^2. \quad (18.3)$$

We note also that

$$\zeta_{\alpha\beta}\zeta^{\gamma\beta} = \delta_\alpha^\gamma a^2. \quad (18.4)$$

To see why this last relation holds, observe that the quantity $\zeta_{\alpha\beta}\zeta_\gamma^{\beta}$, viewed as a second-rank spinor in the indices α and γ , is antisymmetric and hence must be proportional to the metric spinor.

The link between $\zeta^{\alpha\beta}$ and a 4-vector is actually one instance of a broader principle: any symmetric spinor of rank (k, k) can be placed in one-to-one correspondence with an irreducible symmetric 4-tensor of rank k (irreducibility meaning that contraction over every index pair gives zero).

The relationship linking a spinor to a 4-vector can be written compactly with the aid of the two-by-two Pauli matrices²:

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (18.5)$$

Denoting symbolically by ζ the matrix of the quantities $\zeta^{\alpha\beta}$ with upper indices (the first being undotted), formulas (18.2) take the form

$$\zeta = \mathbf{a}\boldsymbol{\sigma} + a^0 \quad (18.6)$$

²For brevity of notation, operators (matrices) that act on spin variables will be denoted by letters without carets.

(in the second term one understands, of course, the product of a^0 with the unit matrix). The inverse relations are:

$$\mathbf{a} = \frac{1}{2} \text{Sp}(\zeta \boldsymbol{\sigma}), \quad a^0 = \frac{1}{2} \text{Sp} \zeta. \quad (18.7)$$

With the help of formulas (18.6), (18.7) one can establish the connection between the transformation laws for the 4-vector and the spinor, thereby expressing the spinor transformation law through the parameters of rotations of the 4-coordinate system.

Let us write the transformation of the spinor ξ^α in the form

$$\xi^{\alpha'} = (B\xi)^\alpha, \quad B = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}, \quad (18.8)$$

where B is a two-by-two matrix assembled from the coefficients of the binary transformation. The transformation of the dotted spinor is then:

$$\eta^{\dot{\beta}'} = (B^* \eta)^{\dot{\beta}} = (\eta B^+)^{\dot{\beta}}, \quad (18.9)$$

and the transformation of the second-rank spinor $\zeta^{\alpha\dot{\beta}} \sim \xi^\alpha \eta^{\dot{\beta}}$ can be written symbolically as $\zeta' = B\zeta B^{+1}$. For an infinitesimally small transformation $B = 1 + \lambda$, where λ is a small matrix, and to first-order accuracy

$$\zeta' = \zeta + (\lambda\zeta + \zeta\lambda^+). \quad (18.10)$$

Consider first the Lorentz transformation to a reference frame moving with an infinitesimally small velocity $\delta\mathbf{V}$ (without altering the orientation of the spatial axes). Under this transformation the 4-vector $a^\mu = (a^0, \mathbf{a})$ changes according to

$$\mathbf{a}' = \mathbf{a} - a^0 \delta\mathbf{V}, \quad a^{0'} = a^0 - \mathbf{a} \delta\mathbf{V}. \quad (18.11)$$

Let us now make use of formulas (18.7). The transformation of a^0 can be expressed, on the one hand, as

$$a^{0'} = a^0 - \mathbf{a} \delta\mathbf{V} = a^0 - \frac{1}{2} \text{Sp}(\zeta \boldsymbol{\sigma} \delta\mathbf{V}),$$

and on the other hand as

$$a^{0'} = \frac{1}{2} \text{Sp} \zeta' = a^0 + \frac{1}{2} \text{Sp} \zeta (\lambda + \lambda^+).$$

These expressions must coincide identically (i.e. for arbitrary ζ). From this we obtain the equality:

$$\lambda + \lambda^+ = -\boldsymbol{\sigma} \delta\mathbf{V}.$$

By the same method, examining the transformation of $\mathbf{a}$, we find

$$\boldsymbol{\sigma} \lambda + \lambda^+ \boldsymbol{\sigma} = -\delta\mathbf{V}.$$

These equalities, regarded as equations for λ , yield the solution:

$$\lambda = \lambda^+ = -\frac{1}{2} \boldsymbol{\sigma} \delta\mathbf{V}.$$

Thus the infinitesimally small Lorentz transformation of the spinor ξ^α is effected by the matrix

$$B = 1 - \frac{1}{2}(\boldsymbol{\sigma}\mathbf{n})\delta V, \quad (18.12)$$

where $\mathbf{n}$ is a unit vector along the velocity $\delta\mathbf{V}$. From this it is straightforward to obtain the transformation for a finite velocity $\mathbf{V}$. To do so, recall that the Lorentz transformation amounts (geometrically) to a rotation of the 4-coordinate system in the $t\mathbf{n}$ plane through an angle φ related to the velocity $\mathbf{V}$ by $\text{th } \varphi = V^3$. An infinitesimally small transformation corresponds to the angle $\delta\varphi = \delta V$, while a rotation through a finite angle φ is achieved by repeating the rotation through $\delta\varphi$ a total of $\varphi/\delta\varphi$ times. Raising the operator (18.12) to the power $\varphi/\delta\varphi$ and passing to the limit $\delta\varphi \rightarrow 0$, we obtain

$$B = e^{-\frac{\varphi}{2}\mathbf{n}\boldsymbol{\sigma}}. \quad (18.13)$$

The mathematical meaning of this operator becomes transparent once one observes that, by the properties of the Pauli matrices, all even powers of $\mathbf{n}\boldsymbol{\sigma}$ equal 1 while all odd powers equal $\mathbf{n}\boldsymbol{\sigma}$. Noting that ch decomposes into even powers and sh into odd powers of its argument, we arrive at the final result

$$B = \text{ch } \frac{\varphi}{2} - \mathbf{n}\boldsymbol{\sigma} \text{sh } \frac{\varphi}{2}, \quad \text{th } \varphi = V. \quad (18.14)$$

We note that the Lorentz-transformation matrices B turn out to be Hermitian: $B = B^+$.

Consider now an infinitesimally small rotation of the spatial coordinate system. Under such a rotation the three-dimensional vector $\mathbf{a}$ transforms according to

$$\mathbf{a}' = \mathbf{a} - [\delta\boldsymbol{\theta} \mathbf{a}], \quad (18.15)$$

where $\delta\boldsymbol{\theta}$ is the vector of the infinitesimally small rotation angle. The corresponding spinor transformation could be found in an analogous manner. There is no need for this, however, since under spatial rotations the behavior of 4-spinors coincides with that of 3-spinors, and for the latter the transformation is known beforehand from the general connection between the spin operator and the operator of an infinitesimally small rotation:

$$B = 1 + \frac{i}{2}\boldsymbol{\sigma} \delta\boldsymbol{\theta}. \quad (18.16)$$

The passage to a rotation through a finite angle θ is carried out in the same way as the passage from (18.12) to (18.14):

$$B = \exp\left(\frac{i\theta}{2}\mathbf{n}\boldsymbol{\sigma}\right) = \cos \frac{\theta}{2} + i\mathbf{n}\boldsymbol{\sigma} \sin \frac{\theta}{2}, \quad (18.17)$$

where $\mathbf{n}$ is the unit vector along the rotation axis. This matrix is unitary ($B^+ = B^{-1}$), as must be the case for a spatial rotation.

³We recall that in planes containing the time axis the metric is pseudo-Euclidean.

§ 19. Inversion of spinors

When developing the three-dimensional spinor theory (in Vol. III) we did not consider how spinors behave with respect to the spatial inversion operation, inasmuch as in the nonrelativistic framework this would not have produced any new physical consequences. It is worth pausing on this matter now, however, for a clearer understanding of the inversion properties of 4-spinors that will be discussed below.

The inversion operation does not alter the sign of an axial vector, and the spin vector is precisely of this type. Consequently the value of its projection s_z remains unchanged as well. It follows that under inversion each component of the spinor ψ^α can transform only into itself, i.e. one must have

$$\psi^\alpha \rightarrow P\psi^\alpha, \quad (19.1)$$

where P is a constant coefficient. Applying inversion twice returns us to the original coordinate system. For spinors, however, the return to the initial configuration can be understood in two distinct senses: as a rotation of the system by 0° or by 360° . For spinors these two interpretations are not equivalent, since ψ^α reverses sign under a 360° rotation. Two alternative definitions of inversion therefore arise: in the first,

$$P^2 = 1, \quad P = \pm 1, \quad (19.2)$$

and in the second,

$$P^2 = -1, \quad P = \pm i. \quad (19.3)$$

A crucial requirement is that inversion be defined in the same manner for every spinor. It would be inconsistent for distinct spinors to follow different inversion rules (one obeying (19.2), another (19.3)), for then one could not always build a scalar (or pseudoscalar) out of an arbitrary pair of spinors: if ψ^α were governed by (19.2) while φ^α were governed by (19.3), the product $\psi^\alpha \varphi_\alpha$ would pick up a factor $\pm i$ upon inversion instead of staying fixed (or simply flipping sign).

It should also be stressed that (under either definition of inversion) the assignment of one or another parity P to a spinor carries no absolute significance, because spinors change sign under a rotation through 2π , which can always be performed simultaneously with the inversion. What does possess an absolute character, however, is the “relative parity” of two spinors, defined as the parity of the scalar $\psi^\alpha \varphi_\alpha$ formed from them; since a 2π rotation reverses the sign of all spinors simultaneously, the associated ambiguity does not affect the parity of the scalar in question.

Let us turn now to four-dimensional spinors.

We note first that since inversion changes the sign of only three (x, y, z) out of the four (t, x, y, z) coordinates, it commutes with spatial rotations but does not commute with transformations that rotate the t axis. If $\widehat{L}$ denotes the Lorentz transformation to a frame moving with velocity $\mathbf{V}$, then $\widehat{P}\widehat{L} = \widehat{L}'\widehat{P}$, where $\widehat{L}'$ is the transformation to a frame moving with velocity $-\mathbf{V}$.

It follows from this that under inversion the components of a 4-spinor ξ^α cannot transform through themselves alone. If the inversion of the spinor ξ^α were still given by the transformation (19.1) (i.e. were represented by a matrix proportional to the identity),

it would commute with all Lorentz transformations whatsoever, which certainly must not be the case (since the operations $\widehat{L}$ and $\widehat{L}'$ applied to ξ^α manifestly do not coincide).

Inversion must therefore map the components of the spinor ξ^α into other quantities. These can only be the components of some other spinor $\eta_{\dot{\alpha}}$ whose transformation properties differ from those of ξ^α . Since inversion leaves the z -projection of the spin unchanged (as already noted above), the components ξ^1 and ξ^2 can pass under inversion only into η_1 and η_2 , which correspond to the same values $s_z = 1/2$ and $s_z = -1/2$. Taking inversion to be an operation that yields 1 when applied twice, one can specify its action by the formulas

$$\xi^\alpha \rightarrow \eta_{\dot{\alpha}}, \quad \eta_{\dot{\alpha}} \rightarrow \xi^\alpha. \quad (19.4)$$

For the covariant components ξ_α and the contravariant $\eta^{\dot{\alpha}}$ these transformations carry the opposite sign:

$$\xi_\alpha \rightarrow -\eta^{\dot{\alpha}}, \quad \eta^{\dot{\alpha}} \rightarrow -\xi_\alpha, \quad (19.4a)$$

since lowering and raising one and the same index are performed with opposite signs, see (17.5) and (17.9)⁴.

A certain distinction between the two definitions of inversion lies in the fact that under the second definition complex-conjugate spinors transform identically: if $\Xi_\alpha = \eta_{\dot{\alpha}}^*$, $H^{\dot{\alpha}} = \xi^{\alpha*}$, then from (19.5) one obtains $\Xi_\alpha \rightarrow -iH^{\dot{\alpha}}$, $H^{\dot{\alpha}} \rightarrow -i\Xi_\alpha$, i.e. the same rule as for $\xi_\alpha, \eta^{\dot{\alpha}}$. Under definition (19.4), by contrast, one would obtain the transformation $\Xi_\alpha \rightarrow H^{\dot{\alpha}}$, $H^{\dot{\alpha}} \rightarrow \Xi_\alpha$, which is the reverse in sign of the transformation of the spinors $\xi_\alpha, \eta^{\dot{\alpha}}$. We shall return to possible physical aspects of this distinction in § 27.

In what follows we shall everywhere adopt definition (19.5) for definiteness.

Under the rotation subgroup the spinors ξ^α and $\eta_{\dot{\alpha}}$ transform, as we know, in the same way. Forming from their components the combinations

$$\xi^\alpha \pm \eta_{\dot{\alpha}}, \quad (19.6)$$

one would obtain quantities transforming under inversion according to (19.1) with $P = \pm i$. These combinations, however, do not behave as spinors with respect to all transformations of the Lorentz group.

Incorporating inversion into the symmetry group therefore demands that one treat a pair of spinors $(\xi^\alpha, \eta_{\dot{\alpha}})$ jointly; such a pair goes by the name of a first-rank *bispinor*. The four components of a bispinor realize one of the irreducible representations that belong to the extended Lorentz group.

Given two bispinors $(\xi^\alpha, \eta_{\dot{\alpha}})$ and $(\Xi^\alpha, H_{\dot{\alpha}})$, their scalar product admits two distinct constructions. The expression

$$\xi^\alpha \Xi_\alpha + \eta_{\dot{\alpha}} H^{\dot{\alpha}} \quad (19.7)$$

⁴If instead inversion is understood in the sense that $P^2 = -1$, its action is specified by the formulas

$$\xi^\alpha \rightarrow i\eta_{\dot{\alpha}}, \quad \eta_{\dot{\alpha}} \rightarrow i\xi^\alpha \quad (19.5)$$

or, equivalently,

$$\xi_\alpha \rightarrow -i\eta^{\dot{\alpha}}, \quad \eta^{\dot{\alpha}} \rightarrow -i\xi_\alpha. \quad (19.5a)$$

is altogether unchanged by inversion, i.e. it constitutes a true scalar. The quantity

$$\xi^\alpha \Xi_\alpha - \eta_{\dot{\alpha}} H^{\dot{\alpha}} \quad (19.8)$$

is likewise invariant under rotations of the 4-coordinate system but reverses sign under inversion; in other words, it is a pseudoscalar.

A second-rank spinor $\zeta^{\alpha\dot{\beta}}$ can also be defined in two ways. Specifying its transformation law as

$$\zeta^{\alpha\dot{\beta}} \sim \xi^\alpha H^{\dot{\beta}} + \Xi^\alpha \eta^{\dot{\beta}}, \quad (19.9)$$

one obtains quantities that transform under inversion according to

$$\zeta^{\alpha\dot{\beta}} \rightarrow \zeta_{\dot{\alpha}\beta}. \quad (19.10)$$

The 4-vector a^μ to which such a spinor is equivalent undergoes, by virtue of (18.1), the mapping $(a^0, \mathbf{a}) \rightarrow (a^0, -\mathbf{a})$, so it constitutes a genuine 4-vector (while the spatial part $\mathbf{a}$ is polar).

One may, however, also define $\zeta^{\alpha\dot{\beta}}$ according to

$$\zeta^{\alpha\dot{\beta}} \sim \xi^\alpha H^{\dot{\beta}} - \Xi^\alpha \eta^{\dot{\beta}}. \quad (19.11)$$

Then⁵

$$\zeta^{\alpha\dot{\beta}} \rightarrow -\zeta_{\dot{\alpha}\beta}. \quad (19.12)$$

The 4-vector corresponding to such a spinor undergoes the inversion transformation $(a^0, \mathbf{a}) \rightarrow (-a^0, \mathbf{a})$, i.e. it is a 4-pseudovector (while the three-dimensional vector $\mathbf{a}$ is axial).

Symmetric second-rank spinors whose indices are of the same type obey the transformation laws:

$$\xi^{\alpha\beta} \sim \xi^\alpha \Xi^\beta + \xi^\beta \Xi^\alpha, \quad \eta_{\dot{\alpha}\dot{\beta}} \sim \eta_{\dot{\alpha}} H_{\dot{\beta}} + \eta_{\dot{\beta}} H_{\dot{\alpha}}. \quad (19.13)$$

Under inversion they pass into one another:

$$\xi^{\alpha\beta} \rightarrow -\eta_{\dot{\alpha}\dot{\beta}}. \quad (19.14)$$

The pair $(\xi^{\alpha\beta}, \eta_{\dot{\alpha}\dot{\beta}})$ constitutes a second-rank bispinor. Its number of independent components equals $3 + 3 = 6$. An antisymmetric second-rank 4-tensor $a^{\mu\nu}$ possesses the same number of independent components. A definite correspondence must therefore exist between them (both furnish equivalent irreducible representations of the extended Lorentz group).

Since the spinors $\xi^{\alpha\beta}$ and $\eta_{\dot{\alpha}\dot{\beta}}$ transform independently under the proper Lorentz group, the components of the 4-tensor $a^{\mu\nu}$ can likewise be partitioned into two groups of quantities that mix only among themselves under all rotations of the 4-coordinate system. This decomposition is carried out as follows.

⁵We emphasize that the transformation laws (19.10) and (19.12), which differ by a sign on the right-hand side, are by no means equivalent, since both sides of each involve the components of one and the same spinor (cf. the footnote on p. 92).

We introduce a polar three-dimensional vector $\mathbf{p}$ and an axial three-dimensional vector $\mathbf{a}$, related to the components of the 4-tensor $a^{\mu\nu}$ by

$$a^{\mu\nu} = \begin{pmatrix} 0 & p_x & p_y & p_z \\ -p_x & 0 & -a_z & a_y \\ -p_y & a_z & 0 & -a_x \\ -p_z & -a_y & a_x & 0 \end{pmatrix} \equiv (\mathbf{p}, \mathbf{a}) \quad (19.15)$$

$(\mathbf{p}, \mathbf{a})$ is a shorthand notation that we shall employ for listing the components of such a tensor). Here $a^{\mu\nu} = (-\mathbf{p}, \mathbf{a})$, and of the two quantities

$$\mathbf{a}^2 - \mathbf{p}^2 = \frac{1}{2} a_{\mu\nu} a^{\mu\nu}, \quad \mathbf{a}\mathbf{p} = \frac{1}{8} e_{\mu\nu\rho\sigma} a^{\mu\nu} a^{\rho\sigma}$$

the first is a scalar and the second a pseudoscalar; with respect to the proper Lorentz group both are equally invariant. Equally invariant alongside them are the squared moduli of the three-dimensional vectors $\mathbf{f}^\pm = \mathbf{p} \pm i\mathbf{a}$. This means that every rotation in 4-space amounts, for the vectors $\mathbf{f}^\pm$, to a “rotation” in three-dimensional space—generally speaking through complex angles (the six rotation angles in 4-space correspond to three complex “rotation angles” of the three-dimensional system). The operation of spatial inversion, on the other hand, reverses the sign of $\mathbf{p}$ (but not of $\mathbf{a}$) and thereby interchanges the vectors $\mathbf{f}^+$ and $-\mathbf{f}^-$. The components of these vectors form precisely the two sought-after groups of quantities built from the components of the tensor $a^{\mu\nu}$.

At the same time the correspondence between the components of the 4-tensor $a^{\mu\nu}$ and those of the spinors $\xi^{\alpha\beta}, \eta_{\dot{\alpha}\dot{\beta}}$ becomes evident. Since spatial rotations enter the Lorentz group as a subgroup, the relations linking the spinor components to the components of the three-dimensional vectors must be the same as for three-dimensional spinors:

$$\begin{aligned} f^+_x &= \frac{1}{2}(\xi^{22} - \xi^{11}), & f^+_y &= \frac{i}{2}(\xi^{22} + \xi^{11}), & f^+_z &= \xi^{12}; \\ f^-_x &= \frac{1}{2}(\eta_{2\dot{2}} - \eta_{1\dot{1}}), & f^-_y &= \frac{i}{2}(\eta_{2\dot{2}} + \eta_{1\dot{1}}), & f^-_z &= \eta_{1\dot{2}}. \end{aligned} \quad (19.16)$$

Problem Derive the general correspondence that connects spinors of even rank to 4-tensors.

Solution. Every spinor whose total rank $k + l$ is even realizes a single-valued (though in general reducible) representation of the extended Lorentz group; it is consequently equivalent to a 4-tensor that realizes the same representation⁶.

A spinor of rank (k, k) transforms under inversion according to

$$\zeta^{\alpha\beta\dots\dot{\gamma}\dot{\delta}\dots} \rightarrow \pm \zeta_{\dot{\alpha}\dot{\beta}\dots\gamma\delta\dots}$$

Such a spinor is equivalent to a symmetric irreducible 4-tensor of rank k —a true tensor or a pseudotensor depending on the sign in (1).

Spinors of ranks (k, l) and (l, k) making up a bispinor transform under inversion according to

$$\underbrace{\zeta^{\alpha\beta\dots}}_k \underbrace{\dot{\gamma}\dot{\delta}\dots}_l \rightarrow (-1)^{\frac{k-l}{2}} \chi_{\dot{\alpha}\dot{\beta}\dots\gamma\delta\dots} \quad (2)$$

⁶Spinors whose rank is odd realize, by contrast, double-valued representations: a spatial rotation through a full turn of 360° reverses the sign of these spinors, so that each group element is associated with a pair of matrices differing in sign.

When $l = k + 2$ the bispinor is equivalent to an irreducible 4-tensor $a_{[\mu\nu]\rho\sigma\dots}$ of rank $k + 2$, antisymmetric in the indices $[\mu\nu]$ and symmetric in all the rest. The irreducibility of this tensor means that it vanishes upon contraction over any pair of indices and upon dualization with respect to any triple of indices (i.e. $e^{\lambda\mu\nu\sigma}a_{[\mu\nu]\rho\sigma\dots} = 0$); the latter condition signifies that the tensor vanishes upon antisymmetrization over any index triple.

More broadly, when $l = k + 2n$ the bispinor corresponds to an irreducible 4-tensor of rank $k + 2n$ that is antisymmetric in n index pairs $[\lambda\mu][\nu\rho]\dots$ and symmetric in the remaining k indices. Tensors possessing antisymmetry of yet higher degree (in triples, quadruples, and so on of indices) are absent from this classification for a transparent reason: a completely antisymmetric tensor of third rank is dual to a pseudovector, one of fourth rank collapses to a scalar (being proportional to the Levi-Civita pseudotensor $e^{\lambda\mu\nu\rho}$), and antisymmetry in more than four indices cannot arise in a four-dimensional space at all.

§ 20. The Dirac equation in spinor representation

In its own rest frame a particle of spin $1/2$ is described by a two-component wave function—a 3-spinor. From the “four-dimensional” standpoint this quantity may originate as either an undotted or a dotted 4-spinor. To describe the particle in an arbitrary frame of reference one needs both kinds of 4-spinor; let us label them ξ^α and $\eta_{\dot{\alpha}}$ ⁷.

In the case of a free particle the sole operator entering the wave equation can be (as was noted already in § 10) the 4-momentum $\hat{p}_\mu = i\partial_\mu$. Expressed through spinor indices, this 4-vector corresponds to an operator spinor $\hat{p}_{\alpha\dot{\beta}}$ with components

$$\begin{aligned}\hat{p}^{11} &= \hat{p}_{22} = \hat{p}_0 + \hat{p}_z, & \hat{p}^{22} &= \hat{p}_{11} = \hat{p}_0 - \hat{p}_z, \\ \hat{p}^{12} &= -\hat{p}_{21} = \hat{p}_x - i\hat{p}_y, & \hat{p}^{21} &= -\hat{p}_{12} = \hat{p}_x + i\hat{p}_y.\end{aligned}\quad (20.1)$$

The wave equation is a linear differential relation linking the components of the spinors through the operator $\hat{p}_{\alpha\dot{\beta}}$. The demand of relativistic invariance singles out the following system of equations:

$$\hat{p}^{\alpha\dot{\beta}}\eta_{\dot{\beta}} = m\xi^\alpha, \quad \hat{p}_{\dot{\beta}\alpha}\xi^\alpha = m\eta_{\dot{\beta}}, \quad (20.2)$$

where m is a dimensional constant. Introducing different constants m_1 and m_2 into these two equations (or reversing the sign in front of m) would be pointless, since by a suitable redefinition of ξ^α or $\eta_{\dot{\alpha}}$ the equations could always be brought back to the form given above.

Eliminating one of the two spinors from equations (20.2) by substituting $\eta_{\dot{\beta}}$ from the second equation into the first, we obtain:

$$\hat{p}^{\alpha\dot{\beta}}\eta_{\dot{\beta}} = \frac{1}{m}\hat{p}^{\alpha\dot{\beta}}\hat{p}_{\gamma\dot{\beta}}\xi^\gamma = m\xi^\alpha.$$

⁷A three-dimensional first-rank spinor can also “descend” from 4-spinors of higher odd rank that become antisymmetric in one or several index pairs in the rest frame. Such variants, however, would lead to equations of higher order (cf. the footnote on p. 52).

But by (18.4) $\widehat{p}^{\alpha\beta}\widehat{p}_{\gamma\beta} = \widehat{p}^2\delta_\gamma^\alpha$, so that we arrive at

$$(\widehat{p}^2 - m^2)\xi^\gamma = 0, \quad (20.3)$$

from which it is clear that m is the mass of the particle.

We draw attention to the fact that the necessity of introducing a mass into the wave equation requires the simultaneous treatment of a pair of spinors (ξ^α and $\eta_{\dot{\alpha}}$): from only one of them it is impossible to compose a relativistically invariant equation that would contain any dimensional parameter. By the same token the wave equation turns out to be automatically invariant with respect to spatial inversion, provided one defines the transformation of the wave function as

$$P : \quad \xi^\alpha \rightarrow i\eta_{\dot{\alpha}}, \quad \eta_{\dot{\alpha}} \rightarrow i\xi^\alpha. \quad (20.4)$$

One readily checks that this substitution (accompanied by the replacement $\widehat{p}^{\dot{\alpha}\beta} \rightarrow \widehat{p}_{\alpha\dot{\beta}}$, obvious from (20.1)) causes the two equations (20.2) to interchange. A pair of spinors that transform into one another under inversion make up a four-component object—a bispinor.

The relativistic wave equation embodied in the system (20.2) bears the name of the *Dirac equation* (it was derived by *Dirac* in 1928). To pursue the further study and application of this equation we shall now examine the different representations in which it can be written.

With the aid of formula (18.6) we rewrite equations (20.2) as

$$(\widehat{p}_0 + \widehat{\mathbf{p}}\boldsymbol{\sigma})\eta = m\xi, \quad (\widehat{p}_0 - \widehat{\mathbf{p}}\boldsymbol{\sigma})\xi = m\eta. \quad (20.5)$$

Here the symbols ξ and η stand for two-component quantities—spinors

$$\xi = \begin{pmatrix} \xi^1 \\ \xi^2 \end{pmatrix}, \quad \eta = \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix} \quad (20.6)$$

(the first carrying upper, the second lower indices), and when the matrices σ are multiplied by any two-component quantity f the ordinary rule of matrix multiplication is always understood here and below:

$$(\sigma f)_\alpha = \sigma_{\alpha\beta} f_\beta. \quad (20.7)$$

For ease of subsequent reference we write out the Pauli matrices once more:

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (20.8)$$

and recall their principal properties:

$$\sigma_i\sigma_k + \sigma_k\sigma_i = 2\delta_{ik}, \quad \sigma_i\sigma_k = ie_{ikl}\sigma_l + \delta_{ik} \quad (20.9)$$

(see III, § 55).

Let us also write down the wave equation obeyed by the complex-conjugate wave function, which is built from the spinors

$$\xi^* = (\xi^{1*}, \xi^{2*}), \quad \eta^* = (\eta_1^*, \eta_2^*). \quad (20.10)$$

Since every operator $\widehat{p}_\mu$ contains the factor i , one has $\widehat{p}_\mu^* = -\widehat{p}_\mu$. In taking the complex conjugate of both sides of equations (20.5) one must also take into account that, by the Hermiticity of the matrices σ ($\sigma^* = \widetilde{\sigma}$),

$$(\sigma f)_\alpha^* = \sigma_{\alpha\beta}^* f_\beta^* = f_\beta^* \sigma_{\beta\alpha} = (f^* \sigma)_\alpha,$$

and one obtains equations of the form

$$\eta^* (\widehat{p}_0 + \widehat{\mathbf{p}} \boldsymbol{\sigma}) = -m \xi^*, \quad \xi^* (\widehat{p}_0 - \widehat{\mathbf{p}} \boldsymbol{\sigma}) = -m \eta^*. \quad (20.11)$$

In this way of writing it is tacitly assumed that the operator $\widehat{p}_\mu$ acts on the function placed to its left. The display of ξ^* and η^* as horizontal rows reflects the matrix multiplication convention adopted in these equations: the row f is contracted with the columns of the σ matrices:

$$(f^* \sigma)_\alpha = f_\beta^* \sigma_{\beta\alpha}. \quad (20.12)$$

The inversion transformation for ξ^* , η^* is defined as the complex conjugate of the transformation (20.4):

$$P : \quad \xi^{\alpha*} \rightarrow -i \eta_{\dot{\alpha}}^*, \quad \eta_{\dot{\alpha}}^* \rightarrow -i \xi^{\alpha*}. \quad (20.13)$$

§ 21. Symmetric form of the Dirac equation

Among all ways of writing the Dirac equation the spinor form stands out as the one that displays relativistic invariance most transparently. For practical calculations, though, it is often advantageous to pass to a different representation by selecting another quartet of independent components of the wave function.

Let ψ stand for the four-component wave function (its entries being ψ_i , $i = 1, 2, 3, 4$). Written in spinor language this object is a bispinor:

$$\psi = \begin{pmatrix} \xi \\ \eta \end{pmatrix}. \quad (21.1)$$

With equal right, however, one may choose as the independent components of ψ any linear combinations of the components of the spinors ξ and η ⁸. We shall restrict the admissible linear transformations solely by the requirement of unitarity; such transformations leave invariant the bilinear forms composed of ψ and ψ^* (see § 28).

For a completely general selection of the components of ψ one can cast the Dirac equation as

$$\widehat{p}_\mu \gamma_{ik}^\mu \psi_k = m \psi_i,$$

where γ^μ ($\mu = 0, 1, 2, 3$) are certain 4×4 matrices (the *Dirac matrices*). It is customary to write this equation in symbolic form, suppressing the matrix indices:

$$(\gamma \widehat{p} - m) \psi = 0, \quad (21.2)$$

⁸For brevity we shall speak of the four-component quantity ψ as a bispinor also in its non-spinor representations.

where

$$\gamma \widehat{p} \equiv \gamma^\mu \widehat{p}_\mu = \widehat{p}_0 \gamma^0 - \widehat{\mathbf{p}} \boldsymbol{\gamma} = i\gamma^0 \frac{\partial}{\partial t} + i\boldsymbol{\gamma} \boldsymbol{\nabla}, \quad \boldsymbol{\gamma} = (\gamma^1, \gamma^2, \gamma^3).$$

In the spinor form of the equation with the components ψ from (21.1) the corresponding matrices are⁹

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \boldsymbol{\gamma} = \begin{pmatrix} 0 & -\boldsymbol{\sigma} \\ \boldsymbol{\sigma} & 0 \end{pmatrix}, \quad (21.3)$$

as is readily seen by writing equations (20.5) in the form

$$\begin{pmatrix} 0 & \widehat{p}_0 + \widehat{\mathbf{p}} \boldsymbol{\sigma} \\ \widehat{p}_0 - \widehat{\mathbf{p}} \boldsymbol{\sigma} & 0 \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix} = m \begin{pmatrix} \xi \\ \eta \end{pmatrix}$$

and comparing with (21.2).

In the general case the matrices γ need only satisfy the conditions ensuring that $\widehat{p}^2 = m^2$. To determine these conditions, we multiply equation (21.2) on the left by $\gamma \widehat{p}$. We have

$$(\gamma^\mu \widehat{p}_\mu) (\gamma^\nu \widehat{p}_\nu) \psi = m (\widehat{p}_\mu \gamma^\mu) \psi = m^2 \psi.$$

Since $\widehat{p}_\mu \widehat{p}_\nu$ is a symmetric tensor (all operators $\widehat{p}_\mu$ commute), this equation can be rewritten as

$$\frac{1}{2} \widehat{p}_\mu \widehat{p}_\nu (\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu) = m^2 \psi,$$

from which it is clear that one must have

$$\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2g^{\mu\nu}. \quad (21.4)$$

Thus all pairs of distinct matrices γ^μ anticommute, and the square of each is

$$(\gamma^1)^2 = (\gamma^2)^2 = (\gamma^3)^2 = -1, \quad (\gamma^0)^2 = 1. \quad (21.5)$$

Under an arbitrary unitary transformation of the components of ψ ($\psi' = U\psi$, with U a unitary 4×4 matrix) the matrices γ transform according to

$$\gamma' = U\gamma U^{-1} = U\gamma U^+ \quad (21.6)$$

(so that the equation $(\gamma \widehat{p} - m)\psi = 0$ goes over into $(\gamma' \widehat{p} - m)\psi' = 0$). The commutation relations (21.4), of course, remain unchanged.

The matrix γ^0 of (21.3) is Hermitian, and the matrices $\boldsymbol{\gamma}$ are anti-Hermitian. These properties are preserved under every unitary transformation (21.6), so that we shall always have¹⁰:

$$\gamma^+ = -\boldsymbol{\gamma}, \quad \gamma^{0+} = \gamma^0. \quad (21.7)$$

⁹Here and below a compact notation is used for 4×4 matrices in terms of 2×2 blocks: each symbol in the expressions (21.3) represents a 2×2 matrix.

¹⁰These relations can be written together as

$$\gamma^{\lambda+} = \gamma^0 \gamma^\lambda \gamma^0. \quad (21.7 \text{ a})$$

We now derive the equation satisfied by ψ^* . Complex-conjugating both sides of (21.2) and invoking (21.7) gives

$$(\widehat{p}_0\gamma^0 - \widehat{\mathbf{p}}\widetilde{\boldsymbol{\gamma}} - m)\psi^* = 0.$$

We rearrange ψ^* by $\widetilde{\gamma}^\mu\psi^* = \psi^*\gamma^\mu$ and then multiply the equation on the right by γ^0 ; noting that $\boldsymbol{\gamma}\gamma^0 = -\gamma^0\boldsymbol{\gamma}$ and introducing the new bispinor

$$\bar{\psi} = \psi^*\gamma^0, \quad \psi^* = \bar{\psi}\gamma^0, \quad (21.8)$$

we obtain

$$\bar{\psi}(\gamma\widehat{p} + m) = 0. \quad (21.9)$$

Just as in (20.11), $\widehat{p}$ is understood to differentiate the function standing to its left. One calls $\bar{\psi}$ the *Dirac conjugate* (or *relativistic conjugate*) of ψ . The purpose of the γ^0 factor is to swap ξ^* and η^* (in the spinor representation), so that within $\bar{\psi} = (\eta^*, \xi^*)$ the undotted spinor occupies the first position—matching the ordering in ψ —while the dotted spinor occupies the second. This is why $\bar{\psi}$, rather than ψ^* , serves as the natural companion of ψ whenever the two enter bilinear combinations together (see § 28).

Under spatial inversion the wave function transforms according to

$$P: \quad \psi \rightarrow i\gamma^0\psi, \quad \bar{\psi} \rightarrow -i\bar{\psi}\gamma^0. \quad (21.10)$$

In the spinor representation of ψ the matrix γ^0 interchanges, as it should under inversion, the components ξ and η . The invariance of the Dirac equation under the transformation (21.10) is in the general case obvious and can be verified directly: replacing $\widehat{\mathbf{p}} \rightarrow -\widehat{\mathbf{p}}$ and simultaneously $\psi \rightarrow i\gamma^0\psi$ in equation (21.2), we get

$$(\widehat{p}_0\gamma^0 + \widehat{\mathbf{p}}\boldsymbol{\gamma} - m)\gamma^0\psi = 0.$$

Multiplying this equation on the left by γ^0 and using the anticommutativity of γ^0 and $\boldsymbol{\gamma}$, we recover the original equation.

Left-multiplying $(\gamma\widehat{p} - m)\psi = 0$ by $\bar{\psi}$, right-multiplying $\bar{\psi}(\gamma\widehat{p} + m) = 0$ by ψ , and summing the two expressions yields

$$\bar{\psi}\gamma^\mu(\widehat{p}_\mu\psi) + (\widehat{p}_\mu\bar{\psi})\gamma^\mu\psi = \widehat{p}_\mu(\bar{\psi}\gamma^\mu\psi) = 0,$$

where parentheses mark the function acted upon by $\widehat{p}$. Because this has the structure of a continuity equation $\partial_\mu j^\mu = 0$, the quantity

$$j^\mu = \bar{\psi}\gamma^\mu\psi = (\psi^*\psi, \psi^*\gamma^0\boldsymbol{\gamma}\psi) \quad (21.11)$$

is the particle current density 4-vector. We note that its time component $j^0 = \psi^*\psi$ is positive definite.

The Dirac equation can be written in a form explicitly solved for the time derivative:

$$i\frac{\partial\psi}{\partial t} = \widehat{H}\psi, \quad (21.12)$$

where $\widehat{H}$ is the particle Hamiltonian¹¹. To this end it suffices to multiply equation (21.2) on the left by γ^0 . For the Hamiltonian one obtains the expression

$$\widehat{H} = \alpha \widehat{\mathbf{p}} + \beta m, \quad (21.13)$$

where the standard notation has been introduced for the matrices appearing here:

$$\alpha = \gamma^0 \boldsymbol{\gamma}, \quad \beta = \gamma^0. \quad (21.14)$$

We note that

$$\alpha_i \alpha_k + \alpha_k \alpha_i = 2\delta_{ik}, \quad \beta \alpha + \alpha \beta = 0, \quad \beta^2 = 1, \quad (21.15)$$

i.e. all the matrices α, β anticommute with one another and their squares are unity; all of them are Hermitian. In the spinor representation

$$\alpha = \begin{pmatrix} \boldsymbol{\sigma} & 0 \\ 0 & -\boldsymbol{\sigma} \end{pmatrix}, \quad \beta = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (21.16)$$

When velocities are small the particle should reduce, as in non-relativistic quantum mechanics, to a single two-component spinor. And indeed, taking $\mathbf{p} \rightarrow 0$, $\varepsilon \rightarrow m$ in (20.5) yields $\xi = \eta$: the two spinors inside the bispinor coincide. A drawback of the spinor representation shows up at this point, however—all four entries of ψ stay non-vanishing despite only two being genuinely independent. One would prefer a representation in which two of the four components of ψ go to zero in the non-relativistic limit.

Accordingly we introduce, in place of ξ and η , their linear combinations φ and χ :

$$\psi = \begin{pmatrix} \varphi \\ \chi \end{pmatrix}, \quad \varphi = \frac{1}{\sqrt{2}}(\xi + \eta), \quad \chi = \frac{1}{\sqrt{2}}(\xi - \eta). \quad (21.17)$$

Then for a particle at rest $\chi = 0$. This representation of ψ will be called the *standard* representation. Under inversion φ and χ transform into themselves:

$$P : \quad \varphi \rightarrow i\varphi, \quad \chi \rightarrow -i\chi. \quad (21.18)$$

The equations for φ and χ are obtained by adding and subtracting equations (20.5):

$$\widehat{p}_0 \varphi - \widehat{\mathbf{p}} \boldsymbol{\sigma} \chi = m\varphi, \quad -\widehat{p}_0 \chi + \widehat{\mathbf{p}} \boldsymbol{\sigma} \varphi = m\chi. \quad (21.19)$$

From these one sees that the standard representation corresponds to the matrices

$$\gamma^0 \equiv \beta = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \boldsymbol{\gamma} = \begin{pmatrix} 0 & \boldsymbol{\sigma} \\ -\boldsymbol{\sigma} & 0 \end{pmatrix}, \quad \alpha = \begin{pmatrix} 0 & \boldsymbol{\sigma} \\ \boldsymbol{\sigma} & 0 \end{pmatrix}. \quad (21.20)$$

Since in (21.17) the first and second components of ξ and η are combined separately, in the standard representation—as in the spinor one—the components ψ_1 and ψ_3 correspond to

¹¹For a particle of spin 0 the wave equation could not be cast in this form: equation (10.5) for the scalar ψ is of second order in time, while the system (10.4) of first-order equations for the five-component quantity (ψ, ψ_k) contains time derivatives not of all of its components.

the eigenvalue $+1/2$ of the spin projection, while ψ_2 and ψ_4 correspond to the projection $-1/2$. In both representations, therefore, the matrix $\frac{1}{2}\Sigma$ with

$$\Sigma = \begin{pmatrix} \sigma & 0 \\ 0 & \sigma \end{pmatrix}, \quad (21.21)$$

is the three-dimensional spin operator: the action of $\frac{1}{2}\Sigma_z$ on a bispinor containing only the components ψ_1, ψ_3 or ψ_2, ψ_4 multiplies the bispinor by $+1/2$ or $-1/2$, respectively. In an arbitrary representation this matrix can be written as

$$\Sigma = -\alpha\gamma^5 = -\frac{i}{2}[\mathbf{a}\mathbf{a}] \quad (21.22)$$

(for the definition of γ^5 see below, (22.14)).

Problems

1. Find the transformation formulae for the wave function under an infinitesimal Lorentz transformation and an infinitesimal spatial rotation.

Solution. In the spinor representation of ψ , under an infinitesimal Lorentz transformation

$$\xi' = \left(1 - \frac{1}{2}\sigma\delta\mathbf{V}\right)\xi, \quad \eta' = \left(1 + \frac{1}{2}\sigma\delta\mathbf{V}\right)\eta$$

(see (18.8), (18.8 a), (18.12)). Both formulae can be written together as

$$\psi' = \left(1 - \frac{1}{2}\alpha\delta\mathbf{V}\right)\psi. \quad (1)$$

Similarly, the transformation law under an infinitesimal rotation is

$$\psi' = \left(1 + \frac{i}{2}\Sigma\delta\theta\right)\psi. \quad (2)$$

In this form the formulae are valid in any representation of ψ , provided one understands α and Σ as the matrices in the same representation.

It is easy to verify that the matrices α and Σ constitute the components of the antisymmetric “matrix 4-tensor”

$$\sigma^{\mu\nu} = \frac{1}{2}(\gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu) = (\alpha, i\Sigma)$$

(the listing of the components follows the rule (19.15)). Let us also introduce the infinitesimal antisymmetric tensor $\delta\varepsilon^{\mu\nu} = (\delta\mathbf{V}, \delta\theta)$. Then

$$\sigma^{\mu\nu}\delta\varepsilon_{\mu\nu} = 2i\Sigma\delta\theta - 2\gamma\delta\mathbf{V}$$

and both formulae (1), (2) can be written in the unified form:

$$\psi' = \left(1 + \frac{1}{4}\sigma^{\mu\nu}\delta\varepsilon_{\mu\nu}\right)\psi. \quad (3)$$

2. Write the Dirac equation in a representation that contains no imaginary coefficients (*E. Majorana, 1937*).

Solution. In the standard representation, in the equation

$$\left(\frac{\partial}{\partial t} + \alpha_x \frac{\partial}{\partial x} + \alpha_y \frac{\partial}{\partial y} + \alpha_z \frac{\partial}{\partial z} + im\beta \right) \psi = 0$$

the only imaginary matrices are α_y and $i\beta$. This imaginary character can be eliminated by performing a transformation $\psi' = U\psi$ that interchanges the imaginary matrix α_y with the real matrix β . To this end one sets

$$U = \frac{1}{\sqrt{2}}(\alpha_y + \beta) = U^{-1}.$$

Then

$$\alpha'_x = U\alpha_x U = -\alpha_x, \quad \alpha'_y = \beta, \quad \alpha'_z = -\alpha_z, \quad \beta' = \alpha_y,$$

and the Dirac equation takes the form

$$\left(\frac{\partial}{\partial t} - \alpha_x \frac{\partial}{\partial x} + \beta \frac{\partial}{\partial y} - \alpha_z \frac{\partial}{\partial z} + im\alpha_y \right) \psi' = 0,$$

in which all the coefficients are real.

§ 22. Algebra of Dirac matrices

Calculations involving the Dirac equation make extensive use of the γ matrices without reference to their explicit form in any particular representation. All the algebraic rules governing these matrices follow entirely from the anticommutation relations

$$\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2g^{\mu\nu}, \quad \mu, \nu = 0, 1, 2, 3, \quad (22.1)$$

which encode every general property that the γ matrices possess.

In this section we collect a number of formulae and algebraic identities for the γ matrices that prove useful in various computations.

The “scalar square” of γ with itself is $g_{\mu\nu} \gamma^\mu \gamma^\nu = 4$. For compact notation we introduce, in analogy with the covariant components of a 4-vector, the symbol $\gamma_\mu = g_{\mu\nu} \gamma^\nu$. Then

$$\gamma_\mu \gamma^\mu = 4. \quad (22.2)$$

When γ_μ and γ^μ are separated by one or more γ -factors, one can bring them next to each other by applying the commutation rule (22.1) one or more times, after which the contraction over μ is performed using (22.2). In this way one obtains the identities

$$\begin{aligned} \gamma_\mu \gamma^\nu \gamma^\mu &= -2\gamma^\nu, \\ \gamma_\mu \gamma^\lambda \gamma^\nu \gamma^\mu &= 4g^{\lambda\nu}, \\ \gamma_\mu \gamma^\lambda \gamma^\nu \gamma^\rho \gamma^\mu &= -2\gamma^\rho \gamma^\nu \gamma^\lambda, \\ \gamma_\mu \gamma^\lambda \gamma^\nu \gamma^\rho \gamma^\sigma \gamma^\mu &= 2(\gamma^\sigma \gamma^\lambda \gamma^\nu \gamma^\rho + \gamma^\rho \gamma^\nu \gamma^\lambda \gamma^\sigma). \end{aligned} \quad (22.3)$$

The γ^μ factors usually appear contracted with various 4-vectors, forming “scalar products”¹²

$$\gamma a \equiv \gamma^\mu a_\mu. \quad (22.4)$$

For products of this kind the relations (22.1) become

$$(a\gamma)(b\gamma) + (b\gamma)(a\gamma) = 2(ab), \quad (a\gamma)(a\gamma) = a^2 \quad (22.5)$$

and the identities (22.3) take the form

$$\begin{aligned} \gamma_\mu(a\gamma)\gamma^\mu &= -2(a\gamma), \\ \gamma_\mu(a\gamma)(b\gamma)\gamma^\mu &= 4(ab), \\ \gamma_\mu(a\gamma)(b\gamma)(c\gamma)\gamma^\mu &= -2(c\gamma)(b\gamma)(a\gamma), \\ \gamma_\mu(a\gamma)(b\gamma)(c\gamma)(d\gamma)\gamma^\mu &= 2[(d\gamma)(a\gamma)(b\gamma)(c\gamma) + (c\gamma)(b\gamma)(a\gamma)(d\gamma)]. \end{aligned} \quad (22.6)$$

A widely used operation is the evaluation of the trace of a product of several γ matrices. Consider the quantities

$$T^{\mu_1\mu_2\ldots\mu_n} \equiv \frac{1}{4} \text{Sp}(\gamma^{\mu_1}\gamma^{\mu_2}\ldots\gamma^{\mu_n}). \quad (22.7)$$

By the well-known cyclic property of the matrix trace, the tensor T is symmetric under cyclic permutations of the indices $\mu_1\mu_2\ldots\mu_n$.

Because the γ matrices take the same form in every reference frame, the quantities T are likewise frame-independent. They therefore constitute a tensor expressible solely through the metric tensor $g_{\mu\nu}$, which is the only tensor sharing this property.

From the rank-2 tensor $g_{\mu\nu}$ one can build only tensors of even rank. It follows at once that the trace of any product containing an odd number of γ factors vanishes. In particular, the trace of each individual γ is zero¹³:

$$\text{Sp} \gamma^\mu = 0. \quad (22.8)$$

The trace of the 4×4 unit matrix (which is implicitly present on the right-hand side of the anticommutation relation (22.1)) equals 4. Taking the trace of both sides of (22.1) therefore gives

$$T^{\mu\nu} = g^{\mu\nu}. \quad (22.9)$$

The trace of a product of four matrices is

$$T^{\lambda\mu\nu\rho} = g^{\lambda\mu}g^{\nu\rho} - g^{\lambda\nu}g^{\mu\rho} + g^{\lambda\rho}g^{\mu\nu}. \quad (22.10)$$

This formula can be derived, for instance, by “dragging” γ^λ to the right through $\text{Sp}(\gamma^\lambda\gamma^\mu\gamma^\nu\gamma^\rho)$ with the help of the anticommutation relation (22.1); each commutation produces one of the terms appearing in (22.10):

$$\begin{aligned} T^{\lambda\mu\nu\rho} &= 2g^{\lambda\mu}T^{\nu\rho} - T^{\mu\lambda\nu\rho} = 2g^{\lambda\mu}g^{\nu\rho} - T^{\mu\lambda\nu\rho}, \\ T^{\mu\lambda\nu\rho} &= 2g^{\lambda\nu}T^{\mu\rho} - T^{\mu\nu\lambda\rho} = 2g^{\lambda\nu}g^{\mu\rho} - T^{\mu\nu\lambda\rho}, \\ T^{\mu\nu\lambda\rho} &= 2g^{\lambda\rho}T^{\mu\nu} - T^{\mu\nu\rho\lambda} \end{aligned} \quad (22.11)$$

¹²In this edition we do not use any special notation for such a product. The literature frequently employs letters with a hat or with a slash (Feynman slash notation).

¹³The trace is invariant under similarity transformations $\gamma = U\gamma U^{-1}$. Hence (22.8) is also obvious from the explicit matrices (21.3).

and so on. After all permutations the right-hand side contains $-T^{\mu\rho\lambda} = -T^{\lambda\mu\nu\rho}$, which is brought to the left. By the same method the trace of a product of six γ 's reduces to traces of products of four, etc. Thus,

$$T^{\lambda\mu\nu\rho\sigma\tau} = g^{\lambda\mu}T^{\nu\rho\sigma\tau} - g^{\lambda\nu}T^{\mu\rho\sigma\tau} + g^{\lambda\rho}T^{\mu\nu\sigma\tau} - \\ - g^{\lambda\sigma}T^{\mu\nu\rho\tau} + g^{\lambda\tau}T^{\mu\nu\rho\sigma}. \quad (22.11)$$

We remark that every trace $T^{\lambda\mu\dots}$ is real, and that it is non-vanishing only when each of the matrices $\gamma^0, \gamma^1, \dots$

occurs an even number of times in the product; both statements are plain from the formulae just obtained. From this one easily deduces that the trace is unchanged when the order of all factors is reversed:

$$T^{\lambda\mu\dots\rho\sigma} = T^{\sigma\rho\dots\mu\lambda}. \quad (22.12)$$

As already noted, the γ factors usually appear as scalar products with various 4-vectors. In that case formulae (22.9) and (22.10), for example, read

$$\frac{1}{4} \text{Sp}(a\gamma)(b\gamma) = ab, \\ \frac{1}{4} \text{Sp}(a\gamma)(b\gamma)(c\gamma)(d\gamma) = (ab)(cd) - (ac)(bd) + (ad)(bc). \quad (22.13)$$

A special role is played by the product $\gamma^0\gamma^1\gamma^2\gamma^3$, for which the standard notation is

$$\gamma^5 = -i\gamma^0\gamma^1\gamma^2\gamma^3. \quad (22.14)$$

One readily verifies that

$$\gamma^5\gamma^\mu + \gamma^\mu\gamma^5 = 0, \quad (\gamma^5)^2 = 1, \quad (22.15)$$

so that γ^5 anticommutes with every γ^μ . With respect to the matrices α and β the rules are

$$\alpha\gamma^5 - \gamma^5\alpha = 0, \quad \beta\gamma^5 + \gamma^5\beta = 0 \quad (22.16)$$

(the commutativity with α follows from the fact that $\alpha = \gamma^0\boldsymbol{\gamma}$ is a product of two γ^μ matrices).

The matrix γ^5 is Hermitian; indeed,

$$\gamma^{5+} = i\gamma^{3+}\gamma^{2+}\gamma^{1+}\gamma^{0+} = -i\gamma^3\gamma^2\gamma^1\gamma^0,$$

and because the sequence 3210 is obtained from 0123 by an even number of transpositions,

$$\gamma^{5+} = \gamma^5. \quad (22.17)$$

We also give the explicit form of this matrix in the two concrete representations:

$$\begin{array}{ll} \text{spinor} & \gamma^5 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}; \\ \text{standard} & \gamma^5 = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}. \end{array} \quad (22.18)$$

The trace of γ^5 vanishes:

$$\text{Sp } \gamma^5 = 0 \quad (22.19)$$

(as is also directly apparent from (22.18)). Likewise zero are the traces of $\gamma^5 \gamma^\mu \gamma^\nu$. For the product of γ^5 with four γ^μ factors, however, one has

$$\frac{1}{4} \text{Sp } \gamma^5 \gamma^\lambda \gamma^\mu \gamma^\nu \gamma^\rho = i e^{\lambda\mu\nu\rho}. \quad (22.20)$$

We note one further identity:

$$\gamma N = i \gamma^5 (\gamma a)(\gamma b)(\gamma c), \quad N^\lambda = e^{\lambda\mu\nu\rho} a_\mu b_\nu c_\rho, \quad (22.21)$$

valid for mutually orthogonal 4-vectors a, b, c : $ab = ac = bc = 0$.

There are situations (problems with non-relativistic particles) where one must compute traces of products containing γ^0 and the spatial “vector” $\boldsymbol{\gamma}$ as separate entities. A non-vanishing trace requires that γ^0 as well as $\boldsymbol{\gamma}$ each enter an even number of times. Every γ^0 factor then simplifies to the identity, and for products carrying two or four $\boldsymbol{\gamma}$ factors the traces read

$$\begin{aligned} \frac{1}{4} \text{Sp}(\mathbf{a}\boldsymbol{\gamma})(\mathbf{b}\boldsymbol{\gamma}) &= -\mathbf{a}\mathbf{b}, \\ \frac{1}{4} \text{Sp}(\mathbf{a}\boldsymbol{\gamma})(\mathbf{b}\boldsymbol{\gamma})(\mathbf{c}\boldsymbol{\gamma})(\mathbf{d}\boldsymbol{\gamma}) &= (\mathbf{a}\mathbf{b})(\mathbf{c}\mathbf{d}) - (\mathbf{a}\mathbf{c})(\mathbf{b}\mathbf{d}) + (\mathbf{a}\mathbf{d})(\mathbf{b}\mathbf{c}). \end{aligned} \quad (22.22)$$

§ 23. Plane waves

A free particle with definite momentum and energy is described by a plane wave, which we write as

$$\psi_p = \frac{1}{\sqrt{2\varepsilon}} u_p e^{-ipx}. \quad (23.1)$$

The subscript p labels the 4-momentum; the wave amplitude u_p is a bispinor normalised in a specific way.

When carrying out the second quantisation later on, we shall need, alongside the wave functions (23.1), also the “negative-frequency” solutions that arise in relativistic theory, as explained in § 11, because of the two-valued square root $\pm\sqrt{\mathbf{p}^2 + m^2}$. Following the convention of § 11 we shall everywhere take ε to be the positive quantity $\varepsilon = +\sqrt{\mathbf{p}^2 + m^2}$, so that the “negative frequency” is $-\varepsilon$; reversing also the sign of $\mathbf{p}$ we arrive at a function that is naturally denoted ψ_{-p} :

$$\psi_{-p} = \frac{1}{\sqrt{2\varepsilon}} u_{-p} e^{ipx}. \quad (23.2)$$

The physical meaning of these functions will become clear in § 26. Below we shall write formulae for ψ_p and ψ_{-p} side by side.

The components of the bispinor amplitudes u_p and u_{-p} satisfy the algebraic systems

$$(\gamma p - m)u_p = 0, \quad (\gamma p + m)u_{-p} = 0, \quad (23.3)$$

obtained by substituting (23.1), (23.2) into the Dirac equation

(which amounts to replacing the operator $\widehat{p}$ by $\pm p$)¹⁴. The relation $p^2 = m^2$ serves here as the consistency condition for each system. We shall always normalise the bispinor amplitudes by the invariant conditions

$$\bar{u}_p u_p = 2m, \quad \bar{u}_{-p} u_{-p} = -2m \quad (23.4)$$

(the bar denotes, as throughout, Dirac conjugation: $\bar{u} = u^* \gamma^0$). Left-multiplying equations (23.3) by $\bar{u}_{\pm p}$ gives $(\bar{u}_{\pm p} \gamma u_{\pm p})p = 2m^2 = 2p^2$, whence

$$\bar{u}_p \gamma u_p = \bar{u}_{-p} \gamma u_{-p} = 2p. \quad (23.5)$$

We note that the formulae for u_{-p} are obtained from those for u_p by changing the sign of m .

The current-density 4-vector is

$$j = \bar{\psi}_{\pm p} \gamma \psi_{\pm p} = \frac{1}{2\varepsilon} \bar{u}_{\pm p} \gamma u_{\pm p} = \frac{p}{\varepsilon}, \quad (23.6)$$

i.e. $j^\mu = (1, \mathbf{v})$, where $\mathbf{v} = \mathbf{p}/\varepsilon$ is the particle velocity. This shows that the functions ψ_p are normalised to “one particle per unit volume $V = 1$ ”.

Because of equations (23.3) the components of the wave amplitude are related to one another; the explicit form of these relations depends, of course, on the choice of representation for ψ . Let us find them for the standard representation.

From equations (21.19) we have, for a plane wave,

$$(\varepsilon - m)\varphi - \mathbf{p}\boldsymbol{\sigma}\chi = 0, \quad (\varepsilon + m)\chi - \mathbf{p}\boldsymbol{\sigma}\varphi = 0. \quad (23.7)$$

These relations yield the connection between φ and χ in two equivalent forms:

$$\varphi = \frac{\mathbf{p}\boldsymbol{\sigma}}{\varepsilon - m}\chi, \quad \chi = \frac{\mathbf{p}\boldsymbol{\sigma}}{\varepsilon + m}\varphi \quad (23.8)$$

(the equivalence is plain: multiplying the first relation on the left by $\mathbf{p}\boldsymbol{\sigma}/(\varepsilon + m)$ and using $(\mathbf{p}\boldsymbol{\sigma})^2 = \mathbf{p}^2$, $\varepsilon^2 - m^2 = \mathbf{p}^2$ reproduces the second). The overall factor in φ and χ is chosen so as to satisfy the normalisation condition (23.4). The result for u_p (and likewise for u_{-p}) is

$$u_p = \begin{pmatrix} \sqrt{\varepsilon + m} w \\ \frac{\mathbf{p}\boldsymbol{\sigma}}{\sqrt{\varepsilon + m}} w \end{pmatrix}, \quad u_{-p} = \begin{pmatrix} \frac{-\mathbf{p}\boldsymbol{\sigma}}{\sqrt{\varepsilon + m}} w' \\ \sqrt{\varepsilon + m} w' \end{pmatrix}, \quad (23.9)$$

(the second expression is obtained from the first by reversing the sign of m and relabelling $w \rightarrow (\mathbf{n}\boldsymbol{\sigma})w'$). Here $\mathbf{n}$ is the unit vector along $\mathbf{p}$, and w is an arbitrary two-component quantity subject only to the normalisation

$$w^* w = 1. \quad (23.10)$$

¹⁴We note also the analogous systems that follow from the Dirac equation (21.9) for the complex-conjugate function:

$$\bar{u}_p (\gamma p - m) = 0, \quad \bar{u}_{-p} (\gamma p + m) = 0. \quad (23.3 \text{ a})$$

For $\bar{u} = u^* \gamma^0$ (with γ^0 from (21.20)) we have

$$\begin{aligned}\bar{u}_p &= (\sqrt{\varepsilon + m} w^*, -\sqrt{\varepsilon - m} w^* (\mathbf{n}\sigma)), \\ \bar{u}_{-p} &= (\sqrt{\varepsilon - m} w'^* (\mathbf{n}\sigma), -\sqrt{\varepsilon + m} w'^*),\end{aligned}\quad (23.11)$$

and direct multiplication confirms that $\bar{u}_{\pm p} u_{\pm p} = \pm 2m$.

In the rest frame, i.e. when $\varepsilon = m$, we have

$$u_p = \sqrt{2m} \begin{pmatrix} w \\ 0 \end{pmatrix}, \quad u_{-p} = \sqrt{2m} \begin{pmatrix} 0 \\ w' \end{pmatrix}, \quad (23.12)$$

i.e. w is precisely the 3-spinor that each wave amplitude becomes in the slow-motion limit. Observe that for the bispinor u_{-p} the components that vanish at rest are the upper two, not the lower two. This behaviour of the negative-frequency Dirac solutions is easy to understand: putting $\mathbf{p} = 0$ and $\varepsilon \rightarrow -m$ in (23.7) yields $\varphi = 0$ ¹⁵.

The plane-wave amplitude contains one arbitrary two-component quantity. In other words, for a given momentum there exist two distinct independent states, corresponding to the two possible values of the spin projection. The projection of the spin on an arbitrary axis z cannot, however, possess a definite value. This is because the Hamiltonian of a particle with definite $\mathbf{p}$ (i.e. the matrix $H = \alpha \mathbf{p} + \beta m$) does not commute with $\Sigma_z = -i\alpha_x \alpha_y$. In agreement with the general statements of § 16, however, the helicity λ —the projection of the spin onto the direction of $\mathbf{p}$ —is conserved: the Hamiltonian commutes with $\mathbf{n}\Sigma$.

Helicity eigenstates correspond to plane waves whose 3-spinor $w = w^{(\lambda)}(\mathbf{n})$ is an eigenfunction of the operator $\mathbf{n}\sigma$:

$$\frac{1}{2}(\mathbf{n}\sigma)w^{(\lambda)} = \lambda w^{(\lambda)}. \quad (23.13)$$

The explicit form of these spinors is

$$w^{(\lambda=1/2)} = \begin{pmatrix} e^{-i\varphi/2} \cos \frac{\theta}{2} \\ e^{i\varphi/2} \sin \frac{\theta}{2} \end{pmatrix}, \quad w^{(\lambda=-1/2)} = \begin{pmatrix} -e^{-i\varphi/2} \sin \frac{\theta}{2} \\ e^{i\varphi/2} \cos \frac{\theta}{2} \end{pmatrix}, \quad (23.14)$$

where θ and φ are the polar angle and azimuth of the direction $\mathbf{n}$ with respect to the fixed axes xyz ¹⁶.

An alternative choice of two independent states of a free particle with given $\mathbf{p}$ (simpler, though less transparent physically) corresponds to the two values of the z -projection of the spin in the rest frame; we denote it by σ . The associated spinors are

$$w^{(\sigma=1/2)} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad w^{(\sigma=-1/2)} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (23.15)$$

As the two linearly independent negative-frequency solutions we choose plane waves with 3-spinors

$$w^{(\sigma)'} = -\sigma_y w^{(-\sigma)} = 2\sigma i w^{(\sigma)} \quad (23.16)$$

¹⁵In the spinor representation one finds $\xi = -\eta$ rather than the relation $\xi = \eta$ that holds at rest for positive-frequency solutions.

¹⁶The solution of equations (23.13) admits multiplication by an arbitrary phase factor, reflecting the freedom of an arbitrary rotation about $\mathbf{n}$.

(the rationale behind this choice will become clear in § 26).

One can construct a representation of the plane wave in which, in every frame (not only in the rest frame), it has just two non-vanishing components that correspond to definite values of the same physical observable—the spin projection in the rest frame (*L. Foldy, S. A. Wouthuysen, 1950*).

Taking u_p in the standard representation (23.9) as the starting point, we look for a unitary map into such a representation, writing

$$u'_p = U u_p, \quad U = e^{W\gamma\mathbf{n}},$$

where W is real; since $\gamma^+ = -\gamma$ the unitarity $U^+ = U^{-1}$ is automatic. Expanding in a series and noting that $(\gamma\mathbf{n})^2 = -1$, we can write U as

$$U = \cos W + \gamma\mathbf{n} \sin W$$

(cf. the passage from (18.13) to (18.14)). Requiring that the last two components of the transformed amplitude u'_p vanish, we find

$$\operatorname{tg} W = \frac{|\mathbf{p}|}{m + \varepsilon},$$

so that

$$U = \frac{m + \varepsilon + (\gamma\mathbf{n})|\mathbf{p}|}{\sqrt{2\varepsilon(\varepsilon + m)}}.$$

In the new representation

$$u'_p = \sqrt{2\varepsilon} \begin{pmatrix} w \\ 0 \end{pmatrix}. \quad (23.17)$$

The particle Hamiltonian in this representation becomes

$$\hat{H}' = U(\alpha\mathbf{p} + \beta m)U^{-1} = \beta\varepsilon \quad (23.18)$$

(here β, α, γ are the standard-representation matrices). This Hamiltonian commutes with

$$\Sigma = -\alpha\gamma^5 = \begin{pmatrix} \sigma & 0 \\ 0 & \sigma \end{pmatrix},$$

which, in the Foldy–Wouthuysen representation, plays the role of the operator for the conserved rest-frame spin.

§ 24. Spherical waves

The wave functions of a free spin-1/2 particle with definite values of the total angular momentum j are spinor spherical waves. We shall determine their form, first recalling the analogous expressions from non-relativistic theory.

The non-relativistic wave function is a 3-spinor $\psi = \begin{pmatrix} \psi^1 \\ \psi^2 \end{pmatrix}$. For a state with definite energy ε (and hence definite magnitude of the momentum p ¹⁷), orbital angular momentum l , total angular momentum j , and projection m , it takes the form

$$\psi = R_{pl}(r)\Omega_{jlm}(\theta, \varphi). \quad (24.1)$$

¹⁷Throughout this section p denotes $|\mathbf{p}|$.

The angular factor Ω_{jlm} is a 3-spinor whose components (for the two values $j = l \pm 1/2$ possible at a given l) are

$$\Omega_{l+1/2, l, m} = \begin{pmatrix} \sqrt{\frac{j+m}{2j}} Y_{l, m-1/2} \\ \sqrt{\frac{j-m}{2j}} Y_{l, m+1/2} \end{pmatrix}, \quad (24.2)$$

$$\Omega_{l-1/2, l, m} = \begin{pmatrix} -\sqrt{\frac{j-m+1}{2j+2}} Y_{l, m-1/2} \\ \sqrt{\frac{j+m+1}{2j+2}} Y_{l, m+1/2} \end{pmatrix}$$

(see III, § 106, problem). We shall call Ω_{jlm} the *spherical spinors*. Their normalisation is

$$\int \Omega_{jlm}^* \Omega_{j'l'm'} d\omega = \delta_{jj'} \delta_{ll'} \delta_{mm'}. \quad (24.3)$$

The radial functions R_{pl} are a common factor in both spinor components of ψ and are given by the formula (33.10) (see III):

$$R_{pl} = \sqrt{\frac{2\pi p}{r}} J_{l+1/2}(pr). \quad (24.4)$$

They are normalised according to

$$\int_0^\infty r^2 R_{p'l} R_{pl} dr = 2\pi \delta(p' - p). \quad (24.5)$$

Turning now to the relativistic case, we recall first of all that for a moving particle no separate conservation laws for spin and orbital angular momentum exist: the operators $\widehat{\mathbf{S}}$ and $\widehat{\mathbf{L}}$ individually fail to commute with the Hamiltonian. The parity of the state, however, remains conserved (for a free particle). Consequently the quantum number l ceases to specify a definite value of the orbital angular momentum, but it does determine (see below) the parity of the state.

We shall work with the desired wave function (a bispinor) in the standard representation: $\psi = \begin{pmatrix} \varphi \\ \chi \end{pmatrix}$. Under rotations φ and χ transform as 3-spinors, so their angular dependence is again carried by the spherical spinors Ω_{jlm} . Let $\varphi \propto \Omega_{jlm}$, where l is one (definite) of the two values $j + 1/2$ or $j - 1/2$. Under inversion $\varphi(\mathbf{r}) \rightarrow i\varphi(-\mathbf{r})$ (see (21.18)), and since $\Omega_{jlm}(-\mathbf{n}) = (-1)^l \Omega_{jlm}(\mathbf{n})$,

$$\varphi(\mathbf{r}) \rightarrow i(-1)^l \varphi(\mathbf{r}).$$

The lower component χ , on the other hand, transforms under inversion as $\chi(\mathbf{r}) \rightarrow -i\chi(-\mathbf{r})$. For the state to possess a definite parity (i.e. for inversion to multiply every component by the same phase), the angular dependence of χ must involve $\Omega_{j'l'm}$ with the other value of l : because the two allowed values differ by unity, $(-1)^{l'} = -(-1)^l$.

Furthermore, the radial dependence of φ and χ is governed by the same functions R_{pl} and $R_{p'l'}$ (with l and l' matching the order of the spherical harmonics entering Ω_{jlm}).

This follows from the fact that each component of ψ satisfies the second-order equation $(\hat{p}^2 - m^2)\psi = 0$, which for a given $|\mathbf{p}|$ reads

$$(\Delta + \mathbf{p}^2)\psi = 0,$$

coinciding in form with the free-particle Schrödinger equation of the non-relativistic theory.

Thus

$$\varphi = AR_{pl}\Omega_{jlm}, \quad \chi = BR_{pl'}\Omega_{jl'm}, \quad (24.6)$$

and it remains to fix the constant coefficients A and B . To this end we examine the far zone, where the spherical wave may be treated as a plane wave. By the asymptotic formula (33.12) (see III)

$$R_{pl} \approx \frac{1}{ir} \left\{ e^{i\left(pr - \frac{\pi l}{2}\right)} - e^{-i\left(pr - \frac{\pi l}{2}\right)} \right\}, \quad (24.7)$$

so that φ is a superposition of two plane waves propagating along $\pm \mathbf{n}$ ($\mathbf{n} = \mathbf{r}/r$). For each of these, by (23.8),

$$\chi = \frac{p}{\varepsilon + m}(\pm \mathbf{n}\boldsymbol{\sigma})\varphi.$$

From what has been said above (formulae (24.6)) it is clear that $(\mathbf{n}\boldsymbol{\sigma})\Omega_{jlm} = a\Omega_{jl'm}$ with a a constant. This constant is easily found by evaluating both sides at $m = 1/2$ with $\mathbf{n}$ along the z -axis. Using (7.2a) one obtains

$$(\mathbf{n}\boldsymbol{\sigma})\Omega_{jlm} = i^{l'-l}\Omega_{jl'm}. \quad (24.8)$$

Collecting the formulae and comparing with (24.6), we find

$$B = -\frac{p}{\varepsilon + m} A.$$

Finally, the coefficient A is fixed by the overall normalisation of ψ . Adopting the condition

$$\int \psi_{pjlm}^* \psi_{p'j'l'm'} d^3x = 2\pi \delta_{jj'} \delta_{ll'} \delta_{mm'} \delta(p - p'), \quad (24.9)$$

we arrive at the final result

$$\psi_{pjlm} = \frac{1}{\sqrt{2\varepsilon}} \begin{pmatrix} \sqrt{\varepsilon + m} R_{pl} \Omega_{jlm} \\ -\sqrt{\varepsilon - m} R_{pl'} \Omega_{jl'm} \end{pmatrix}, \quad l' = 2j - l. \quad (24.10)$$

At fixed j and m (and energy ε) there are therefore two states that differ in parity. Which parity a state carries is uniquely determined by l ($= j \pm 1/2$): inversion multiplies the bispinor (24.10) by $i(-1)^l$. The upper and lower components of this bispinor do, however, involve spherical harmonics of both orders l and l' —a direct manifestation of the fact that the orbital angular momentum is not separately well-defined.

As $r \rightarrow \infty$ one can, in any small region of space, regard the spherical waves (24.7) as plane waves with momentum $\mathbf{p} = \pm p\mathbf{n}$. It is therefore clear that the momentum-space wave functions differ from (24.10) mainly in the

absence of radial factors, with $\mathbf{n}$ now denoting the direction of the momentum.

To pass directly to the momentum representation one performs the Fourier transform:

$$\psi(\mathbf{p}') = \int \psi(\mathbf{r}) e^{-i\mathbf{p}'\mathbf{r}} d^3x, \quad (24.11)$$

The integral is evaluated with the aid of the plane-wave expansion in spherical harmonics (see III, (34.3)):

$$e^{i\mathbf{p}\mathbf{r}} = \frac{2\pi}{p} \sum_{l=0}^{\infty} \sum_{m=-l}^l i^l R_{pl}(r) Y_{lm}^*\left(\frac{\mathbf{p}}{p}\right) Y_{lm}\left(\frac{\mathbf{r}}{r}\right). \quad (24.12)$$

Substituting $e^{-i\mathbf{p}'\mathbf{r}}$ into (24.11) via this expansion and using (24.5), the Fourier components of the function

$$\psi(\mathbf{r}) = R_{pl}(r) \Omega_{jlm}\left(\frac{\mathbf{r}}{r}\right)$$

are found to be

$$\psi(\mathbf{p}') = \frac{(2\pi)^2}{p} \delta(p' - p) i^{-l} Y_{lm'}\left(\frac{\mathbf{p}'}{p'}\right) \int \Omega_{jlm}\left(\frac{\mathbf{r}}{r}\right) Y_{lm'}^*\left(\frac{\mathbf{r}}{r}\right) d\Omega.$$

The integral here equals the coefficients of the spherical harmonics in the definition of the spherical spinors (24.2), and together with the factor $Y_{lm'}(\mathbf{p}'/p')$ it reconstructs the same spherical spinor, now depending on $\mathbf{p}'/p'$:

$$\psi(\mathbf{p}') = \frac{(2\pi)^2}{p} \delta(p' - p) i^{-l} \Omega_{jlm}\left(\frac{\mathbf{p}'}{p'}\right).$$

Applying this result to the bispinor wave function (24.10), we obtain its momentum-space form

$$\psi_{pjlm}(\mathbf{p}') = \delta(p' - p) \frac{(2\pi)^2}{p\sqrt{2\varepsilon}} \begin{pmatrix} \sqrt{\varepsilon + m} i^{-l} \Omega_{jlm}(\mathbf{p}'/p') \\ -\sqrt{\varepsilon - m} i^{-l'} \Omega_{jl'm}(\mathbf{p}'/p') \end{pmatrix}. \quad (24.13)$$

The states $|pjlm\rangle$ coincide with the states $|pjm|\lambda\rangle$ considered in § 16 (with $|\lambda| = 1/2$): both sets possess definite values of pjm and parity. The spherical spinors Ω_{jlm} are therefore expressible through the functions $D_m^{(j)}$ (both depending on the argument $\mathbf{p}/p$). In the limit $p \rightarrow 0$ the wave functions (24.13) reduce to 3-spinors Ω_{jlm} with parity $P = \eta(-1)^l$ (where $\eta = i$ is the “intrinsic parity” of the spinor). Comparison with the results of § 16 leads to the formula

$$\Omega_{jlm} = i^l \sqrt{\frac{2j+1}{8\pi}} \left(w^{(-1/2)} D_{-1/2,m}^{(j)} \pm w^{(1/2)} D_{1/2,m}^{(j)} \right) \quad (24.14)$$

(with $l = j \mp 1/2$, where $w^{(\lambda)}$ are the 3-spinors (23.14)).

§ 25. The connection between spin and statistics

Second quantisation of the field of spin-1/2 particles (the spinor field) proceeds in exactly the same way as was done in § 11 for the scalar field.

Without repeating the entire argument, we write down at once the expressions for the field operators, fully analogous to (11.2):

$$\begin{aligned}\hat{\psi} &= \sum_{\mathbf{p}\sigma} \frac{1}{\sqrt{2\varepsilon}} (\hat{a}_{\mathbf{p}\sigma} u_{\mathbf{p}\sigma} e^{-i\mathbf{p}x} + \hat{b}_{\mathbf{p}\sigma}^+ u_{-\mathbf{p}-\sigma} e^{i\mathbf{p}x}), \\ \hat{\bar{\psi}} &\equiv \hat{\psi}^+ \gamma^0 = \sum_{\mathbf{p}\sigma} \frac{1}{\sqrt{2\varepsilon}} (\hat{a}_{\mathbf{p}\sigma}^+ \bar{u}_{\mathbf{p}\sigma} e^{i\mathbf{p}x} + \hat{b}_{\mathbf{p}\sigma} \bar{u}_{-\mathbf{p}-\sigma} e^{-i\mathbf{p}x});\end{aligned}\quad (25.1)$$

the summation runs over all values of $\mathbf{p}$ and over $\sigma = \pm 1/2$. The antiparticle annihilation operators $\hat{b}_{\mathbf{p}\sigma}$ (like the particle annihilation operators $\hat{a}_{\mathbf{p}\sigma}$) stand as coefficients of functions whose coordinate dependence ($e^{i\mathbf{p}x}$) corresponds to a state with momentum $\mathbf{p}$ ¹⁸.

To construct the Hamiltonian of the spinor field there is no need to determine its energy–momentum tensor (as we did for the scalar field), because here the single-particle Hamiltonian exists and can be used to write down the wave equation (the Dirac equation) (21.12). The mean energy of a particle in a state with wave function ψ is the integral

$$\int \psi^* \hat{H} \psi d^3x = \int \psi^* \frac{\partial \psi}{\partial t} d^3x = i \int \bar{\psi} \gamma^0 \frac{\partial \psi}{\partial t} d^3x. \quad (25.2)$$

We draw attention to the fact that the “energy density” (the integrand) is not positive definite here.

Replacing ψ and $\bar{\psi}$ in (25.2) by ψ -operators, taking into account the mutual orthogonality of wave functions with different $\mathbf{p}$ or σ , and also the relation $\bar{u}_{\pm\mathbf{p}\sigma} \gamma^0 u_{\pm\mathbf{p}\sigma} = 2\varepsilon$ for the wave amplitudes, we obtain the field Hamiltonian

$$\hat{H} = \sum_{\mathbf{p}\sigma} \varepsilon (\hat{a}_{\mathbf{p}\sigma}^+ \hat{a}_{\mathbf{p}\sigma} - \hat{b}_{\mathbf{p}\sigma} \hat{b}_{\mathbf{p}\sigma}^+). \quad (25.3)$$

It follows that quantisation must here be carried out according to Fermi statistics:

$$\{\hat{a}_{\mathbf{p}\sigma}, \hat{a}_{\mathbf{p}\sigma}^+\}_+ = 1, \quad \{\hat{b}_{\mathbf{p}\sigma}, \hat{b}_{\mathbf{p}\sigma}^+\}_+ = 1, \quad (25.4)$$

while all other pairs of operators $\hat{a}, \hat{a}^+, \hat{b}, \hat{b}^+$ anticommute (see III, § 65). Indeed, the Hamiltonian (25.3) then takes the form

$$\hat{H} = \sum_{\mathbf{p}\sigma} \varepsilon (\hat{a}_{\mathbf{p}\sigma}^+ \hat{a}_{\mathbf{p}\sigma} + \hat{b}_{\mathbf{p}\sigma}^+ \hat{b}_{\mathbf{p}\sigma} - 1),$$

and the energy eigenvalues (after subtracting, as always, the infinite additive constant) are

$$E = \sum_{\mathbf{p}\sigma} \varepsilon (N_{\mathbf{p}\sigma} + \bar{N}_{\mathbf{p}\sigma}), \quad (25.5)$$

which are, as required, positive definite. Had we quantised according to Bose statistics, equation (25.3) would have given the meaningless eigenvalues

$$\sum_{\mathbf{p}\sigma} \varepsilon (N_{\mathbf{p}\sigma} - \bar{N}_{\mathbf{p}\sigma}),$$

¹⁸Both types of function also correspond to the same value of σ , the rest-frame spin projection: for the functions $\bar{\psi}_{-\mathbf{p}, -\sigma}$ this will be demonstrated in § 26, see (26.10).

which are not positive definite.

The analogous expression

$$\mathbf{P} = \sum_{\mathbf{p}\sigma} \mathbf{p}(N_{\mathbf{p}\sigma} + \bar{N}_{\mathbf{p}\sigma}) \quad (25.6)$$

is obtained for the momentum of the system—the eigenvalues of the operator $\int \hat{\psi}^+ \hat{\mathbf{p}} \hat{\psi} d^3x$.

The 4-current operator is

$$\hat{j}^\mu = \hat{\psi} \gamma^\mu \hat{\psi}, \quad (25.7)$$

and for the “charge” operator of the field we get

$$\hat{Q} = \int \hat{\psi} \gamma^0 \hat{\psi} d^3x = \sum_{\mathbf{p}\sigma} (\hat{a}_{\mathbf{p}\sigma}^+ \hat{a}_{\mathbf{p}\sigma} + \hat{b}_{\mathbf{p}\sigma} \hat{b}_{\mathbf{p}\sigma}^+) = \sum_{\mathbf{p}\sigma} (\hat{a}_{\mathbf{p}\sigma}^+ \hat{a}_{\mathbf{p}\sigma} - \hat{b}_{\mathbf{p}\sigma}^+ \hat{b}_{\mathbf{p}\sigma} + 1), \quad (25.8)$$

with eigenvalues

$$Q = \sum_{\mathbf{p}\sigma} (N_{\mathbf{p}\sigma} - \bar{N}_{\mathbf{p}\sigma}). \quad (25.9)$$

We thus arrive once more at the picture of particles and antiparticles; everything said about them in § 11 applies here as well.

Whereas spin-0 particles are bosons, however, spin-1/2 particles turn out to be fermions. Tracing this difference to its formal origin, one finds that it stems from a difference in the character of the “energy density” for the scalar and spinor fields. In the scalar case the energy density is positive definite, so that both terms in the Hamiltonian (11.3)

($\hat{a}^+ \hat{a}$ and $\hat{b} \hat{b}^+$) enter with a plus sign. To guarantee positive energy eigenvalues, the replacement of $\hat{b} \hat{b}^+$ by $\hat{b}^+ \hat{b}$ must then proceed without a change of sign, i.e. by the Bose commutation rule. For the spinor field, on the other hand, the energy density is not positive definite, so that the $\hat{b} \hat{b}^+$ term in the Hamiltonian (25.3) carries a minus sign; obtaining positive eigenvalues now requires that the rearrangement $\hat{b} \hat{b}^+ \rightarrow \hat{b}^+ \hat{b}$ be accompanied by a sign change, i.e. that it follow the Fermi commutation rule.

On the other hand, the form of the energy density is directly tied to the transformation properties of the wave function and to the demands of relativistic invariance. One can therefore say that the connection between a particle’s spin and the statistics it obeys is itself a direct consequence of these requirements.

From the fact that spin-1/2 particles are fermions one can also draw a general conclusion: every particle with half-integer spin is a fermion, and every particle with integer spin is a boson (the latter includes the spin-0 result proved in § 11)¹⁹.

This becomes obvious once one notes that a particle of spin s may be thought of as “assembled” from $2s$ particles of spin 1/2. When s is half-integer the number $2s$ is odd, while for integer s it is even. Now a composite object built from an even count

¹⁹The origin of the connection between the spin of a particle and the statistics it obeys was elucidated by Pauli (W. Pauli, 1940).

of fermions behaves as a boson, whereas one containing an odd count of fermions is a fermion²⁰.

When a system contains particles of several species, each species must be equipped with its own creation and annihilation operators. Operators belonging to different bosons, or to a boson and a fermion, commute with one another. As for operators belonging to different fermions, in non-relativistic theory they could be taken either commuting or anticommuting (III, § 65). In a relativistic theory, however, which allows mutual transformations of particles, the creation and annihilation operators of distinct fermion species must be regarded as anticommuting—just

like operators referring to different states of the same fermion.

Problem

Find the Lagrangian of the spinor field.

Solution. A real, Lorentz-scalar Lagrangian that yields the Dirac equation is

$$L = \frac{i}{2} (\bar{\psi} \gamma^\mu \partial_\mu \psi - \partial_\mu \bar{\psi} \cdot \gamma^\mu \psi) - m \bar{\psi} \psi. \quad (1)$$

Treating the components of ψ and $\bar{\psi}$ as “generalised coordinates” q , one easily verifies that the corresponding Euler–Lagrange equations (10.10) coincide with the Dirac equations for $\bar{\psi}$ and ψ . The overall sign of the Lagrangian (as well as the overall numerical coefficient) is a matter of convention here. Because L is linear in the derivatives of ψ and $\bar{\psi}$, the action $S = \int L d^4x$ can possess neither a minimum nor a maximum. The condition $\delta S = 0$ determines in this case only a stationary point, not an extremum, of the integral.

The field-theoretic Lagrangian of the spinor field is obtained by replacing ψ in (1) with the operator $\hat{\psi}$. Applying to this Lagrangian the formula (12.12), one recovers the current operator (25.7).

§ 26. Charge conjugation and time reversal of spinors

The factors $\psi_{p\sigma} = u_{p\sigma} e^{-ipx}$ that multiply the operators $\hat{a}_{p\sigma}$ in (25.1) are the wave functions of free particles (which we shall call “electrons”) carrying momentum $\mathbf{p}$ and polarisation σ :

$$\psi_{p\sigma}^{(e)} = \psi_{p\sigma}.$$

The factors $\bar{\psi}_{-p-\sigma}$ accompanying the operators $\hat{b}_{p\sigma}$, on the other hand, should be regarded as positron wave functions for the same $\mathbf{p}$ and σ . It turns out, however, that the electron and positron functions are expressed in different bispinor representations. This is apparent from the fact that ψ and $\bar{\psi}$ have distinct transformation properties, their components obeying different systems of equations. To eliminate this drawback one must carry out a specific unitary transformation of the components of $\bar{\psi}_{-p-\sigma}$ —one that

²⁰This reasoning tacitly assumes that all particles of the same spin obey the same statistics, regardless of how they are “assembled”. That this is indeed so can be seen from an analogous argument. If spin-0 fermions existed, then combining a spin-0 fermion with a spin-1/2 fermion would produce a spin-1/2 particle that is a boson—in contradiction with the general result established above for spin 1/2.

ensures the resulting four-component function satisfies the same equation as $\psi_{p\sigma}$ ²¹. It is precisely such a function that we shall call the positron wave function (with momentum $\mathbf{p}$ and polarisation σ). Denoting the matrix of the required unitary transformation by U_C , we write

$$\psi_{p\sigma}^{(p)} = U_C \bar{\psi}_{-p-\sigma}. \quad (26.1)$$

The operation C by which this function is constructed from $\psi_{-p-\sigma}$ is called the *charge conjugation* of the wave function (*H. A. Kramers*, 1937). This notion is by no means limited to plane waves. For any function ψ whatsoever there exists a “charge-conjugate” function

$$\widehat{C}\psi(t, \mathbf{r}) = U_C \bar{\psi}(t, \mathbf{r}), \quad (26.2)$$

which transforms in the same way as ψ and obeys the same equation.

The properties of U_C follow from this definition. If ψ solves the Dirac equation $(\gamma\widehat{p} - m)\psi = 0$, then $\bar{\psi}$ satisfies

$$\bar{\psi}(\gamma\widehat{p} + m) = 0, \quad \text{i.e.} \quad (\bar{\gamma}\widehat{p} + m)\bar{\psi} = 0.$$

Multiplying this equation on the left by U_C :

$$U_C \bar{\gamma}\widehat{p}\bar{\psi} + mU_C \bar{\psi} = 0,$$

and demanding that $U_C \bar{\psi}$ obey the same equation as ψ :

$$(\gamma\widehat{p} - m)U_C \bar{\psi} = 0,$$

we compare the two equations and obtain the following “commutation relation” between U_C and the matrices γ^μ ⁻¹:

$$U_C \bar{\gamma}^\mu = -\gamma^\mu U_C. \quad (26.3)$$

We shall henceforth assume that the wave functions are given in the spinor or standard representation (the case of an arbitrary representation will be discussed only at the end of this section). In these representations

$$\begin{aligned} \gamma^{0,2} &= \bar{\gamma}^{0,2}, & \gamma^{1,3} &= -\bar{\gamma}^{1,3}, \\ (\gamma^{0,1,3})^* &= \gamma^{0,1,3}, & \gamma^{2*} &= -\gamma^2. \end{aligned} \quad (26.4)$$

The conditions (26.3) are then met by the matrix $U_C = \eta_C \gamma^2 \gamma^0$ with an arbitrary constant η_C . The requirement $\widehat{C}^2 = 1$ implies $|\eta_C|^2 = 1$, so U_C is fixed up to a phase factor. We shall set $\eta_C = 1$, giving

$$U_C = \gamma^2 \gamma^0 = -\alpha_y. \quad (26.5)$$

²¹For spin-0 particles this question did not arise, since the scalar functions ψ and ψ^* both satisfy the same equation, and ψ_{-p}^* simply coincides with ψ_p .

Noting further that $\bar{\psi} = \psi^* \gamma^0 = \bar{\gamma}^0 \psi^* = \gamma^0 \psi^*$, one can write the action of $\widehat{C}$ as

$$\widehat{C}\psi = \gamma^2 \gamma^0 \bar{\psi} = \gamma^2 \psi^*. \quad (26.6)$$

Written out explicitly, the transformation (26.6) reads in the spinor representation

$$\begin{aligned} C : \quad \xi^\alpha &\rightarrow -i\eta^{\alpha*}, & \eta_\alpha &\rightarrow -i\xi_\alpha^*, \\ C : \quad \xi_\alpha &\rightarrow i\xi^{\alpha*}, & \eta^\alpha &\rightarrow -i\eta_\alpha^*. \end{aligned} \quad (26.7a, b)$$

The charge-conjugation transformation for the plane waves $\psi_{\pm p\sigma}$ is easily carried out using their explicit forms (23.9) together with the matrix U_C in the standard representation:

$$U_C = \begin{pmatrix} 0 & -\sigma_y \\ -\sigma_y & 0 \end{pmatrix}. \quad (26.8)$$

Noting that

$$\sigma_y \sigma^* = -\sigma \sigma_y,$$

and using the definition of $w^{(\sigma)'} from (23.16), we obtain$

$$U_C \bar{u}_{-p-\sigma} = u_{p\sigma}, \quad U_C u_{-p-\sigma} = \bar{u}_{p\sigma}. \quad (26.9)$$

Consequently,

$$\widehat{C}\psi_{-p-\sigma} = \psi_{p\sigma}, \quad (26.10)$$

confirming that the functions $\psi_{-p-\sigma}$ appearing in the ψ -operators (25.1) alongside $\widehat{b}_{p\sigma}$ do indeed describe states of a particle with momentum $\mathbf{p}$ and polarisation σ . We also see that the electron and positron states are described by the same functions:

$$\psi_{p\sigma}^{(e)} = \psi_{p\sigma}^{(p)} = \psi_{p\sigma}.$$

This is entirely natural, since the functions $\psi_{p\sigma}$ encode only the momentum and polarisation of the particle.

The operation of time reversal can be treated in an analogous manner. Reversing the sign of time must be accompanied by complex conjugation of the wave function. In order for the resulting “time-reversed” wave function ($\widehat{T}\psi$) to belong to the same representation as the original ψ , one must

perform a further unitary transformation on the components of ψ^* (or $\bar{\psi}$). Proceeding in analogy with (26.2), we represent the action of $\widehat{T}$ on ψ as

$$\widehat{T}\psi(t, \mathbf{r}) = U_T \bar{\psi}(-t, \mathbf{r}), \quad (26.11)$$

where U_T is a unitary matrix.

Writing once more the Dirac equation for ψ :

$$\left(i\gamma^0 \frac{\partial}{\partial t} + i\boldsymbol{\gamma} \nabla - m \right) \psi(t, \mathbf{r}) = 0,$$

and the corresponding equation for $\bar{\psi}$:

$$\left(i\bar{\gamma}^0 \frac{\partial}{\partial t} + i\bar{\boldsymbol{\gamma}} \nabla + m \right) \bar{\psi}(t, \mathbf{r}) = 0.$$

Replacing $t \rightarrow -t$ in the latter and multiplying on the left by $-U_T$:

$$\left(iU_T\tilde{\gamma}^0\frac{\partial}{\partial t} - iU_T\tilde{\gamma}\nabla\right)\tilde{\psi}(-t, \mathbf{r}) - mU_T\tilde{\psi}(-t, \mathbf{r}) = 0.$$

We require $U_T\tilde{\psi}(-t, \mathbf{r})$ to satisfy the same equation as $\psi(t, \mathbf{r})$:

$$\left(i\gamma^0\frac{\partial}{\partial t} + i\gamma\nabla\right)U_T\tilde{\psi}(-t, \mathbf{r}) - mU_T\tilde{\psi}(-t, \mathbf{r}) = 0.$$

Comparison yields the conditions

$$U_T\tilde{\gamma}^0 = \gamma^0 U_T, \quad U_T\tilde{\gamma} = -\gamma U_T. \quad (26.12)$$

In the spinor and standard representations, these conditions are satisfied by the matrix²²

$$U_T = i\gamma^3\gamma^1\gamma^0. \quad (26.13)$$

The action of $\hat{T}$ is therefore given by

$$\hat{T}\psi(t, \mathbf{r}) = i\gamma^3\gamma^1\gamma^0\tilde{\psi}(-t, \mathbf{r}) = i\gamma^3\gamma^1\psi^*(-t, \mathbf{r}). \quad (26.14)$$

Explicitly, in the spinor representation this transformation reads

$$T : \quad \xi^\alpha \rightarrow -i\xi_\alpha^*, \quad \eta_{\dot{\alpha}} \rightarrow i\eta^{\dot{\alpha}*} \quad (26.15 \text{ a})$$

or

$$T : \quad \xi_\alpha \rightarrow i\xi^{\alpha*}, \quad \eta^\alpha \rightarrow -i\eta_{\dot{\alpha}}^*. \quad (26.15 \text{ b})$$

In the standard representation

$$U_T = \begin{pmatrix} \sigma_y & 0 \\ 0 & -\sigma_y \end{pmatrix}. \quad (26.16)$$

Let us now find the result of applying all three operations P , T , and C to ψ . Writing them out in succession:

$$\begin{aligned} \hat{T}\psi(t, \mathbf{r}) &= -i\gamma^1\gamma^3\psi^*(-t, \mathbf{r}), \\ \hat{P}\hat{T}\psi(t, \mathbf{r}) &= i\gamma^0(\hat{T}\psi) = \gamma^0\gamma^1\gamma^3\psi^*(-t, -\mathbf{r}), \\ \hat{C}\hat{P}\hat{T}\psi(t, \mathbf{r}) &= \gamma^2(\gamma^0\gamma^1\gamma^3\psi^*)^* = \gamma^2\gamma^0\gamma^1\gamma^3\psi(-t, -\mathbf{r}), \end{aligned}$$

which gives

$$\hat{C}\hat{P}\hat{T}\psi(t, \mathbf{r}) = i\gamma^5\psi(-t, -\mathbf{r}). \quad (26.17)$$

In the spinor representation

$$CPT : \quad \xi^\alpha \rightarrow -i\xi_\alpha^*, \quad \eta_{\dot{\alpha}} \rightarrow i\eta^{\dot{\alpha}*}, \quad (26.18)$$

²²The choice of phase factor in (26.13) is tied to the choice made in (26.5) by considerations discussed below in the footnote on p. 124.

as can also be verified directly from the transformation rules (20.4), (26.7), (26.15)²³.

The expressions for the matrices U_C and U_T written above assumed the spinor or standard representation of ψ . Let us finally clarify which properties of these expressions remain valid for an arbitrary representation.

If ψ undergoes a unitary transformation:

$$\psi' = U\psi, \quad \gamma' = U\gamma U^{-1}, \quad \bar{\psi}' = \psi'^* \gamma'^0 = \bar{\psi} U^+ = \bar{\psi} \tilde{U}^{-1}, \quad (26.19)$$

then in the new representation

$$(\hat{C}\psi)' = U(\hat{C}\psi) = UU_C\tilde{\psi} = UU_C(\widetilde{\psi'U}) = UU_C\tilde{U}\tilde{\psi}'.$$

Comparing with the definition of U'_C in the new representation ($(\hat{C}\psi)' = U'_C\tilde{\psi}'$), we find

$$U'_C = UU_C\tilde{U}. \quad (26.20)$$

The transformation (26.20) coincides with the transformation law of the γ matrices only when U is real. For this reason the expression (26.5) holds only in representations obtainable from the spinor or standard representation by a real transformation.

The matrix (26.5) is unitary, and transposition reverses its sign:

$$U_C U_C^+ = 1, \quad \tilde{U}_C = -U_C. \quad (26.21)$$

These properties are invariant under the transformation (26.20) and hence hold in every representation. The matrix (26.5) is also Hermitian ($U_C = U_C^+$), but this property is in general not preserved by (26.20).

Everything just said (including (26.21)) applies equally to the properties of the matrix U_T .

Within the second-quantisation formalism, the transformations C , P , T for ψ -operators must be formulated as transformation rules for the creation and annihilation operators of the particles. These rules can be established (similarly to how it was done in § 13 for spin-0 particles) by requiring that the transformed ψ -operators admit a representation of the form

$$\begin{aligned} \hat{\psi}^C(t, \mathbf{r}) &= U_C \tilde{\hat{\psi}}(t, \mathbf{r}), \\ \hat{\psi}^P(t, \mathbf{r}) &= i\gamma^0 \hat{\psi}(t, -\mathbf{r}), \\ \hat{\psi}^T(t, \mathbf{r}) &= U_T \tilde{\hat{\psi}}(-t, \mathbf{r}). \end{aligned} \quad (26.22)$$

Problem

Find the charge-conjugation operator in the Majorana representation (see Problem 2, § 21).

Solution. The matrix U'_C in the Majorana representation is obtained from $U_C = -\alpha_y$ in the standard representation via the transformation (26.20) with $U = (\alpha_y + \beta)/\sqrt{2}$,

²³The notation $\hat{C}\hat{P}\hat{T}$ implies that the operators act from right to left. The overall sign in (26.17), (26.18) depends on this ordering because $\hat{T}$ does not commute with $\hat{C}$ and $\hat{P}$ (in their action on a bispinor).

and equals $U'_C = \alpha_y$ (α_y and β denote the standard-representation matrices). Denoting quantities in the Majorana representation by a prime, we have $\widehat{C}\psi' = U'_C(\psi'^*\beta')$, and since $\beta' = \alpha_y$,

$$\widehat{C}\psi' = \alpha_y(\psi'^*\alpha_y) = \alpha_y\widetilde{\alpha}_y\psi'^* = \psi'^*,$$

i.e. charge conjugation reduces to complex conjugation.

§ 27. Internal symmetry of particles and antiparticles

In its rest frame the wave function of a spin-1/2 particle reduces to a single 3-spinor (which we denote Φ^α). The behaviour of this spinor under inversion is linked to the notion of the particle's intrinsic parity. However (as already noted in § 19), although the two possible transformation laws for 3-spinors ($\Phi^\alpha \rightarrow \pm i\Phi^\alpha$) are inequivalent, assigning a definite parity to a spinor has no absolute meaning. It is therefore meaningless to speak of the intrinsic parity of a spin-1/2 particle by itself. One can, however, speak of the *relative* intrinsic parity of two such particles.

From two (three-dimensional) spinors $\Phi^{(e)}$ and $\Phi^{(p)}$ one can form the scalar $\Phi^{(e)}_\alpha \Phi^{(p)\alpha}$. If this quantity is a true scalar, the particles described by these spinors are said to have the same parity; if it is a pseudoscalar, their intrinsic parities are said to be opposite.

We shall show that the intrinsic parities of a particle and its antiparticle (with spin 1/2) are opposite (V. B. Berestetskii, 1948).

To this end, note that if the operation C (26.7) is applied to both sides of the P -transformation (19.5) (in the spinor representation)

$$P : \quad \xi^\alpha \rightarrow i\eta_{\dot{\alpha}}, \quad \eta_{\dot{\alpha}} \rightarrow i\xi^\alpha, \quad (27.1)$$

one obtains

$$\eta^{c\alpha*} \rightarrow i\xi_{\dot{\alpha}}^{c*}, \quad \xi_{\dot{\alpha}}^{c*} \rightarrow i\eta^{c\alpha*},$$

where the superscript c labels the components of the bispinor $\psi^C = \begin{pmatrix} \xi^c \\ \eta^c \end{pmatrix}$ that is charge-conjugate to $\psi = \begin{pmatrix} \xi \\ \eta \end{pmatrix}$. Taking the complex conjugate and rearranging the indices, we find

$$P : \quad \eta_{\dot{\alpha}}^c \rightarrow i\xi^{c\alpha}, \quad \xi^{c\alpha} \rightarrow i\eta_{\dot{\alpha}}^c. \quad (27.2)$$

We see that charge-conjugate bispinors transform under inversion according to the same law.

Let $\psi^{(e)}$ be the wave function of the particle (electron) and $\psi^{(p)}$ that of the antiparticle (positron). The latter is a bispinor obtained by charge-conjugating a certain “negative-frequency” solution of the Dirac equation. In the rest frame each reduces to a 3-spinor:

$$\xi^{(e)\alpha} = \eta_{\dot{\alpha}}^{(e)} = \Phi^{(e)\alpha}, \quad \xi^{(p)\alpha} = \eta_{\dot{\alpha}}^{(p)} = \Phi^{(p)\alpha}.$$

According to (27.1) and (27.2), these spinors transform under inversion as

$$\Phi^\alpha \rightarrow i\Phi^\alpha, \quad (27.3)$$

with the same rule for both $\Phi^{(e)}$ and $\Phi^{(p)}$. The product $\Phi^{(e)}\Phi^{(p)}$, however, changes sign—which proves the assertion.

A particle that coincides with its own antiparticle is called truly neutral (see § 12). The ψ -operator of the field of such particles satisfies the condition

$$\widehat{\psi}(t, \mathbf{r}) = \widehat{\psi}^C(t, \mathbf{r}).$$

For spin-1/2 particles this yields the conditions (in the spinor representation)²⁴

$$\xi^\alpha = -i\bar{\eta}^{\dot{\alpha}+}, \quad \widehat{\eta}_{\dot{\alpha}} = -i\widehat{\xi}_{\alpha}^+ \quad (27.4)$$

Like any relations expressing physical properties, these conditions are invariant under the *CPT* transformation²⁵. It is readily verified that they are in fact invariant not only under *CPT* as a whole, but also under each of the three transformations individually.

In § 19 we agreed to define spinor inversion as the transformation for which $\widehat{P}^2 = -1$, and we have followed this convention throughout. It is easy to see that the result obtained above concerning the relative parity of particle and antiparticle is independent, as it should be, of the particular convention adopted for the definition of inversion.

If inversion is defined by the condition $\widehat{P}^2 = 1$, then in place of (27.1) we have

$$P: \quad \xi^\alpha \rightarrow \eta_{\dot{\alpha}}, \quad \eta_{\dot{\alpha}} \rightarrow \xi^\alpha. \quad (27.5)$$

The charge-conjugate function then transforms according to the rule

$$\xi^{c\alpha} \rightarrow -\eta_{\dot{\alpha}}^c, \quad \eta_{\dot{\alpha}}^c \rightarrow -\xi^{c\alpha},$$

which differs from (27.5) by a sign. Accordingly, the three-dimensional spinors Φ transform as

$$\Phi^{(e)\alpha} \rightarrow \Phi^{(e)\alpha}, \quad \Phi^{(p)\alpha} \rightarrow -\Phi^{(p)\alpha},$$

so that the product $\Phi^{(e)}\Phi^{(p)}$ remains a pseudoscalar, as before.

The only possible physical distinction between the two conventions for inversion is that under the definition (27.5) the true-neutrality condition for the field would not be invariant under this transformation (or under *CP*): it would reverse the relative sign of the two sides of the equalities (27.4). In practice, truly neutral particles with spin 1/2 are unknown, and at present one cannot say whether the difference between the two definitions of inversion carries any real physical significance²⁶.

Problem

Determine the charge parity of positronium (a hydrogen-like system of an electron and a positron).

Solution. The wave function of two fermions must be antisymmetric under the simultaneous interchange of the coordinates, spins, and charge variables of the particles

²⁴In the Majorana representation, true neutrality simply means Hermiticity of the operator $\widehat{\psi}'$ (see the problem to § 26).

²⁵More precisely, the *CPT* transformation must be defined in this case so as to leave relations of the type (27.4) invariant. This is achieved by an appropriate choice of the phase factor in the definition of U_T (see the footnote on p. 121).

²⁶The incomplete equivalence of the two definitions of inversion was pointed out by *Racah* (G. Racah, 1937).

(cf. the problem to § 13). Interchanging the first multiplies the function by $(-1)^l$, the second by $(-1)^{l+S}$ (where $S = 0$ or 1 is the total spin of the system), and the third by the sought quantity C . From the condition $(-1)^l(-1)^{l+S}C = -1$ we find

$$C = (-1)^{l+S}.$$

Since the intrinsic parities of electron and positron are opposite, the spatial parity of the system is $P = (-1)^{l+1}$. The combined parity is $CP = (-1)^{S+1}$.

§28. Bilinear forms

Let us examine the transformation properties of the various bilinear forms that can be constructed from the components of ψ and ψ^* . Such forms are of great importance throughout quantum mechanics; among them is the 4-current density (21.11).

Since ψ and ψ^* each have four components, one can construct $4 \cdot 4 = 16$ independent bilinear combinations. Their classification by transformation properties follows immediately from the multiplication rules for two arbitrary bispinors listed in § 19 (which in the present case are ψ and ψ^*). Specifically, one can form a scalar (denoted S), a pseudoscalar (P), a mixed spinor of rank two equivalent to a true 4-vector V^μ (four independent quantities), a mixed rank-two spinor equivalent to a 4-pseudovector A^μ (four quantities), and a rank-two bispinor equivalent to an antisymmetric 4-tensor $T^{\mu\nu}$ (six quantities).

In a representation-independent notation these combinations are written as follows:

$$S = \bar{\psi}\psi, \quad P = i\bar{\psi}\gamma^5\psi, \quad (\text{III.2})$$

$$V^\mu = \bar{\psi}\gamma^\mu\psi, \quad A^\mu = \bar{\psi}\gamma^\mu\gamma^5\psi, \quad T^{\mu\nu} = i\bar{\psi}\sigma^{\mu\nu}\psi, \quad (\text{28.1})$$

where

$$\sigma^{\mu\nu} = \frac{1}{2}(\gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu) = (\boldsymbol{\alpha}, i\boldsymbol{\Sigma}) \quad (\text{28.2})$$

(the listing of components in (28.2) follows (19.15))²⁷. All the expressions written above are real.

The scalar and pseudoscalar character of S and P is evident from their spinor-representation forms: $S = \xi^*\eta + \eta^*\xi$, $P = i(\xi^*\eta - \eta^*\xi)$, which are precisely the expressions (19.7) and (19.8). That V^μ is a vector then follows from the Dirac equation: multiplying $\hat{p}_\mu\gamma^\mu\psi = m\psi$ on the left by $\bar{\psi}$ gives

$$(\bar{\psi}\hat{p}_\mu\gamma^\mu\psi) = m\bar{\psi}\psi;$$

the right-hand side is a scalar, so the left-hand side must be one as well.

The pattern governing the construction of the quantities (28.1) is straightforward: they are assembled as though the matrices γ^μ formed a 4-vector, γ^5 were a pseudoscalar, and the flanking factors $\bar{\psi}$ and ψ combined into a scalar²⁸.

²⁷Under a unitary change of representation $\psi \rightarrow U\psi$, $\gamma \rightarrow U\gamma U^{-1}$, $\bar{\psi} \rightarrow \bar{\psi}U^{-1}$, so the invariance of the bilinear forms is obvious.

²⁸The absence of bilinear forms having the character of a symmetric 4-tensor, obvious from the spinor representation, is clear from this rule as well: the symmetric combination $\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = 2g^{\mu\nu}$ would reduce the form to a scalar.

Second-quantised bilinear forms are obtained by replacing the ψ -functions in (28.1) with ψ -operators. For greater generality we shall assume that the two ψ -operators refer to the fields of different particles, distinguishing them by subscripts a and b . Let us determine how these operator forms transform under charge conjugation. Noting that ²⁹

$$\widehat{\psi}^C = U_C \widetilde{\widehat{\psi}}, \quad \widehat{\bar{\psi}}^C = U_C^\dagger \widehat{\bar{\psi}}, \quad (28.3)$$

and employing (26.3) and (26.21):

$$\begin{aligned} \widehat{\bar{\psi}}_a^C \widehat{\psi}_b^C &= \widehat{\bar{\psi}}_a U_C^\dagger U_C \widetilde{\widehat{\psi}}_b = -\widehat{\bar{\psi}}_a U_C^\dagger U_C \widehat{\bar{\psi}}_b = -\widehat{\bar{\psi}}_a \widehat{\bar{\psi}}_b, \\ \widehat{\bar{\psi}}_a^C \gamma^\mu \widehat{\psi}_b^C &= \widehat{\bar{\psi}}_a U_C^\dagger \gamma^\mu U_C \widetilde{\widehat{\psi}}_b = -\widehat{\bar{\psi}}_a U_C^\dagger \gamma^\mu U_C \widehat{\bar{\psi}}_b = \widehat{\bar{\psi}}_a \gamma^\mu \widehat{\bar{\psi}}_b. \end{aligned}$$

Restoring the operators to their original order ($\widehat{\bar{\psi}}$ to the left of $\widehat{\psi}$) reverses the sign of the product by the Fermi commutation rules (25.4) (and also generates state-independent terms, which we discard as in the analogous derivation of § 13). We thus obtain

$$\widehat{\bar{\psi}}_a^C \widehat{\psi}_b^C = \widehat{\bar{\psi}}_b \widehat{\psi}_a, \quad \widehat{\bar{\psi}}_a^C \gamma^\mu \widehat{\psi}_b^C = -\widehat{\bar{\psi}}_b \gamma^\mu \widehat{\psi}_a.$$

Treating the remaining forms in the same way, we find the following charge-conjugation rules ³⁰:

$$\begin{aligned} C : \quad \widehat{S}_{ab} &\rightarrow \widehat{S}_{ba}, \quad \widehat{P}_{ab} \rightarrow \widehat{P}_{ba}, \quad (\widehat{V}^0, \widehat{\mathbf{V}})_{ab} \rightarrow (\widehat{V}^0, -\widehat{\mathbf{V}})_{ba}, \\ (\widehat{A}^0, \widehat{\mathbf{A}})_{ab} &\rightarrow (\widehat{A}^0, -\widehat{\mathbf{A}})_{ba}, \quad \widehat{T}_{ab}^{\mu\nu} \rightarrow (\widehat{\mathbf{p}}, \widehat{\mathbf{a}})_{ab} \rightarrow (\widehat{\mathbf{p}}, -\widehat{\mathbf{a}})_{ba} \end{aligned} \quad (28.4)$$

($\widehat{\mathbf{p}}, \widehat{\mathbf{a}}$ are the three-dimensional vectors equivalent to the components of $\widehat{T}^{\mu\nu}$ according to (19.15)).

The behaviour of the same forms under time reversal is found analogously. One must bear in mind (see § 13) that this operation entails a reversal of the operator ordering, so that, for instance,

$$(\widehat{\bar{\psi}}_a \widehat{\psi}_b)^T = \widehat{\bar{\psi}}_b^T \widehat{\psi}_a^T.$$

Substituting

$$\widehat{\psi}^T = U_T \widetilde{\widehat{\psi}}, \quad \widehat{\bar{\psi}}^T = -U_T^\dagger \widehat{\bar{\psi}}, \quad (28.5)$$

we get

$$(\widehat{\bar{\psi}}_a \widehat{\psi}_b)^T = -\widehat{\bar{\psi}}_b U_T U_T^\dagger \widehat{\psi}_a = \widehat{\bar{\psi}}_b \widehat{\psi}_a.$$

Carrying out the same procedure for all the other forms yields

$$\begin{aligned} T : \quad \widehat{S}_{ab} &\rightarrow \widehat{S}_{ba}, \quad \widehat{P}_{ab} \rightarrow -\widehat{P}_{ba}, \quad (\widehat{V}^0, \widehat{\mathbf{V}})_{ab} \rightarrow (\widehat{V}^0, -\widehat{\mathbf{V}})_{ba}, \\ (\widehat{A}^0, \widehat{\mathbf{A}})_{ab} &\rightarrow (\widehat{A}^0, -\widehat{\mathbf{A}})_{ba}, \quad \widehat{T}_{ab}^{\mu\nu} \rightarrow (\widehat{\mathbf{p}}, \widehat{\mathbf{a}})_{ab} \rightarrow (\widehat{\mathbf{p}}, -\widehat{\mathbf{a}})_{ba} \end{aligned} \quad (28.6)$$

²⁹To derive the second relation from the first, write $\bar{\psi}^C = [U_C (\widehat{\bar{\psi}}^* \gamma^{0*})] \gamma^0 = -\bar{\gamma}^0 U_C^\dagger \gamma^0 \widehat{\bar{\psi}} = \bar{\gamma}^0 \gamma^{0*} U_C^\dagger \widehat{\bar{\psi}} = U_C^\dagger \widehat{\bar{\psi}}$ (using (26.3), (26.21) and the Hermiticity of γ^0).

³⁰Note that for bilinear forms built from ψ -functions (rather than ψ -operators), the transformations (28.4) would carry the opposite sign, since restoring the original ordering of the factors $\bar{\psi}$ and ψ would not entail a sign change.

($\hat{\mathbf{p}}, \hat{\mathbf{a}}$ are the three-dimensional vectors equivalent to the components of $\hat{T}^{\mu\nu}$ according to (19.15)).

Under spatial inversion, on the other hand, in accordance with the tensor character of the quantities³¹,

$$P: \quad \hat{S}_{ab} \rightarrow \hat{S}_{ab}, \quad \hat{P}_{ab} \rightarrow -\hat{P}_{ab}, \quad (\hat{V}^0, \hat{\mathbf{V}})_{ab} \rightarrow (\hat{V}^0, -\hat{\mathbf{V}})_{ab}, \\ (\hat{A}^0, \hat{\mathbf{A}})_{ab} \rightarrow (-\hat{A}^0, \hat{\mathbf{A}})_{ab}, \quad \hat{T}_{ab}^{\mu\nu} \rightarrow (\hat{\mathbf{p}}, \hat{\mathbf{a}})_{ab} \rightarrow (-\hat{\mathbf{p}}, \hat{\mathbf{a}})_{ab}. \quad (28.7)$$

Finally, the joint application of all three operations leaves $\hat{S}_{ab}, \hat{P}_{ab}, \hat{T}_{ab}^{\mu\nu}$ unchanged and reverses the sign of every $\hat{V}_{ab}^\mu, \hat{A}_{ab}^\mu$. This is exactly what one expects from the interpretation of this combined transformation as a 4-inversion: since a 4-inversion is equivalent to a rotation of the four-dimensional coordinate system, it draws no distinction between true tensors and pseudotensors of any rank.

Consider now pairwise products of bilinear forms constructed from four distinct functions $\psi^a, \psi^b, \psi^c, \psi^d$. Different results are obtained depending on which pairs of functions are multiplied together. It turns out, however, that any such product can be reduced to products of bilinear forms with fixed pairings of the factors (*W. Pauli, M. Fierz, 1936*). We shall derive the identity underlying this reduction.

Consider the set of 4×4 matrices

$$1, \quad \gamma^5, \quad \gamma^\mu, \quad i\gamma^\mu\gamma^5, \quad i\sigma^{\mu\nu} \quad (28.8)$$

(1 being the unit matrix). Numbering these 16 ($= 1 + 1 + 4 + 4 + 6$) matrices in some definite order, we denote them by γ^A ($A = 1, \dots, 16$), and the same matrices with the 4-tensor indices (μ, ν) lowered by γ_A . They possess the following properties:

$$\text{Sp } \gamma^A = 0 \quad (\gamma^A \neq 1), \quad \gamma^A \gamma_A = 1, \quad \frac{1}{4} \text{Sp } \gamma^A \gamma_B = \delta_B^A. \quad (28.9)$$

By virtue of the last of these properties, the matrices γ^A are linearly independent. Since their number equals the number of elements (4×4) of a 4×4 matrix, the γ^A form a complete set in terms of which any 4×4 matrix Γ can be expanded:

$$\Gamma = \sum_A c_A \gamma^A, \quad c_A = \frac{1}{4} \text{Sp } \gamma_A \Gamma, \quad (28.10)$$

or, written out explicitly with matrix indices ($i, k = 1, 2, 3, 4$):

$$\Gamma_{ik} = \frac{1}{4} \sum_A \Gamma_{lm} \gamma_{ml}^A \gamma_{Ai k}.$$

Supposing, in particular, that Γ has only one non-vanishing element (Γ_{lm}), we obtain the desired identity (the “completeness relation”)

$$\delta_{il} \delta_{km} = \frac{1}{4} \sum_A \gamma_{Aik} \gamma_{ml}^A. \quad (28.11)$$

³¹To avoid misunderstanding, recall that the T and P transformations also require a change in the arguments of the functions on the right-hand sides (the transformed forms in (28.6), (28.7)) to $x^T = (-t, \mathbf{r})$ and $x^P = (t, -\mathbf{r})$, respectively, if the left-hand sides are functions of $x = (t, \mathbf{r})$.

Multiplying both sides by $\bar{\psi}_i^a \psi_k^b \bar{\psi}_m^c \psi_l^d$ gives

$$(\bar{\psi}^a \psi^b)(\bar{\psi}^c \psi^d) = \frac{1}{4} \sum_A (\bar{\psi}^a \gamma_A \psi^b)(\bar{\psi}^c \gamma^A \psi^d). \quad (28.12)$$

This is one identity of the kind described above: it expresses the product of two scalar bilinear forms in terms of products of forms built from different pairings of the factors³².

Further identities of this type are obtained from (28.12) by substituting

$$\psi^d \rightarrow \gamma^B \psi^d, \quad \psi^b \rightarrow \gamma^C \psi^b$$

and using the expansion (see the problem)

$$\gamma^A \gamma^B = \sum_R c_R \gamma^R, \quad c_R = \frac{1}{4} \text{Sp } \gamma^A \gamma^B \gamma_R.$$

We also record here, for later reference, the analogue of (28.11) for 2×2 matrices. A complete set of linearly independent 2×2 matrices σ^A ($A = 1, \dots, 4$) is formed by

$$1, \quad \sigma_x, \quad \sigma_y, \quad \sigma_z. \quad (28.13)$$

For these

$$\text{Sp } \sigma^A = 0 \quad (\sigma^A \neq 1), \quad \frac{1}{2} \text{Sp } \sigma^A \sigma^B = \delta_{AB}. \quad (28.14)$$

The completeness relation reads:

$$\delta_{\alpha\gamma} \delta_{\beta\delta} = (1/2) \sum_A \sigma_{\alpha\beta}^A \sigma_{\delta\gamma}^A = (1/2) \delta_{\alpha\beta} \delta_{\delta\gamma} + (1/2) \delta_{\alpha\delta} \delta_{\beta\gamma} \quad (28.15)$$

($\alpha, \beta, \dots = 1, 2$), or equivalently:

$$\sigma_{\alpha\beta} \sigma_{\delta\gamma} = -(1/2) \sigma_{\alpha\gamma} \sigma_{\delta\beta} + (3/2) \delta_{\alpha\gamma} \delta_{\delta\beta}. \quad (28.16)$$

Problem

Derive formulae analogous to (28.12) for the scalar products of two bilinear forms of types P, V, A, T .

Solution. Denote:

$$\begin{aligned} J_S &= (\bar{\psi}^a \psi^b)(\bar{\psi}^c \psi^d), & J_P &= (\bar{\psi}^a \gamma^5 \psi^b)(\bar{\psi}^c \gamma^5 \psi^d), \\ J_V &= (\bar{\psi}^a \gamma^\mu \psi^b)(\bar{\psi}^c \gamma_\mu \psi^d), & J_A &= (\bar{\psi}^a i \gamma^\mu \gamma^5 \psi^b)(\bar{\psi}^c i \gamma_\mu \gamma^5 \psi^d), \\ J_T &= (\bar{\psi}^a i \sigma^{\mu\nu} \psi^b)(\bar{\psi}^c i \sigma_{\mu\nu} \psi^d), \end{aligned}$$

³²We recall, to avoid confusion, that the forms here are built from ψ -functions. For forms built from anticommuting ψ -operators the sign of the rearrangement would be reversed.

and let primed letters denote the same products with ψ^b and ψ^d interchanged. Applying the procedure described in the text yields

$$\begin{aligned} 4J'_S &= J_S + J_V + J_T + J_A + J_P, \\ 4J'_V &= 4J_S - 2J_V + 2J_A - 4J_P, \\ 4J'_T &= 6J_S - 2J_T + 6J_P, \\ 4J'_A &= 4J_S + 2J_V - 2J_A - 4J_P, \\ 4J'_P &= J_S - J_V - J_T - J_A + J_P \end{aligned}$$

(the first row is formula (28.12)).

§ 29. Polarisation density matrix

The coordinate dependence of the wave function ψ describing free motion with momentum $\mathbf{p}$ (a plane wave) reduces to the common factor $e^{i\mathbf{p}\mathbf{r}}$, while the amplitude u_p plays the role of a spin wave function. In such a (pure) state the particle is fully polarised (see III, § 59). In the non-relativistic theory

this means that the particle's spin has a definite spatial direction (more precisely, there exists a direction along which the spin projection takes the definite value $+1/2$). In a relativistic theory such a characterisation of the state is impossible in an arbitrary reference frame, owing to the non-conservation of the spin vector (already noted in § 23). Purity of the state signifies only that the spin has a definite direction in the particle's rest frame.

In a state of partial polarisation no definite amplitude exists; there is only a *polarisation density matrix* ρ_{ik} ($i, k = 1, 2, 3, 4$ being bispinor indices). We define this matrix so that in a pure state it reduces to the product

$$\rho_{ik} = u_{pi} \bar{u}_{pk}. \quad (29.1)$$

The corresponding normalisation condition for ρ is

$$\text{Sp } \rho = 2m \quad (29.2)$$

(cf. (23.4)).

In a pure state the mean value of the spin is given by

$$\mathbf{s} = \frac{1}{2} \int \psi^* \boldsymbol{\Sigma} \psi d^3x = \frac{1}{4\varepsilon} u_p^+ \boldsymbol{\Sigma} u_p = \frac{1}{4\varepsilon} \bar{u}_p \gamma^0 \boldsymbol{\Sigma} u_p. \quad (29.3)$$

The corresponding expression for a partially polarised state is

$$\mathbf{s} = \frac{1}{4\varepsilon} \text{Sp}(\rho \gamma^0 \boldsymbol{\Sigma}) = \frac{1}{4\varepsilon} \text{Sp}(\rho \gamma^5 \boldsymbol{\gamma}). \quad (29.4)$$

The amplitudes u_p and $\bar{u}_p$ satisfy the algebraic equations

$$(\gamma p - m)u_p = 0, \quad \bar{u}_p(\gamma p - m) = 0.$$

The matrix (29.1) therefore obeys

$$(\gamma p - m)\rho = 0, \quad \rho(\gamma p - m) = 0. \quad (29.5)$$

The density matrix must satisfy the same linear equations in the general case of a mixed (in spin) state as well (cf. the analogous argument in III, § 14).

When a free particle is considered in its rest frame, the non-relativistic theory applies. In that theory the state of partial polarisation is completely specified by three parameters—the components of the mean spin vector $\mathbf{s}$ (see III, § 59). It is therefore clear that the same parameters will determine the polarisation state after any Lorentz boost, i.e. for a moving particle.

Denote the doubled mean spin vector in the rest frame by ζ (in a pure state $|\zeta| = 1$; in a mixed state $|\zeta| < 1$). For a four-dimensional description of the polarisation state it is convenient to introduce a 4-vector a^μ that coincides in the rest frame with the three-vector ζ ; since ζ is an axial vector, a^μ is a 4-pseudovector. This 4-vector is orthogonal to the 4-momentum in the rest frame (where $a^\mu = (0, \zeta)$ and $p^\mu = (m, 0)$), and hence in every frame:

$$a_\mu p^\mu = 0. \quad (29.6)$$

In an arbitrary frame one also has

$$a_\mu a^\mu = -\zeta^2. \quad (29.7)$$

The components of a^μ in a frame where the particle moves with velocity $\mathbf{v} = \mathbf{p}/\varepsilon$ are obtained by Lorentz-boosting from the rest frame:

$$a^0 = \frac{|\mathbf{p}|}{m} \zeta_{\parallel}, \quad \mathbf{a}_{\perp} = \zeta_{\perp}, \quad a_{\parallel} = \frac{\varepsilon}{m} \zeta_{\parallel}, \quad (29.8)$$

where the subscripts $\parallel$ and $\perp$ denote the components of ζ and $\mathbf{a}$ parallel and perpendicular to $\mathbf{p}$ ³³. These formulae can be written in vector form:

$$\mathbf{a} = \zeta + \frac{\mathbf{p}(\zeta \cdot \mathbf{p})}{m(\varepsilon + m)}, \quad a^0 = \frac{\mathbf{a} \cdot \mathbf{p}}{\varepsilon} = \frac{\mathbf{p} \cdot \zeta}{m}, \quad \mathbf{a}^2 = \zeta^2 + \frac{(\mathbf{p} \cdot \zeta)^2}{m^2}. \quad (29.9)$$

Consider first the unpolarised state ($\zeta = 0$). The density matrix can then depend only on the 4-momentum p . The unique matrix of this kind satisfying the equations (29.5) is

$$\rho = \frac{1}{2}(\gamma p + m) \quad (29.10)$$

(I. E. Tamm, 1930; H. B. G. Casimir, 1933). The numerical coefficient is fixed by the normalisation condition (29.2).

³³By their transformation properties, the components of the mean spin vector $\mathbf{s}$ (like those of any angular momentum) are, in relativistic mechanics, the spatial components of an antisymmetric tensor $S^{\lambda\mu}$. The 4-vector a^λ is related to this tensor by

$$S^{\lambda\mu} = \frac{1}{2m} \epsilon^{\lambda\mu\nu\rho} a_\nu p_\rho, \quad a^\lambda = -\frac{2}{m} \epsilon^{\lambda\mu\nu\rho} S_{\mu\nu} p_\rho.$$

Note that in an arbitrary frame the spatial part of a^λ does not coincide with $2\mathbf{s}$. One easily finds $2\mathbf{s}_{\parallel} = \frac{1}{m}(a_{\parallel}\varepsilon - a^0|\mathbf{p}|) = \zeta_{\parallel}$, $2\mathbf{s}_{\perp} = \frac{\varepsilon}{m}\mathbf{a}_{\perp} = \frac{\varepsilon}{m}\zeta_{\perp}$.

In the general case of partial polarisation ($\zeta \neq 0$) we seek the density matrix in the form

$$\rho = \frac{1}{4m}(\gamma p + m)\rho'(\gamma p + m), \quad (29.11)$$

which automatically satisfies (29.5). When $\zeta \neq 0$ the auxiliary matrix ρ' must reduce to the identity; since

$$(\gamma p + m)^2 = 2m(\gamma p + m),$$

equation (29.11) then coincides with (29.10). Furthermore, ρ' must contain the 4-vector a linearly as a parameter, i.e. it must have the form

$$\rho' = 1 - A\gamma^5(\gamma a); \quad (29.12)$$

the second term involves the scalar product of the pseudovector a with the “matrix 4-pseudovector” $\gamma^5\gamma$. To fix the coefficient A , write the density matrix in the rest frame:

$$\rho = \frac{m}{4}(1 + \gamma^0)(1 + A\gamma^5\gamma \cdot \zeta)(1 + \gamma^0) = \frac{m}{2}(1 + \gamma^0)(1 + A\gamma^5\gamma \cdot \zeta),$$

and compute the mean spin via (29.4). Using the trace rules listed in § 22, one readily finds that the only non-vanishing contribution to the required trace is

$$2s = \frac{1}{2m} \text{Sp}(\rho\gamma^5\gamma) = -\frac{A}{4} \text{Sp}((\gamma \cdot \zeta)\gamma) = A\zeta.$$

Setting this equal to ζ gives $A = 1$. To obtain the final expression for ρ , substitute (29.12) into (29.11) and commute ρ' past $(\gamma p + m)$; because a and p are orthogonal, γp anticommutes with γa :

$$(\gamma a)(\gamma p) = 2a \cdot p - (\gamma p)(\gamma a) = -(\gamma p)(\gamma a),$$

and therefore commutes with $\gamma^5(\gamma a)$.

The density matrix of a partially polarised electron is thus

$$\rho = \frac{1}{2}(\gamma p + m)[1 - \gamma^5(\gamma a)] \quad (29.13)$$

(*L. Michel, A. S. Wightman, 1955*). If ρ is known, the 4-vector a characterising the state (and with it the vector ζ) can be recovered from

$$a^\mu = \frac{1}{2m} \text{Sp}(\rho\gamma^5\gamma^\mu). \quad (29.14)$$

The formulae for the positron density matrix are analogous to those for the electron. Had we described a positron (with

4-momentum p) by a positron amplitude $u_p^{(\text{pos})}$ and a correspondingly defined density matrix $\rho^{(\text{pos})}$, there would be no difference from the electron case and $\rho^{(\text{pos})}$ would be given by the same formula (29.13). In actual calculations of scattering cross sections involving positrons, however, one deals (as we shall see later) not with $u_p^{(\text{pos})}$ but with the “negative-frequency” amplitudes u_{-p} . Accordingly, the polarisation density matrix (which we denote $\rho^{(-)}$) should be defined so that for a pure state it reduces to $u_{-p}i\bar{u}_{-p}k$.

By (26.1) the positron amplitude $u_p^{(\text{pos})} = U_C \bar{u}_{-p}$. Inversely:

$$u_{-p} = U_C u_p^{(\text{pos})}, \quad \bar{u}_{-p} = U_C^+ u_p^{(\text{pos})} = u_p^{(\text{pos})} U_C^*$$

(cf. (28.3)). If

$$\rho_{ik}^{(-)} = u_{-p i} \bar{u}_{-p k}, \quad \rho_{ik}^{(\text{pos})} = u_{pi}^{(\text{pos})} \bar{u}_{pk}^{(\text{pos})},$$

then these relations give

$$\rho^{(-)} = U_C \tilde{\rho}^{(\text{pos})} U_C^*. \quad (29.15)$$

Substituting (29.13) for $\rho^{(\text{pos})}$ and carrying out straightforward algebra (using (26.3) and (26.21)), we obtain

$$\rho^{(-)} = \frac{1}{2}(\gamma p - m)[1 - \gamma^5(\gamma a)]. \quad (29.16)$$

In particular, for an unpolarised state

$$\rho^{(-)} = \frac{1}{2}(\gamma p - m). \quad (29.17)$$

Henceforth, when speaking of the positron density matrix we shall mean $\rho^{(-)}$ and drop the superscript $(-)$ (the matrix $\rho^{(\text{pos})}$ is in practice never needed).

In various calculations one frequently has to average over spin states expressions of the form $\bar{u} F u (\equiv \bar{u}_i F_{ik} u_k)$, where F is some (4×4) matrix and u is the bispinor amplitude for a state with definite 4-momentum p . Such averaging is equivalent to replacing the products $u_k \bar{u}_i$ by the density matrix ρ_{ik} of a partially polarised state.

In particular, a complete average over the two independent spin states amounts to passing to the unpolarised state; by (29.10) we then have

$$\frac{1}{2} \sum_{\text{pol}} \bar{u}_p F u_p = \frac{1}{2} \text{Sp}(\gamma p + m) F. \quad (29.18)$$

Similarly, for the negative-frequency wave functions

$$\frac{1}{2} \sum_{\text{pol}} \bar{u}_{-p} F u_{-p} = \frac{1}{2} \text{Sp}(\gamma p - m) F. \quad (29.19)$$

If one is summing over spin states rather than averaging, the result is twice as large.

Let us trace how the density matrix (29.13) passes to its non-relativistic limit. To this end we go to the electron rest frame. In the standard representation the amplitudes u_p become two-component in this frame, and the density matrix must correspondingly become a 2×2 matrix. Indeed, in the rest frame

$$\rho = \frac{m}{2}(\gamma^0 + 1)(1 + \gamma^5 \boldsymbol{\gamma} \cdot \boldsymbol{\zeta}),$$

and using the explicit forms of the γ matrices (21.20) and (22.18) we find

$$\rho = \begin{pmatrix} \rho_{\text{nr}} & 0 \\ 0 & 0 \end{pmatrix}, \quad \rho_{\text{nr}} = m(1 + \boldsymbol{\sigma} \cdot \boldsymbol{\zeta}) \quad (29.20)$$

(the zeros denote 2×2 null matrices). Adopting the normalisation $\text{Sp } \rho_{\text{nr}} = 1$ conventional in the non-relativistic theory (instead of $2m$), one must divide this expression by $2m$, yielding

$$\frac{1}{2}(1 + \boldsymbol{\sigma} \cdot \boldsymbol{\zeta}),$$

in agreement with formula (59.6) (see III).

The non-relativistic limit of the positron density matrix is obtained in a similar way:

$$\rho = \begin{pmatrix} 0 & 0 \\ 0 & \rho_{\text{nr}} \end{pmatrix}, \quad \rho_{\text{nr}} = -m(1 + \boldsymbol{\sigma} \cdot \boldsymbol{\zeta}).$$

Finally, let us write down a simplified expression for the density matrix in the ultrarelativistic case. Setting $|\mathbf{p}| \approx \varepsilon$ in (29.8) (thereby neglecting quantities of relative order $(m/\varepsilon)^2$), substituting into (29.13) or (29.16), and choosing the direction of $\mathbf{p}$ as the x -axis, we have

$$\rho = \frac{1}{2}[\varepsilon(\gamma^0 - \gamma^1) \pm m] \left[1 - \gamma^5 \left(\frac{\varepsilon}{m}(\gamma^0 - \gamma^1)\zeta_{\parallel} - \zeta_{\perp}\gamma_{\perp} \right) \right],$$

where the upper sign refers to the electron and the lower to the positron. On expanding the product the leading terms cancel, and the next-order terms give

$$\rho = \frac{1}{2}\varepsilon(\gamma^0 - \gamma^1)[1 + \gamma^5(\pm\zeta_{\parallel} + \zeta_{\perp}\gamma_{\perp})]$$

or, writing $\varepsilon(\gamma^0 - \gamma^1)$ as γp :

$$\rho = \frac{1}{2}(\gamma p)[1 + \gamma^5(\pm\zeta_{\parallel} + \zeta_{\perp} \cdot \boldsymbol{\gamma}_{\perp})]. \quad (29.21)$$

This is the desired ultrarelativistic density matrix. Note that all components of the polarisation vector $\boldsymbol{\zeta}$ enter on an equal footing, as terms of the same order. Recall that $\zeta_{\parallel}$ is the component of this vector parallel ($\zeta_{\parallel} > 0$) or antiparallel ($\zeta_{\parallel} < 0$) to the particle's momentum. In particular, for a helicity eigenstate $\zeta_{\parallel} = 2\lambda = \pm 1$; the density matrix then takes the especially simple form

$$\rho = \frac{1}{2}(\gamma p)(1 \pm 2\lambda\gamma^5), \quad (29.22)$$

which coincides, as it should, with the density matrix of a neutrino or antineutrino—a particle with zero mass and definite helicity (see § 30).

§ 30. Two-component fermions

We saw in § 20 that the need to describe a spin-1/2 particle by two spinors (ξ and η) is tied to the particle having a mass. This requirement disappears when the mass vanishes. The wave equation for such a particle can be formulated in terms of a single spinor—say the dotted spinor η :

$$\widehat{p}^{\alpha\beta}\eta_{\beta} = 0, \quad (30.1)$$

or equivalently,

$$(\widehat{p}_0 + \widehat{\mathbf{p}} \cdot \boldsymbol{\sigma})\eta = 0. \quad (30.2)$$

In § 20 it was also noted that the wave equation containing the mass m is automatically symmetric under inversion (the transformation (20.4)). When the particle is described by a single spinor this symmetry is lost. There is, however, no necessity for it, since symmetry under inversion is not a universal law of nature.

The energy and momentum of a particle with $m = 0$ are related by $\varepsilon = |\mathbf{p}|$. For a plane wave ($\eta_p \propto e^{-ipx}$) the equation (30.2) therefore gives

$$(\mathbf{n} \cdot \boldsymbol{\sigma})\eta_p = -\eta_p, \quad (30.3)$$

where $\mathbf{n}$ is the unit vector along $\mathbf{p}$. The same equation

$$(\mathbf{n} \cdot \boldsymbol{\sigma})\eta_{-p} = -\eta_{-p} \quad (30.4)$$

holds also for the “negative-frequency” waves ($\eta_{-p} \propto e^{ipx}$). The second-quantised ψ -operator is

$$\widehat{\eta} = \sum_{\mathbf{p}} (\eta_p \widehat{a}_{\mathbf{p}} + \eta_{-p} \widehat{b}_{\mathbf{p}}^+), \quad \widehat{\eta}^+ = \sum_{\mathbf{p}} (\eta_p^* \widehat{a}_{\mathbf{p}}^+ + \eta_{-p}^* \widehat{b}_{\mathbf{p}}). \quad (30.5)$$

From this it follows, as usual, that η_{-p}^* are the antiparticle wave functions.

From the definition of the operators $\widehat{p}^{\alpha\beta}$ (20.1) it is clear that $\widehat{p}^{\alpha\beta*} = -\widehat{p}^{\dot{\alpha}\dot{\beta}}$. Hence the complex-conjugate spinor η^* satisfies $\widehat{p}^{\dot{\alpha}\dot{\beta}}\eta_{\dot{\beta}}^* = 0$, i.e.

$$\widehat{p}_{\dot{\alpha}\beta}\eta^{\beta*} = 0.$$

Denoting $\eta^{\beta*} = \xi^{\dot{\beta}}$ (thus expressing the fact that complex conjugation converts a dotted spinor into an undotted one), we see that the antiparticle wave functions satisfy

$$\widehat{p}_{\dot{\alpha}\beta}\xi^{\dot{\beta}} = 0, \quad (30.6)$$

or

$$(\widehat{p}_0 - \widehat{\mathbf{p}} \cdot \boldsymbol{\sigma})\xi = 0. \quad (30.7)$$

For a plane wave this gives

$$(\mathbf{n} \cdot \boldsymbol{\sigma})\xi_p = \xi_p. \quad (30.8)$$

Now $\frac{1}{2}(\mathbf{n} \cdot \boldsymbol{\sigma})$ is the operator of the spin projection on the direction of motion. Equations (30.3) and (30.8) therefore mean that states with a definite momentum are automatically helicity eigenstates—the spin projection along the direction of motion takes a definite value. If the particle’s spin is antiparallel to its momentum (helicity $-1/2$), then the antiparticle’s spin is aligned with its momentum (helicity $+1/2$). Particles with these properties are, perhaps, realised in nature as **neutrinos**. The particle with helicity $-1/2$ is conventionally called the neutrino, while the one with helicity $+1/2$ is called the antineutrino³⁴.

³⁴The existence of the neutrino was predicted theoretically by *Pauli* to explain β -decay properties (1931). Equation (30.1) was first considered by *Weyl* (*H. Weyl*, 1929). A neutrino theory based on these equations was formulated by *L. D. Landau*; *Lee, Yang and Salam* (*T. D. Lee, C. N. Yang, A. Salam*) in 1957.

The experimental question of whether the neutrino mass is exactly zero has not been settled definitively up to the present time. We shall henceforth use the term “neutrino” conventionally to designate a particle described by equation (30.3).

In connection with the non-degeneracy of the neutrino states with respect to the spin direction, recall the remark made in § 8 that a massless particle (with spin 1/2) possesses only an axial symmetry about the direction of its momentum. For a truly neutral particle—the photon—this symmetry includes both rotations about the axis and reflections in planes passing through it. For the neutrino, however, there is no symmetry under reflections, and one is left only with the group of rotations about the axis, which preserves the momentum projection of the spin but does not reverse its sign. Symmetry under reflections exists only on the condition that particle and antiparticle are simultaneously interchanged.

It should also be noted that the obligatory longitudinal polarisation means that, for the neutrino, spin is altogether inseparable from orbital angular momentum (just as, for the photon, it is inseparable from the obligatory transversality of the field; see § 6).

From a single spinor η (or ξ) one can form only four bilinear combinations, which together make up a 4-vector

$$j^\mu = (\eta^* \eta, \eta^* \sigma \eta). \quad (30.9)$$

It is easily verified that, by virtue of the equations

$$(\widehat{p}_0 + \widehat{\mathbf{p}} \cdot \boldsymbol{\sigma})\eta = 0, \quad \eta^*(\widehat{p}_0 - \widehat{\mathbf{p}} \cdot \boldsymbol{\sigma}) = 0,$$

the continuity equation $\partial_\mu j^\mu = 0$ is satisfied, i.e. j^μ serves as the particle current-density 4-vector.

The neutrino plane waves are conveniently normalised in the same fashion as was done in § 23 for massive particles:

$$\eta_p = \frac{1}{\sqrt{2\varepsilon}} u_p e^{-ipx}, \quad \eta_{-p} = \frac{1}{\sqrt{2\varepsilon}} u_{-p} e^{ipx}, \quad (30.10)$$

with the spinor amplitudes satisfying the invariant normalisation condition

$$u_{\pm p}^*(1, \boldsymbol{\sigma}) u_{\pm p} = 2(\varepsilon, \mathbf{p}). \quad (30.11)$$

The current density is then $j^0 = 1$, $\mathbf{j} = \mathbf{p}/\varepsilon = \mathbf{n}$.

Since a free neutrino with a given momentum is always fully polarised, the concept of a mixed spin state does not arise. It may nonetheless be useful to introduce a 2×2 polarisation “density matrix” defined simply as a rank-two spinor:

$$\rho_{\dot{\alpha}\dot{\beta}} = u_{\dot{\alpha}} u_{\dot{\beta}}^* \quad (30.12)$$

(with $\text{Sp } \rho = 2\varepsilon$). The expression for this matrix is obtained by noting that it must satisfy

$$(\varepsilon + \mathbf{p} \cdot \boldsymbol{\sigma})\rho = \rho(\varepsilon + \mathbf{p} \cdot \boldsymbol{\sigma}) = 0.$$

It follows that

$$\rho = \varepsilon - \mathbf{p} \cdot \boldsymbol{\sigma}. \quad (30.13)$$

When considering various interaction processes, the neutrino may appear alongside other particles (with spin 1/2) that have a mass and are therefore described by four-component wave functions. In such cases it is convenient to maintain a uniform notation

by formally introducing a “bispinor” wave function for the neutrino as well, $\psi = \begin{pmatrix} 0 \\ \eta \end{pmatrix}$, two of whose components are, however, identically zero. But such a form of ψ would, generally speaking, be spoiled by a change to a different (non-spinor) representation. This difficulty is circumvented by observing that in the spinor representation one has the identity

$$\frac{1 + \gamma^5}{2} \begin{pmatrix} \xi \\ \eta \end{pmatrix} = \begin{pmatrix} 0 \\ \eta \end{pmatrix}, \quad (\eta^* \eta^*) \frac{1 - \gamma^5}{2} = (\eta^* 0),$$

where ξ is an arbitrary “ballast” spinor that drops out thanks to the matrix γ^5 from (22.18). The condition of true “two-componentness” of the neutrino is therefore maintained when describing it by a four-component ψ in any representation, provided one means by ψ a solution of the Dirac equation with $m = 0$:

$$(\gamma p)\psi = 0, \quad (30.14)$$

subject to the additional constraint $\frac{1}{2}(1 + \gamma^5)\psi = \psi$, i.e.

$$\gamma^5\psi = \psi. \quad (30.15)$$

This condition can be taken into account, once and for all, in every formula involving ψ and $\bar{\psi}$ by making the substitution

$$\psi \rightarrow \frac{1 + \gamma^5}{2}\psi, \quad \bar{\psi} \rightarrow \bar{\psi} \frac{1 - \gamma^5}{2}. \quad (30.16)$$

The current-density 4-vector then takes the form (cf. the replacement (30.16) in the expression $\bar{\psi}\gamma^\mu\psi$)

$$j^\mu = \frac{1}{4}\bar{\psi}(1 - \gamma^5)\gamma^\mu(1 + \gamma^5)\psi = \frac{1}{2}\bar{\psi}\gamma^\mu(1 + \gamma^5)\psi. \quad (30.17)$$

In accordance with the same rule, the 4×4 density matrix of the neutrino should be written as

$$\rho = \frac{1}{4}(1 + \gamma^5)(\gamma p)(1 - \gamma^5) = \frac{1}{2}(1 + \gamma^5)(\gamma p). \quad (30.18)$$

In the spinor representation it reduces, as it should, to the 2×2 matrix (30.13):

$$\rho = \begin{pmatrix} 0 & 0 \\ \varepsilon - \boldsymbol{\sigma} \cdot \mathbf{p} & 0 \end{pmatrix}.$$

The corresponding formulae for the antineutrino differ from those written above by a change of sign in front of γ^5 .

The neutrino is an electrically neutral particle. A neutrino with the properties described above is not, however, a truly neutral particle. We note in this connection that the “neutrino field” described by a two-component spinor is, by the number of particle states it permits (but not, of course, by any other physical property), equivalent to a truly neutral field described by a four-component bispinor. In place of particles and antiparticles with definite helicities, there would be the same number of states of a

single particle with two possible helicity values, and symmetry under inversion would be automatically preserved.

Note, however, that the vanishing of the “four-component” neutrino mass would be, so to speak, “accidental”—it would not be related to any symmetry property of the wave equation (which permits a non-zero mass as well). For this reason the account of various interactions of such a particle would automatically generate a mass that, while small, would not be strictly zero.

§ 31. Wave equation for a spin-3/2 particle

A spin-3/2 particle is described in its rest frame by a symmetric 3-spinor of rank three (with $2s + 1 = 4$ independent components). Accordingly, in an arbitrary frame the description may involve the 4-spinors $\xi^{\alpha\beta\dot{\gamma}}$, $\eta_{\dot{\alpha}\beta\gamma}$, $\zeta^{\alpha\beta\gamma}$, and $\chi_{\dot{\alpha}\beta\dot{\gamma}}$, each symmetric under interchange of all identical (dotted or undotted) indices; under inversion the spinors in the first pair, and likewise those in the second pair, transform into each other.

For the rest-frame 4-spinors $\xi^{\alpha\beta\dot{\gamma}}$ and $\eta_{\dot{\alpha}\beta\gamma}$, symmetric over all three indices, to reduce to 3-spinors they must satisfy the conditions

$$\widehat{p}^{\dot{\alpha}\beta} \eta_{\dot{\alpha}\beta\gamma} = 0, \quad \widehat{p}_{\alpha\dot{\beta}} \xi^{\alpha\beta\dot{\gamma}} = 0. \quad (31.1)$$

Indeed, in the rest frame

$$\widehat{p}^{\dot{\alpha}\beta} \rightarrow \widehat{p}_0 \delta_{\alpha}^{\beta} = m \delta_{\alpha}^{\beta}$$

(as is clear from (20.1)). The conditions (31.1) therefore lead to the equalities

$$\delta_{\alpha}^{\beta} \eta'^{\alpha}{}_{\beta\gamma} = 0, \quad \delta_{\alpha}^{\beta} \xi'^{\alpha}{}_{\beta\gamma} = 0,$$

where the primed letters denote the corresponding 3-spinors; in other words, these spinors vanish upon contraction over the in-

indices $\alpha\beta$, which means that they are symmetric in those indices, and hence in all three.

The differential relation between the spinors ξ and η is established by the equations

$$\widehat{p}^{\dot{\beta}\gamma} \eta'_{\dot{\alpha}\beta} = m \xi^{\beta\gamma}{}_{\dot{\alpha}}, \quad \widehat{p}_{\dot{\beta}\gamma} \xi^{\beta\gamma}{}_{\dot{\alpha}} = m \eta_{\dot{\alpha}\beta\gamma}. \quad (31.2)$$

The symmetry of the left-hand sides of these equations (in the indices β, γ or $\alpha, \dot{\beta}$) is ensured by the conditions (31.1), by virtue of which they vanish upon contraction over all indices. In the rest frame the 3-spinors ξ' and η' coincide because of equations (31.2). Eliminating η (or ξ) from (31.2), one finds that each component of the spinors ξ and η satisfies the second-order equation

$$(\widehat{p}^2 - m^2) \xi^{\alpha\beta\dot{\gamma}} = 0. \quad (31.3)$$

The system of equations (31.1), (31.2) constitutes the complete set of wave equations for a spin-3/2 particle³⁵. Adding the spinors ζ, χ would contribute nothing new. They are constructed according to

$$m \zeta^{\alpha\beta\gamma} = \widehat{p}^{\alpha\dot{\delta}} \eta'^{\beta\gamma}{}_{\dot{\delta}}, \quad m \chi_{\dot{\alpha}\beta\dot{\gamma}} = \widehat{p}_{\dot{\alpha}\delta} \xi^{\delta}{}_{\beta\dot{\gamma}}.$$

³⁵For the Lagrangian formulation of these equations see the article by *Fierz* and *Pauli* cited on p. 76.

The equations for spin-3/2 particles can also be cast in a different form that exploits the vector properties of the spinors (W. Rarita, J. Schwinger, 1941; A. S. Davydov, I. E. Tamm, 1942). A pair of spinor indices $\alpha\dot{\beta}$ is associated with a single four-dimensional vector index μ . The components of a rank-three spinor $\xi^{\alpha\beta\dot{\gamma}}$ can therefore be brought into correspondence with the components of “mixed” quantities ψ^μ_l carrying one vector and one spinor index. Likewise, the spinor $\eta^{\dot{\beta}\alpha\gamma}$ corresponds to quantities ψ^μ_l , and the totality of both spinors to a “vector” bispinor ψ_μ (with the bispinor index suppressed). The wave equation is then written as a “Dirac equation” for each of the vector components ψ_μ :

$$(\gamma\hat{p} - m)\psi_\mu = 0 \quad (31.4)$$

with the supplementary condition

$$\gamma^\mu\psi_\mu = 0. \quad (31.5)$$

Using the expressions for the matrices γ^μ in the spinor representation (18.6), (18.7), it is straightforward to verify that the equations (31.2) are contained in (31.4), and the condition (31.5) is equivalent to the symmetry requirement on the spinors $\xi^{\alpha\beta\dot{\gamma}}$ and $\eta^{\dot{\alpha}\beta\gamma}$ in the indices $\beta\dot{\gamma}$ or $\beta\gamma$. Multiplying equation (31.4) on γ^μ and using (31.5):

$$\gamma^\mu\gamma^\nu\hat{p}_\nu\psi_\mu = 0$$

or, applying the commutation rules for the γ matrices,

$$2g^{\mu\nu}\hat{p}_\nu\psi_\mu - \gamma^\nu\hat{p}_\nu\gamma^\mu\psi_\mu = 0. \quad (31.6)$$

The second term vanishes by (31.5), and the first gives

$$\hat{p}^\mu\psi_\mu = 0. \quad (31.7)$$

It is easy to see that this condition, which follows automatically from (31.4) and (31.5), is equivalent to (31.1).

Finally, there is yet another way to formulate the wave equation. It consists in introducing quantities ψ_{ikl} ($i, k, l = 1, 2, 3, 4$) with three bispinor indices, symmetric under permutation (V. Bargmann, E. P. Wigner, 1948). The totality of these quantities is equivalent to the set of all four spinors ξ, η, ζ, χ . The wave equation is written as a system of “Dirac equations”

$$\hat{p}_\mu\gamma_{im}^\mu\psi_{mkl} = m\psi_{ikl}. \quad (31.8)$$

It is readily seen that these equations already yield the correct number of independent components (four), and no additional constraints are needed. Indeed, in the rest frame (31.8) reduces to

$$\gamma_{im}^0\psi_{mkl} = \psi_{ikl},$$

by virtue of which all components with $i, k, l = 3, 4$ (in the standard representation) vanish, i.e. ψ_{ikl} reduces to the components of a rank-three 3-spinor.

The results presented here generalise in an obvious way to particles of any half-integer spin s . In the description via equations of the type (31.4), (31.5) the wave function is a symmetric 4-tensor of rank $(2s - 1)/2$ carrying one bispinor index. In the description via equations of the type (31.8) the wave function has $2s$ bispinor indices over which it is symmetric.

A Particle in an External Field

§ 32. The Dirac equation for an electron in an external field

The wave equations for free particles express, in essence, only those properties that follow from the general requirements of space-time symmetry. The physical processes that occur with particles depend on the nature of their interactions.

A description of the electromagnetic interactions of particles in a relativistic quantum theory turns out to be possible by generalising the method employed for this purpose in both classical and non-relativistic quantum theories.

This method is, however, applicable only to the electromagnetic interactions of particles that are incapable of strong interactions. Electrons (and positrons) belong to this class, and thus the entire vast domain of quantum electrodynamics of electrons becomes accessible to the present theory. Muons are likewise incapable of strong interactions; they too are described by quantum electrodynamics in the regime of phenomena whose time scales are short compared with the muon lifetime (which is governed by weak interactions).

In this chapter we consider the class of problems in quantum electrodynamics that fall within the framework of the single-particle theory, where the number of particles is fixed and the interaction can be described in terms of an external electromagnetic field. Besides the conditions that permit the external field to be treated as given, the range of applicability of such a theory is further restricted by considerations connected with the so-called radiative corrections.

The wave equation of an electron in a given external field is obtained in the same way as in the non-relativistic theory (see III, § 111). Let $A^\mu = (\Phi, \mathbf{A})$ be the 4-potential of the external electromagnetic field ($\mathbf{A}$ is the vector potential, Φ the scalar potential). The desired equation is obtained by replacing in the Dirac equation the 4-momentum operator $\hat{p}$ with the difference $\hat{p} - eA$, where e is

the charge of the particle¹:

$$[\gamma(\hat{p} - eA) - m]\psi = 0. \quad (32.1)$$

The Hamiltonian corresponding to this equation is obtained from (21.13) by the same substitution:

$$\hat{H} = \alpha(\hat{\mathbf{p}} - e\mathbf{A}) + \beta m + e\Phi. \quad (32.2)$$

The invariance of the Dirac equation under a gauge transformation of the electromagnetic potentials is expressed by the fact that its form remains unchanged when the transformation $A \rightarrow A + i\hat{p}\chi$ (where χ is an arbitrary function) is accompanied by the

¹The charge is understood together with its sign, so that for the electron $e = -|e|$.

transformation of the wave function²

$$\psi \rightarrow \psi e^{ie\chi} \quad (32.3)$$

(cf. the analogous transformation for the Schrödinger equation in III, § 111).

The current density, expressed in terms of the wave function, is given by the same formula (21.11) $j = \bar{\psi} \gamma \psi$ as in the absence of an external field. It is easily verified that, on substituting (32.1) (together with equation (32.4) below) into the same calculation that was used to derive (21.11), the external field drops out and the continuity equation remains valid in its previous form.

Let us apply to equation (32.1) the operation of charge conjugation. To this end, write the equation as

$$\bar{\psi}[\gamma(\hat{p} + eA) + m] = 0, \quad (32.4)$$

which is obtained by complex conjugation from (32.1) in the same way as was done previously for equation (19.9) (bearing in mind that the 4-vector A is real). Rewriting this equation as

$$[\bar{\gamma}(\hat{p} + eA) + m]\bar{\psi} = 0,$$

multiplying on the left by the matrix U_C and using the relations (26.3), we find

$$[\gamma(\hat{p} + eA) - m](\hat{C}\psi) = 0. \quad (32.5)$$

Thus the charge-conjugate wave function satisfies an equation differing from the original one only by a

reversal of the sign of the charge. On the other hand, the operation of charge conjugation signifies a transition from particles to antiparticles. We see, therefore, that if the particles carry an electric charge, the charges of electron and positron are automatically opposite.

The first-order equation (32.1) can be converted into a second-order equation by applying the operator $\gamma(\hat{p} - eA) + m$:

$$[\gamma^\mu \gamma^\nu (\hat{p}_\mu - eA_\mu)(\hat{p}_\nu - eA_\nu) - m^2]\psi = 0.$$

Replacing $\gamma^\mu \gamma^\nu$ by

$$\gamma^\mu \gamma^\nu = \frac{1}{2}(\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu) + \frac{1}{2}(\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu) = g^{\mu\nu} + \sigma^{\mu\nu},$$

where $\sigma^{\mu\nu}$ is the antisymmetric “matrix 4-tensor” (28.2). On multiplying by $\sigma^{\mu\nu}$ one may antisymmetrise, i.e.

$$\begin{aligned} (\hat{p}_\mu - eA_\mu)(\hat{p}_\nu - eA_\nu) &\rightarrow \frac{1}{2}\{(\hat{p}_\mu - eA_\mu), (\hat{p}_\nu - eA_\nu)\}_- = \\ &= \frac{1}{2}e(-A_\mu \hat{p}_\nu + \hat{p}_\nu A_\mu - \hat{p}_\mu A_\nu + A_\nu \hat{p}_\mu) = \\ &= \frac{1}{2}ie(\partial_\nu A_\mu - \partial_\mu A_\nu) = -\frac{ie}{2}F_{\mu\nu} \end{aligned}$$

²The transformation (32.3) with a function $\chi(t, \mathbf{r})$ is sometimes called a “local gauge transformation”, as distinct from the “global gauge transformation” (12.10) with a constant phase α .

($F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ being the electromagnetic field tensor). The resulting second-order equation is

$$\left[(\hat{p} - eA)^2 - m^2 - \frac{i}{2} e F_{\mu\nu} \sigma^{\mu\nu} \right] \psi = 0. \quad (32.6)$$

The product $F_{\mu\nu} \sigma^{\mu\nu}$ can be written in three-dimensional form by expressing it in terms of its components:

$$\sigma^{\mu\nu} = (\boldsymbol{\alpha}, i\boldsymbol{\Sigma}), \quad F^{\mu\nu} = (-\mathbf{E}, \mathbf{H}).$$

We then obtain the second-order equation as

$$[(\hat{p} - eA)^2 - m^2 + e\boldsymbol{\Sigma} \cdot \mathbf{H} - ie\boldsymbol{\alpha} \cdot \mathbf{E}] \psi = 0, \quad (32.7)$$

or, in ordinary units,

$$\left[\left(\frac{i\hbar}{c} \frac{\partial}{\partial t} - \frac{e}{c} \Phi \right)^2 - \left(i\hbar \nabla + \frac{e}{c} \mathbf{A} \right)^2 - m^2 c^2 + \frac{e\hbar}{c} \boldsymbol{\Sigma} \cdot \mathbf{H} - i \frac{e\hbar}{c} \boldsymbol{\alpha} \cdot \mathbf{E} \right] \psi = 0. \quad (32.7a)$$

The appearance in these equations of the terms containing $\mathbf{E}$ and $\mathbf{H}$ is connected with the spin of the particle; we shall return to their discussion in the next section.

Among the solutions of the second-order equation there are, of course, “spurious” ones—solutions that do not satisfy the original first-order equation (32.1) (they represent solutions

of equation (32.1) with the opposite sign in front of m). Selecting the correct solutions in specific cases is usually straightforward. A systematic selection procedure is the following: if φ is an arbitrary solution of the second-order equation, the correct solution of the first-order equation is

$$\psi = [\gamma(\hat{p} - eA) + m] \varphi. \quad (32.8)$$

Indeed, multiplying this relation by $\gamma(\hat{p} - eA) - m$, one sees that the right-hand side vanishes if φ satisfies equation (32.6).

It should be emphasised that the method of introducing an external field into the relativistic wave equation—by replacing $\hat{p}$ with $\hat{p} - eA$ —is not entirely self-evident. In carrying it out we essentially relied on an additional principle: the indicated replacement should be performed in the first-order equation. It is precisely for this reason that in equation (32.6) there appeared extra terms that would not have arisen had the substitution been made directly in the second-order equation.

Among the stationary solutions of the Dirac equation in an external field there may exist states belonging to both the continuous and the discrete spectrum. As in the non-relativistic theory, the continuous spectrum corresponds to infinite motion, in which the particle can be found at arbitrarily large distances where it may be regarded as free. Since the eigenvalues of the free-particle Hamiltonian are $\pm \sqrt{\mathbf{p}^2 + m^2}$, it is clear that the continuous spectrum of eigenvalues lies at $\varepsilon > m$ and at $\varepsilon \leq -m$. For $-m < \varepsilon < m$ the particle cannot escape to infinity, so that the motion is finite and the state belongs to the discrete spectrum.

As for free particles, so also for particles in an external field the wave functions with “positive frequency” ($\varepsilon > 0$) and “negative frequency” ($\varepsilon < 0$) enter the scheme of second quantisation in a definite way. For a particle in an external field this scheme is naturally generalised by replacing the plane waves in the formulae (25.1) with the correspondingly normalised eigenfunctions of the Dirac equation $\psi_n^{(+)}$ and $\psi_n^{(-)}$, belonging to the positive ($\varepsilon_n^{(+)}$) and negative ($-\varepsilon_n^{(-)}$) frequencies:

$$\begin{aligned}\widehat{\psi} &= \sum_n \{ \widehat{a}_n \psi_n^{(+)} \exp(-i\varepsilon_n^{(+)} t) + \widehat{b}_n^+ \psi_n^{(-)} \exp(i\varepsilon_n^{(-)} t) \}, \\ \widehat{\bar{\psi}} &= \sum_n \{ \widehat{a}_n^+ \bar{\psi}_n^{(+)} \exp(i\varepsilon_n^{(+)} t) + \widehat{b}_n \bar{\psi}_n^{(-)} \exp(-i\varepsilon_n^{(-)} t) \}.\end{aligned}\quad (32.9)$$

One must bear in mind here that, as the potential well deepens, the energy levels may cross the boundary $\varepsilon = 0$, i.e. the energies that were

positive may become negative (or, for a potential of the opposite sign, initially negative energies may become positive). Nonetheless, on grounds of continuity one must continue to regard these levels as electronic (rather than positronic). In other words, an electron includes all states which, in the limit of an infinitely slow removal of the field, tend to the positive boundary of the continuous spectrum ($\varepsilon = m$).

Although the Dirac equation for an electron in an external field does, as stated above, permit the solution of a wide class of quantum-electrodynamic problems, one must at the same time stress that the applicability of the external-field concept in relativistic theory is itself limited. This limitation is connected with the spontaneous creation of electron–positron pairs that arises in sufficiently strong fields (see below, §§ 35, 36).

We shall not consider in this book the introduction of an external field into the wave equations of particles with spin other than $1/2$, since this has no direct physical relevance—the real particles with such spins are hadrons, and their electromagnetic interactions cannot be described by wave equations. In this connection it should be noted that for spin-0 particles the wave equation has complex eigenvalues (with imaginary parts of both signs) in a sufficiently deep potential well. The wave equation for a spin-3/2 particle leads to a violation of causality, manifested in the appearance of solutions that propagate faster than light.

Problem

Determine the energy levels of an electron in a constant magnetic field.

Solution. Vector potential: $A_x = A_z = 0$, $A_y = Hx$ (the field H directed along the z -axis). The quantities that are conserved (along with the energy) are the components p_y and p_z of the generalised momentum.

We use the second-order equation for the auxiliary function φ (see (32.8)) and assume that φ is an eigenfunction of the operator Σ_z (with eigenvalue $\sigma = \pm 1$), as well as of the operators $\widehat{p}_y$ and $\widehat{p}_z$. The equation for φ then takes the form

$$\left\{ -\frac{d^2}{dx^2} + (eHx - p_y)^2 - eH\sigma \right\} \varphi = (\varepsilon^2 - m^2 - p_z^2) \varphi.$$

This equation is identical in form to the Schrödinger equation for a linear harmonic oscillator. Its eigenvalues are given by the formula

$$\varepsilon^2 - m^2 - p_z^2 = |e|H(2n+1) - eH\sigma, \quad n = 0, 1, 2, \dots$$

(cf. III, § 112). Note that the wave function ψ , which should be determined from φ by formula (32.8), is not an eigenfunction of the operator Σ_z , in accordance with the fact that for a moving particle the spin is not a conserved quantity.

§ 33. Expansion in powers of $1/c$

We saw (see § 21) that in the non-relativistic limit ($v \rightarrow 0$) the two components (χ) of the bispinor $\psi = \begin{pmatrix} \varphi \\ \chi \end{pmatrix}$ vanish. At low electron velocities, therefore, $\chi \ll \varphi$, and it becomes possible to derive an approximate equation involving only the two-component quantity φ , by formally expanding the wave function in powers of $1/c$ ³.

Starting from the Dirac equation for an electron in an external field written in the form

$$i\hbar \frac{\partial \psi}{\partial t} = \left\{ c\alpha \left(\hat{\mathbf{p}} - \frac{e}{c} \mathbf{A} \right) + \beta mc^2 + e\Phi \right\} \psi. \quad (33.1)$$

The relativistic energy of the particle includes the rest energy mc^2 . To pass to the non-relativistic approximation this must be removed, which we do by introducing in place of ψ a function ψ' defined by

$$\psi = \psi' e^{-imc^2 t / \hbar}.$$

Then

$$\left(i\hbar \frac{\partial}{\partial t} + mc^2 \right) \psi' = \left\{ c\alpha \left(\hat{\mathbf{p}} - \frac{e}{c} \mathbf{A} \right) + \beta mc^2 + e\Phi \right\} \psi'.$$

Writing $\psi' = \begin{pmatrix} \varphi' \\ \chi' \end{pmatrix}$ yields the system of equations:

$$\left(i\hbar \frac{\partial}{\partial t} - e\Phi \right) \varphi' = c\sigma \left(\hat{\mathbf{p}} - \frac{e}{c} \mathbf{A} \right) \chi', \quad (33.2)$$

$$\left(i\hbar \frac{\partial}{\partial t} - e\Phi + 2mc^2 \right) \chi' = c\sigma \left(\hat{\mathbf{p}} - \frac{e}{c} \mathbf{A} \right) \varphi' \quad (33.3)$$

(hereafter we drop the primes on φ and χ , since this will cause no confusion given that in this section we work exclusively with the transformed function ψ').

In the leading approximation we retain only the term $2mc^2\chi$ on the left of equation (33.3) and obtain

$$\chi = \frac{1}{2mc} \sigma \left(\hat{\mathbf{p}} - \frac{e}{c} \mathbf{A} \right) \varphi \quad (33.4)$$

(note that $\chi \sim \varphi/c$). Substituting this expression into (33.2) gives

$$\left(i\hbar \frac{\partial}{\partial t} - e\Phi \right) \varphi = \frac{1}{2m} \left(\sigma \left(\hat{\mathbf{p}} - \frac{e}{c} \mathbf{A} \right) \right)^2 \varphi.$$

³In this section we use ordinary units.

For the Pauli matrices one has the identity

$$(\sigma \mathbf{a})(\sigma \mathbf{b}) = \mathbf{a} \cdot \mathbf{b} + i\sigma \cdot [\mathbf{a} \times \mathbf{b}], \quad (33.5)$$

where $\mathbf{a}$ and $\mathbf{b}$ are arbitrary vectors (see (20.9)). Here $\mathbf{a} = \mathbf{b} = \hat{\mathbf{p}} - \frac{e}{c}\mathbf{A}$, but the vector product $[\mathbf{a} \times \mathbf{b}]$ does not vanish because $\hat{\mathbf{p}}$ and $\mathbf{A}$ do not commute:

$$\left[\left(\hat{\mathbf{p}} - \frac{e}{c}\mathbf{A} \right) \times \left(\hat{\mathbf{p}} - \frac{e}{c}\mathbf{A} \right) \right] \varphi = i \frac{e\hbar}{c} \{ [\mathbf{A} \cdot \nabla] + [\nabla \cdot \mathbf{A}] \} \varphi = i \frac{e\hbar}{c} \text{rot } \mathbf{A} \cdot \varphi.$$

Thus

$$\left(\sigma \left(\hat{\mathbf{p}} - \frac{e}{c}\mathbf{A} \right) \right)^2 = \left(\hat{\mathbf{p}} - \frac{e}{c}\mathbf{A} \right)^2 - \frac{e\hbar}{c} \sigma \cdot \mathbf{H} \quad (33.6)$$

(where $\mathbf{H} = \text{rot } \mathbf{A}$ is the magnetic field), and for φ we obtain the equation

$$i\hbar \frac{\partial \varphi}{\partial t} = \hat{H}\varphi = \left[\frac{1}{2m} \left(\hat{\mathbf{p}} - \frac{e}{c}\mathbf{A} \right)^2 + e\Phi - \frac{e\hbar}{2mc} \sigma \cdot \mathbf{H} \right] \varphi. \quad (33.7)$$

This is the so-called *Pauli equation*. It differs from the non-relativistic Schrödinger equation by the presence in the Hamiltonian of the last term, which represents the potential energy of a magnetic dipole in an external field (cf. III, § 111). In the leading ($1/c$) approximation, therefore, the electron behaves as a particle that possesses, in addition to its charge, a magnetic moment⁴:

$$\boldsymbol{\mu} = \frac{e}{mc} \hbar \mathbf{s}. \quad (33.8)$$

The gyromagnetic ratio (e/mc) is twice as large as it would be for a magnetic moment associated with orbital motion.

The density is $\rho = \psi^* \psi = \varphi^* \varphi + \chi^* \chi$. In the leading approximation the second term is to be dropped, so that $\rho = |\varphi|^2$, as it should be for the Schrödinger equation.

The current density is:

$$\mathbf{j} = c\psi^* \boldsymbol{\alpha} \psi = c(\varphi^* \boldsymbol{\sigma} \chi + \chi^* \boldsymbol{\sigma} \varphi).$$

Substituting (33.4):

$$\chi = \frac{1}{2mc} \boldsymbol{\sigma} \left(-i\hbar \nabla - \frac{e}{c}\mathbf{A} \right) \varphi, \quad \chi^* = \frac{1}{2mc} \left(i\hbar \nabla - \frac{e}{c}\mathbf{A} \right) \varphi^* \boldsymbol{\sigma},$$

and using the products involving two $\boldsymbol{\sigma}$ factors, which are rearranged with the help of (33.5) in the form

$$(\sigma \mathbf{a}) \boldsymbol{\sigma} = \mathbf{a} + i[\boldsymbol{\sigma} \times \mathbf{a}], \quad \boldsymbol{\sigma} (\sigma \mathbf{a}) = \mathbf{a} + i[\mathbf{a} \times \boldsymbol{\sigma}]. \quad (33.9)$$

The result is

$$\mathbf{j} = \frac{i\hbar}{2m} (\varphi \nabla \varphi^* - \varphi^* \nabla \varphi) - \frac{e}{mc} \mathbf{A} \varphi^* \varphi + \frac{\hbar}{2m} \text{rot}(\varphi^* \boldsymbol{\sigma} \varphi), \quad (33.10)$$

⁴This remarkable result was obtained by Dirac in 1928. The two-component wave function satisfying equation (33.7) had been introduced by Pauli (1927) even before Dirac discovered his equation.

in agreement with expression (115.4) (see III) from the non-relativistic theory.

Let us now find the second approximation, continuing the expansion to terms $\sim 1/c^2$ ⁵. We shall assume that only an electric field is present ($\mathbf{A} = 0$).

First of all, we note that with the inclusion of terms $\sim 1/c^2$ the density

$$\rho = |\varphi|^2 + |\chi|^2 = |\varphi|^2 + \frac{\hbar^2}{4m^2c^2} |\boldsymbol{\sigma} \cdot \nabla \varphi|^2.$$

This expression differs from the Schrödinger form. To find (in the second approximation) a wave equation analogous to the Schrödinger equation, we must introduce in place of φ a different (two-component) function φ_{Sch} , for which the time-independent integral $\int |\varphi_{\text{Sch}}|^2 d^3x$ has the same form as in the Schrödinger equation.

To determine the required transformation we write the condition

$$\int \varphi_{\text{Sch}}^* \varphi_{\text{Sch}} d^3x = \int \left\{ \varphi^* \varphi + \frac{\hbar^2}{4m^2c^2} (\nabla \varphi^* \cdot \boldsymbol{\sigma})(\boldsymbol{\sigma} \cdot \nabla \varphi) \right\} d^3x$$

and integrate by parts:

$$\int (\nabla \varphi^* \cdot \boldsymbol{\sigma})(\boldsymbol{\sigma} \cdot \nabla) \varphi d^3x = - \int \varphi^* (\boldsymbol{\sigma} \cdot \nabla)(\boldsymbol{\sigma} \cdot \nabla) \varphi d^3x = - \int \varphi^* \Delta \varphi d^3x$$

(or the same with φ and φ^* interchanged). Thus

$$\int \varphi_{\text{Sch}}^* \varphi_{\text{Sch}} d^3x = \int \left\{ \varphi^* \varphi - \frac{\hbar^2}{8m^2c^2} (\varphi^* \Delta \varphi + \varphi \Delta \varphi^*) \right\} d^3x,$$

from which one sees that

$$\varphi_{\text{Sch}} = \left(1 + \frac{\hat{\mathbf{p}}^2}{8m^2c^2} \right) \varphi, \quad \varphi = \left(1 - \frac{\hat{\mathbf{p}}^2}{8m^2c^2} \right) \varphi_{\text{Sch}}. \quad (33.11)$$

For simplicity let us assume a stationary state, i.e. replace the operator $i\hbar\partial/\partial t$ by the energy ε (with the rest energy subtracted). In the next (after (33.4)) approximation we have from (33.3):

$$\chi = \frac{1}{2mc} \left(1 - \frac{\varepsilon - e\Phi}{2mc^2} \right) (\boldsymbol{\sigma} \cdot \hat{\mathbf{p}}) \varphi.$$

This expression is to be substituted into (33.2), after which we replace φ by φ_{Sch} according to (33.11), dropping throughout all terms of order higher than $1/c^2$. A straightforward computation yields the

equation $\varepsilon \varphi_{\text{Sch}} = \hat{H} \varphi_{\text{Sch}}$, with the Hamiltonian

$$\hat{H} = \frac{\hat{\mathbf{p}}^2}{2m} + e\Phi - \frac{\hat{\mathbf{p}}^4}{8m^3c^2} + \frac{e}{4m^2c^2} (\boldsymbol{\sigma} \cdot \hat{\mathbf{p}}) \Phi (\boldsymbol{\sigma} \cdot \hat{\mathbf{p}}) - \frac{1}{2} (\hat{\mathbf{p}}^2 \Phi + \Phi \hat{\mathbf{p}}^2). \quad (33.12)$$

The expression in braces is rearranged using the formulae

$$(\boldsymbol{\sigma} \cdot \hat{\mathbf{p}}) \Phi (\boldsymbol{\sigma} \cdot \hat{\mathbf{p}}) + (\boldsymbol{\sigma} \cdot \hat{\mathbf{p}} \Phi) (\boldsymbol{\sigma} \cdot \hat{\mathbf{p}}) = \Phi \hat{\mathbf{p}}^2 + i\hbar (\boldsymbol{\sigma} \cdot \mathbf{E}) (\boldsymbol{\sigma} \cdot \hat{\mathbf{p}}),$$

⁵Here we follow the method of V. B. Berestetskii and L. D. Landau (1949).

$$\hat{\mathbf{p}}^2\Phi - \Phi\hat{\mathbf{p}}^2 = -\hbar^2\Delta\Phi + 2i\hbar\mathbf{E} \cdot \hat{\mathbf{p}},$$

where $\mathbf{E} = -\nabla\Phi$ is the electric field. The final expression for the Hamiltonian is:

$$\hat{H} = \frac{\hat{\mathbf{p}}^2}{2m} + e\Phi - \frac{\hat{\mathbf{p}}^4}{8m^3c^2} - \frac{e\hbar}{4m^2c^2}\boldsymbol{\sigma} \cdot [\mathbf{E} \times \hat{\mathbf{p}}] - \frac{e\hbar^2}{8m^2c^2}\text{div } \mathbf{E}. \quad (33.12)$$

The last three terms are the sought-for corrections of order $1/c^2$. The first of them is a consequence of the relativistic dependence of kinetic energy on momentum (the expansion of $\sqrt{\hat{\mathbf{p}}^2 + m^2c^2} - mc^2$). The second, which may be called the *spin-orbit interaction* energy, represents the interaction of the magnetic moment of a moving charge with the electric field⁶. The last term differs from zero only at those points where the field-producing charges are located; for the Coulomb field of a point charge Ze : $\Delta\Phi = -4\pi Ze\delta(\mathbf{r})$ (C. G. Darwin, 1928).

If the electric field is centrally symmetric,

$$\mathbf{E} = -\frac{\mathbf{r}}{r} \frac{d\Phi}{dr}$$

and the spin-orbit interaction operator can be written as

$$\frac{e\hbar}{4m^2c^2r}\boldsymbol{\sigma} \cdot [\mathbf{r} \times \hat{\mathbf{p}}] \frac{d\Phi}{dr} = \frac{\hbar^2}{2m^2c^2r} \frac{dU}{dr} \hat{\mathbf{I}} \cdot \hat{\mathbf{s}}. \quad (33.13)$$

Here $\hat{\mathbf{I}}$ is the orbital angular momentum operator, $\hat{\mathbf{s}} = \frac{1}{2}\boldsymbol{\sigma}$ is the electron spin operator, and $U = e\Phi$ is the potential energy of the electron in the field.

§ 34. Fine structure of hydrogen atom levels

Let us determine the relativistic corrections to the energy levels of the hydrogen atom—an electron in the Coulomb field of a stationary nucleus⁷. The velocity of the electron in the hydrogen atom $v/c \sim \alpha \ll 1$. Therefore the desired corrections can be calculated by perturbation theory—as an average over the unperturbed state (i.e. the non-relativistic wave function) of the relativistic terms in the approximate Hamiltonian (33.12). For somewhat greater generality we set the nuclear charge equal to Ze , assuming, however, that $Z\alpha \ll 1$.

The field of the nucleus $\mathbf{E} = Ze\mathbf{r}/r^3$, and its potential satisfies the equation $\Delta\Phi = -4\pi Ze\delta(\mathbf{r})$. Substituting this into (33.12) (the last three terms), and taking into account the negativity of the electron's charge, we obtain the perturbation operator:

$$\hat{V} = -\frac{\hat{\mathbf{p}}^4}{8m^3} + \frac{Z\alpha}{2r^3m^2}\hat{\mathbf{I}} \cdot \hat{\mathbf{s}} + \frac{Z\alpha\pi}{2m^2}\delta(\mathbf{r}). \quad (34.1)$$

⁶Introducing the magnetic moment (33.8) and the velocity $\mathbf{v} = \mathbf{p}/m$, one can write this energy as $-\frac{1}{2}\boldsymbol{\mu} \cdot [\mathbf{E} \times \mathbf{v}]$. At first sight this result may seem surprising, since on passing to the rest frame of the particle the magnetic field $\mathbf{H} = \frac{1}{c}[\mathbf{E} \times \mathbf{v}]$ arises, relative to which the magnetic moment should carry energy $-\boldsymbol{\mu} \cdot \mathbf{H}$. In reality the factor $1/2$ ("Thomas precession", L. Thomas, 1926) is connected with the general requirements of relativistic invariance in conjunction with the specific properties of the electron as a "spinor" particle with its associated value of the gyromagnetic ratio (see § 41).

⁷The effect of nuclear motion on the values of these corrections is of higher-order smallness, which does not interest us here.

Since according to the non-relativistic Schrödinger equation

$$\hat{\mathbf{p}}^2 \psi = 2m \left(\varepsilon_0 + \frac{Z\alpha}{r} \right) \psi$$

($\varepsilon_0 = -mZ^2\alpha^2/2n^2$ is the unperturbed level, n the principal quantum number), the mean value

$$\overline{\mathbf{p}^4} = 4m^2 \overline{\left(\varepsilon_0 + \frac{Z\alpha}{r} \right)^2}.$$

This quantity, as well as the mean value of the second term in (34.1), is computed using the formulae (see III, § 36)

$$\overline{r^{-1}} = \frac{m\alpha Z}{n^2}, \quad \overline{r^{-2}} = \frac{(m\alpha Z)^2}{n^3(l + \frac{1}{2})}, \quad \overline{r^{-3}} = \frac{(m\alpha Z)^3}{n^3 l(l + \frac{1}{2})(l + 1)} \quad (34.2)$$

(the last formula applies to $l \neq 0$); the eigenvalue

$$\mathbf{l} \cdot \mathbf{s} = \begin{cases} \frac{1}{2}[j(j+1) - l(l+1) - \frac{3}{4}], & l \neq 0, \\ 0, & l = 0. \end{cases}$$

Finally, the averaging of the third term is performed using the formula

$$|\psi(0)|^2 = \begin{cases} \frac{1}{\pi} \left(\frac{Z\alpha m}{n} \right)^3, & l = 0, \\ 0, & l \neq 0. \end{cases} \quad (34.3)$$

The result of a simple calculation using the formulae written above can be presented in all cases (for all j and l) in the form

$$\Delta\varepsilon = -\frac{m(Z\alpha)^4}{2n^3} \left(\frac{1}{j + \frac{1}{2}} - \frac{3}{4n} \right). \quad (34.4)$$

Formula (34.4) gives the desired relativistic correction to the energy of the hydrogen levels—the fine-structure energy⁸. Recall that in the non-relativistic theory degeneracy occurs both with respect to the direction of spin and the Coulomb degeneracy with respect to l . The fine structure (spin–orbit interaction) removes this degeneracy, but not completely: there remain doubly degenerate levels with the same n, j , but different $l = j \pm \frac{1}{2}$ (non-degenerate turn out to be only the levels with the largest possible $j = j_{\max} = l_{\max} + \frac{1}{2} = n - \frac{1}{2}$ for a given n). Thus the sequence of hydrogen levels with fine structure taken into account is as follows:

$$\begin{aligned} &1s_{1/2}; \\ &2s_{1/2}, \quad 2p_{1/2}, \quad 2p_{3/2}; \\ &3s_{1/2}, \quad 3p_{1/2}, \quad 3p_{3/2}, \quad 3d_{3/2}, \quad 3d_{5/2}. \end{aligned}$$

⁸This formula (and the more exact formula (36.10)) was obtained by A. Sommerfeld from the old Bohr theory even before the creation of quantum mechanics.

A level with principal quantum number n splits into n components of fine structure.

Recall that in non-relativistic mechanics the “accidental” degeneracy of energy levels in the Coulomb field is connected with the existence of a conservation law specific to this field: the quantity $\mathbf{A}$ is conserved, whose operator is

$$\hat{\mathbf{A}} = \frac{\mathbf{r}}{r} + \frac{1}{2m\alpha} \{[\hat{\mathbf{l}}\hat{\mathbf{p}}] - [\hat{\mathbf{p}}\hat{\mathbf{l}}]\}$$

(see III, (36.30)). A specific conservation law is connected with, and persists in, the relativistic case as a twofold degeneracy: the Hamiltonian of the Dirac equation $\hat{H} = \alpha\hat{\mathbf{p}} + \beta m - e^2/r$ commutes with the operator

$$\hat{\Gamma} = \frac{\mathbf{r}}{r} \cdot \boldsymbol{\Sigma} + \frac{i}{m\alpha} \beta (\boldsymbol{\Sigma} \cdot \hat{\mathbf{l}} + 1) \gamma^5 (\hat{H} - m\beta)$$

(*M. H. Johnson, B. A. Lippmann, 1950*). In the non-relativistic limit this operator $\hat{\Gamma} \rightarrow \boldsymbol{\Sigma}\hat{\mathbf{A}}$.

We shall see later (§ 123) that this remaining degeneracy is lifted by so-called radiative corrections (the *Lamb shift*), not accounted for by the Dirac equation for the single-electron problem.

Looking ahead, let us note here that the order of magnitude of these corrections is $\sim mZ^4\alpha^5 \ln(1/\alpha)$. The second-order correction due to the spin-orbit interaction was $\sim m(Z\alpha)^6$, so that its ratio to the radiative corrections is $\sim Z^2\alpha/\ln(1/\alpha)$. For hydrogen ($Z = 1$) this ratio is quite small, and therefore the problem of an exact solution of the Dirac equation in this case is of no practical interest. This problem may, however, have meaning for the energy levels of an electron in the field of a nucleus with large Z (see § 36).

§ 35. Motion in a centrally symmetric field

Consider the motion of an electron in a centrally symmetric electric field. Since angular momentum and parity are conserved in motion in a central field (relative to the centre of the field, chosen as the coordinate origin), everything said in § 24 about the angular dependence of wave functions of spherical waves of free particles applies to this motion as well. Only the radial functions change. Accordingly we seek the wave function of stationary states (in the standard representation) in the form

$$\varphi = \begin{pmatrix} \varphi \\ \chi \end{pmatrix} = \begin{pmatrix} f(r) \Omega_{jlm} \\ (-1)^{(1+l-l')/2} g(r) \Omega_{jl'm} \end{pmatrix}, \quad (35.1)$$

where $l = j \pm \frac{1}{2}$, $l' = 2j - l$, and the factor -1 is introduced to simplify the subsequent formulae.

The Dirac equation in the standard representation gives the following system of equations for φ and χ :

$$(\varepsilon - m - U)\varphi = \sigma\hat{\mathbf{p}}\chi, \quad (\varepsilon + m - U)\chi = \sigma\hat{\mathbf{p}}\varphi, \quad (35.2)$$

where $U(r) = e\Phi(r)$ is the potential energy of the electron in the field. The calculation of the result of substituting the expressions (35.1) into these equations reduces to computing the right-hand sides.

Expressing the spherical spinor $\Omega_{jl'm}$ through Ω_{jlm} according to

$$\Omega_{jl'm} = i^{l-l'} \left(\frac{\boldsymbol{\sigma} \cdot \mathbf{r}}{r} \right) \Omega_{jlm}$$

(see (24.8)), we write:

$$(\boldsymbol{\sigma} \hat{\mathbf{p}}) \chi = -i(\boldsymbol{\sigma} \nabla)(\boldsymbol{\sigma} \mathbf{r}) \frac{g}{r} \Omega_{jlm}.$$

Transforming now the product $(\boldsymbol{\sigma} \hat{\mathbf{p}})(\boldsymbol{\sigma} \mathbf{r})$ with the aid of formula (33.5), after expanding the vector operations we find

$$\begin{aligned} (\boldsymbol{\sigma} \hat{\mathbf{p}}) \chi &= -i\{\hat{p}_r + i\boldsymbol{\sigma}[\hat{\mathbf{p}} \hat{\mathbf{r}}]\} \frac{g}{r} \Omega_{jlm} = \\ &= \{-\operatorname{div} \mathbf{r} - (\mathbf{r} \nabla) - \boldsymbol{\sigma}[\mathbf{r} \hat{\mathbf{r}}]\} \frac{g}{r} \Omega_{jlm} = -\left\{g' + \frac{2}{r}g + \frac{g}{r} \boldsymbol{\sigma} \cdot \hat{\mathbf{I}}\right\} \Omega_{jlm}, \end{aligned}$$

where $\hat{\mathbf{I}} = [\mathbf{r} \hat{\mathbf{p}}]$ is the orbital angular momentum operator; the prime denotes differentiation with respect to r . The eigenvalues of the product $\boldsymbol{\sigma} \cdot \hat{\mathbf{I}} = 2\hat{\mathbf{l}} \cdot \hat{\mathbf{s}}$ are

$$2\hat{\mathbf{l}} \cdot \hat{\mathbf{s}} = \mathbf{j}^2 - \mathbf{l}^2 - \mathbf{s}^2 = j(j+1) - l(l+1) - \frac{3}{4} = \begin{cases} j - \frac{1}{2}, & l = j - \frac{1}{2}, \\ -j - \frac{3}{2}, & l = j + \frac{1}{2}. \end{cases}$$

For uniformity of notation in both cases ($l = j \pm \frac{1}{2}$) it is convenient to introduce the notation

$$\kappa = \begin{cases} -(j + \frac{1}{2}) = -(l + 1), & j = l + \frac{1}{2}, \\ +(j + \frac{1}{2}) = l, & j = l - \frac{1}{2}. \end{cases} \quad (35.3)$$

The number κ takes all integer values excluding zero (positive numbers correspond to $j = l - \frac{1}{2}$, and negative ones to $j = l + \frac{1}{2}$). Then $\mathbf{l} \cdot \boldsymbol{\sigma} = -(1 + \kappa)$, so that

$$(\boldsymbol{\sigma} \hat{\mathbf{p}}) \chi = -\left(g' + \frac{1 - \kappa}{r}g\right) \Omega_{jlm}.$$

On substituting this expression into the first of equations (35.2), the spherical spinor Ω_{jlm} cancels on both sides of the equation. Proceeding analogously with the second equation, we obtain the following system for the radial functions:

$$\begin{aligned} f' + \frac{1 + \kappa}{r}f - (\varepsilon + m - U)g &= 0, \\ g' + \frac{1 - \kappa}{r}g + (\varepsilon - m - U)f &= 0, \end{aligned} \quad (35.4)$$

or

$$\begin{aligned} (fr)' + \frac{\kappa}{r}(fr) - (\varepsilon + m - U)gr &= 0, \\ (gr)' - \frac{\kappa}{r}(gr) + (\varepsilon - m - U)fr &= 0. \end{aligned} \quad (35.5)$$

Let us investigate the behaviour of f and g at small distances, assuming that the field $U(r)$ grows as $r \rightarrow 0$ faster than $1/r$. Then in the region of small r equations (35.4) take the form

$$f' + Ug = 0, \quad g' - Uf = 0.$$

They have real solutions of the form

$$f = \text{const} \cdot \sin\left(\int U dr + \delta\right), \quad g = \text{const} \cdot \cos\left(\int U dr + \delta\right), \quad (35.6)$$

where δ is an arbitrary constant. These functions oscillate as $r \rightarrow 0$, without tending to any limit. It is easy to see that this situation corresponds in the non-relativistic theory to “falling” of the particle to the centre.

First of all we note that the region of small distances imposes in this case no restrictions on the choice of solution: the condition at $r = 0$ for the oscillating function is absent, and the choice of the constant δ remains arbitrary (the correct behaviour of the wave function at large r can be achieved for any ε by an appropriate choice of δ). This indeterminacy can be removed by considering the singular (at $r = 0$) potential as the limit $r_0 \rightarrow 0$ of the potential “cut off” at some small r_0 (i.e. equal to $U(r)$ for $r > r_0$ and $U(r_0)$ for $r < r_0$). With a finite r_0 one obtains, of course, a definite system of energy levels. However, the energy of the ground state tends to $-\infty$ as $r_0 \rightarrow 0$.

In the non-relativistic theory this is precisely the “falling” to the centre, since a particle at a deep level is localised in a small region around $r = 0$. In the relativistic theory, however, such a situation is altogether inadmissible, since it signifies the instability of the system with respect to the spontaneous creation of electron–positron pairs. Indeed, if in vacuum the energy needed to create such a pair exceeds $2m$, in the field a smaller energy already suffices. If there exists a bound state of the electron with energy $\varepsilon < m$, pair creation is possible at the cost of only the energy $\varepsilon + m < 2m$, with a free positron created and the electron in the bound state. If the energy of the bound state $\varepsilon < -m$, such a field can create positrons (with energy $-\varepsilon > m$) spontaneously, without any input of energy from an external source. In the field being considered, as $r_0 \rightarrow 0$ there is an infinite number of such “anomalous” levels with $\varepsilon < -m$. Therefore a potential $\Phi(r)$ growing as $r \rightarrow 0$ faster than $1/r$ cannot at all be considered in the Dirac theory. Let us stress that this applies to potentials of either sign. The “falling” occurs, of course, only in the case of attraction, since the sign of $U = e\Phi$ depends also on the sign of the charge; in one case the electronic, and in the other the positronic levels behave anomalously; in the second case the field creates free electrons.

Let us consider further the behaviour of wave functions at large distances. If the field $U(r)$ falls off sufficiently rapidly at $r \rightarrow \infty$, then in determining the asymptotic form of the wave functions at large distances the field can be completely neglected in the equations. For $\varepsilon > m$, i.e. in the continuous-spectrum region, we return then to the free-motion equation, so that the asymptotic form of the wave functions (spherical waves) differs from that for a free particle only by the appearance of additional “phase shifts”, whose values are determined by the form of the field at close distances⁹. These shifts depend on the values of j and l , or, equivalently, on the number κ (and also, of course, on the energy ε). Denoting them by δ_κ and using the expression for the free spherical

⁹Cf. III, § 33. As in the non-relativistic theory, $U(r)$ must fall off faster than $1/r$. The case $U \sim 1/r$ will be considered separately in § 36.

wave (24.7), we can immediately write the desired asymptotic formula

$$\psi \approx \frac{2}{r} \frac{1}{\sqrt{2\varepsilon}} \left(\begin{array}{l} \sqrt{\varepsilon + m} \Omega_{jlm} \sin\left(pr - \frac{\pi l}{2} + \delta_\kappa\right) \\ -\sqrt{\varepsilon - m} \Omega_{jl'm} \sin\left(pr - \frac{\pi l'}{2} + \delta_\kappa\right) \end{array} \right), \quad (35.7)$$

or, taking into account the definition (35.1):

$$\left. \begin{array}{l} f \\ g \end{array} \right\} = \frac{\sqrt{2}}{r} \sqrt{\frac{\varepsilon \pm m}{\varepsilon}} \sin\left(pr - \frac{\pi l}{2} + \delta_\kappa\right), \quad (35.8)$$

where $p = \sqrt{\varepsilon^2 - m^2}$. The overall coefficient here corresponds to normalisation of the radial functions according to (24.5).

The wave functions of the discrete spectrum ($\varepsilon < m$) decay exponentially at $r \rightarrow \infty$ according to the law

$$f = -\sqrt{\frac{m + \varepsilon}{m - \varepsilon}} g = \frac{A_0}{r} \exp\left(-r\sqrt{m^2 - \varepsilon^2}\right), \quad (35.9)$$

where A_0 is a constant.

As in the non-relativistic theory, the phase shifts δ_κ (more precisely, the quantities $e^{2i\delta_\kappa} - 1$) determine the scattering amplitude in the given field (this will be discussed in more detail in § 37). We shall not investigate here the analytic properties of these quantities (cf. III, § 128). We note only that $e^{2i\delta_\kappa}$ as a function of the energy still has poles at the points corresponding to the bound-state energy levels of the particle. The residue of the function $e^{2i\delta_\kappa}$ at such a pole is related in a definite way to the coefficient in the asymptotic expression of the corresponding wave function of the discrete spectrum. Let us find this relation, which generalises the non-relativistic formula (128.17) (see III). The necessary calculations are entirely analogous to those performed in III, § 128.

Differentiate equations (35.5) with respect to the energy:

$$\begin{aligned} \left(\frac{\partial r f}{\partial \varepsilon}\right)' + \frac{\kappa}{r} \frac{\partial r f}{\partial \varepsilon} - (\varepsilon + m - U) \frac{\partial r g}{\partial \varepsilon} &= r g, \\ \left(\frac{\partial r g}{\partial \varepsilon}\right)' - \frac{\kappa}{r} \frac{\partial r g}{\partial \varepsilon} + (\varepsilon - m - U) \frac{\partial r f}{\partial \varepsilon} &= -r f. \end{aligned}$$

Multiply these two equations respectively by rg and $-rf$, and the two equations (35.5) respectively by $-rg$ and rf ; then add all four equations term by term. After all cancellations we get

$$\frac{\partial}{\partial r} \left[r^2 \left(g \frac{\partial f}{\partial \varepsilon} - f \frac{\partial g}{\partial \varepsilon} \right) \right] = r^2 (f^2 + g^2).$$

Integrate this relation over r :

$$r^2 \left(g \frac{\partial f}{\partial \varepsilon} - f \frac{\partial g}{\partial \varepsilon} \right) = \int_0^r (f^2 + g^2) r^2 dr,$$

and then pass to the limit $r \rightarrow \infty$. By the normalisation condition the integral on the right-hand side becomes unity. On the left-hand side we take into account that in the asymptotic region the functions f and g are related by

$$rg = \frac{(rf)'}{\varepsilon + m},$$

obtained from (35.5) by neglecting the terms with U and $1/r$. As a result we get

$$\frac{1}{\varepsilon + m} \left[(rf)' \frac{\partial rf}{\partial \varepsilon} - rf \left(\frac{\partial rf}{\partial \varepsilon} \right)' \right] = 1. \quad (35.10)$$

This formula differs from the analogous non-relativistic formula (for the function χ) only by the coefficient ($\varepsilon + m$ instead of $2m$). There is therefore no need to repeat all the subsequent calculations, and we immediately give the final formula, valid near the point $\varepsilon = \varepsilon_0$ (ε_0 being the energy level):

$$e^{2i\delta_\kappa} = (-1)^l \frac{2A_0^2}{\varepsilon - \varepsilon_0} \sqrt{\frac{m - \varepsilon_0}{m + \varepsilon_0}}, \quad (35.11)$$

where A_0 is the coefficient in the asymptotic expression (35.9).

Problem

Find the limiting form of the wave function at small r in the field $U \sim r^{-s}$, $s < 1$.

Solution. For a free particle we have at small r : $f \sim r^l$, $g \sim r^{l'}$, so that for $l < l'$: $f \gg g$, and for $l > l'$: $f \ll g$. We make the assumption (justified by the result) that such a relation is preserved also in the field being considered. For $l < l'$ (i.e. $l = j - \frac{1}{2}$, $\kappa = -l - 1$) the g term in the first of equations (35.4) can be dropped, so that $f \sim r^l$ as before. From the second equation we then have $g \sim rfU$, i.e. $g \sim r^{l+1-s}$. The case $l > l'$ is treated analogously. As a result we find:

$$\begin{aligned} \text{for } l < l': \quad f &\sim r^l, & g &\sim r^{l'-s}; \\ \text{for } l > l': \quad f &\sim r^{l-s}, & g &\sim r^{l'}. \end{aligned}$$

Radiation

Scattering of Light

The Scattering Matrix

Invariant Perturbation Theory

Interaction of Electrons

Interaction of Electrons with Photons

Exact Propagators and Vertex Parts

Radiative Corrections

Asymptotic Formulae of Quantum Electrodynamics

Electrodynamics of Hadrons