

THEMATIC REVIEW

Metabolomics in diabetes research

Nele Friedrich

Institute for Clinical Chemistry and Laboratory Medicine, University of Greifswald, Ferdinand-Sauerbruch-Strasse, D-17475 Greifswald, Germany
(Correspondence should be addressed to N Friedrich; Email: nele.friedrich@uni-greifswald.de)

Abstract

Diabetes represents one of the most important global health problems because it is associated with a large economic burden on the health systems of many countries. Whereas the diagnosis and treatment of manifest diabetes have been well investigated, the identification of novel pathways or early biomarkers indicative of metabolic alterations or insulin resistance related to the development of diabetes is still in progress. Over half of the type 2 diabetes patients show manifestations of diabetes-related diseases, which highlight the need for early screening markers of diabetes. During the last decade, the rapidly growing research field of

metabolomics has introduced new insights into the pathology of diabetes as well as methods to predict disease onset and has revealed new biomarkers. Recent epidemiological studies first used metabolism to predict incident diabetes and revealed branched-chain and aromatic amino acids including isoleucine, leucine, valine, tyrosine and phenylalanine as highly significant predictors of future diabetes. This review summarises the current findings of metabolic research regarding diabetes in animal models and human investigations.

Journal of Endocrinology (2012) **215**, 29–42

Introduction

Diabetes mellitus, particularly type 2 diabetes (T2DM), represents one of the most significant global health problems because it is associated with a large economic burden on the health systems of many countries. The World Health Organization (WHO) reported that worldwide 171 million individuals were affected by diabetes in 2000, which is equivalent to a prevalence of 2.8%, and predicted an estimated future number of 366 million affected individuals in 2030, which would be equivalent to a diabetes prevalence of 4.4% (Wild *et al.* 2004). However, recent data from the International Diabetes Federation (IDF) revealed that this number has already been reached in 2011. The IDF expected an even higher number of 552 million affected persons in 2030. Due to the broad range of diabetes-related complications, including diabetic nephropathy, peripheral neuropathy and cardiovascular disease, diabetes is a major cause of both morbidity and mortality.

Diabetes mellitus is a chronic disease that is characterised by high blood glucose levels, which may be due either to the progressive failure of pancreatic β -cell function and consequently a lack of insulin production (type 1: T1DM) or to the

development of insulin resistance and subsequently the loss of β -cell function (T2DM). Approximately 90% of patients with diabetes have T2DM. T1DM is an autoimmune disease with a strong genetic component. Genetic susceptibility to T1DM has been intensively investigated, and the major histocompatibility complex was reported to be the main genetic determinant (Polychronakos & Li 2011, Noble & Erlich 2012). The predominant cause of T2DM is related to lifestyle factors including diet, insufficient physical activity, an overweight or obese state and stress. Furthermore, at least 36 genes, accounting for 10% of the total genetic component, have been significantly associated with an increased risk for T2DM (Herder & Roden 2011). Moreover, a further 18 genes were related to glucose and HbA1c levels as well as insulin resistance. Detailed reviews on the genetics of T1DM and T2DM have been presented elsewhere (Herder & Roden 2011, Polychronakos & Li 2011).

The diagnosis of diabetes is mainly based on the results of blood tests examining fasting plasma glucose or HbA1c levels (World Health Organization 2006). Additionally, the treatment of diabetes is linked to these measures, as the main goal of antidiabetic therapy is to reduce blood glucose and HbA1c levels via the administration of insulin (T1DM) or antidiabetic medication (T2DM). Furthermore, lifestyle changes, such as eating a more healthy diet, performing regular physical activity, achieving a normal body weight and smoking cessation, are recommended for diabetic patients.

This paper is one of three papers that form part of a thematic review section on Metabolomics. The Guest Editor for this section was Henri Wallaschofski, Ernst-Moritz-Arndt University, Greifswald, Germany.

Whereas the diagnosis and treatment of manifest diabetes have been thoroughly investigated, the identification of novel pathways or early biomarkers indicative of metabolic alterations or insulin resistance related to the development of T2DM is still underway. Data from the National Health and Nutrition Examination Survey (NHANES) showed that an estimated 57.9% of subjects with diagnosed diabetes are affected by one or more macro- or microvascular complications (American Association of Clinical Endocrinologists 2007), which highlights the need for early screening markers to monitor the development of T2DM. During the last decade, the rapidly growing research field of metabolomics has introduced new insights into the pathology of diabetes as well as methods to predict disease onset. In accordance with the other 'omics' techniques, such as genomics, transcriptomics or proteomics, the term metabolomics, which is synonymous with metabolomics or metabolic profiling, was first introduced by Nicholson *et al.* (1999). Metabolomics relates to measurements of the metabolome, which represents the entire collection of all small-molecule metabolites present in any biological organism (Oliver *et al.* 1998, Tweeddale *et al.* 1998). The advantages of metabolomics over other 'omics' technologies include its high level of sensitivity and its ability to enable the analysis of relatively few metabolites compared with the unwieldy number of corresponding genes or mRNA molecules. The comprehensive Human Metabolome Database (HMDB) contained ~7900 metabolite entries in January 2009. Another advantage of metabolomics is that metabolites are the final downstream products of the interaction between genes and influences like environmental factors, health behaviour or pharmaceutical interventions, and metabolite levels reflect the activity of metabolic pathways. Therefore, metabolomics enables the detection of short-term as well long-term physiological or pathological changes in cells, tissues or body fluids and represents a useful tool for biomarker detection.

The purpose of the present review is to summarise current metabolomics technologies and to provide an overview of the contribution of metabolomics to diabetes research.

Metabolomics technologies

The two main high-throughput metabolomics tools consist of nuclear magnetic resonance (NMR) spectroscopy and mass spectrometry (MS). Both methods enable the comprehensive investigation of metabolic profiles (Dunn *et al.* 2005, Hollywood *et al.* 2006, Lenz & Wilson 2007) and can provide complementary snapshots of the metabolome of body fluids such as plasma, urine or cerebrospinal fluid (Bictash *et al.* 2010).

NMR spectroscopy

NMR is a widely used spectroscopic technique for metabolomics that is based on the magnetic properties of the atomic nucleus (e.g. ^1H , ^{13}C or ^{31}P). This method was

first used for the analysis of body fluids in the 1980s (Nicholson *et al.* 1984, Iles *et al.* 1985, Bell *et al.* 1989). The behaviour of NMR active nuclei in a strong magnetic field provides information about the structural and chemical properties of a molecule. Due to the high abundance of the ^1H nucleus, ^1H -NMR spectroscopy has been heavily used to investigate biological fluids such as plasma and urine as well as tissues. Each separate signal in an ^1H -NMR spectrum corresponds to a particular compound. Exemplary ^1H -NMR spectra for human urine and plasma are shown in Fig. 1. Based on measurements of the following:

- i) chemical shift,
- ii) spin-spin coupling: neighbouring nuclei influence the effective magnetic field, which results in spin interaction. The so-called spin-spin coupling can cause splitting of the signal into two or more peaks,
- iii) relaxation: describes the return to equilibrium of net magnetisation and included two types of relaxation: spin-lattice (T_1 , also called longitudinal relaxation) and spin-spin (T_2 , also called transverse relaxation),
- iv) diffusion

the identification of single metabolites and absolute quantification are possible. Detailed information regarding NMR theory, its application and typical chemical shift values is available elsewhere (Blümich 2005). The application of NMR spectroscopy in metabolomics ranges from pharmaceutical studies (Lindon *et al.* 2007) to cardiovascular biomarker identification (Griffin *et al.* 2011, Rhee & Gerszten 2012).

Advantages of NMR spectroscopy include the non-destructive nature of the analysis, the robust and reproducible measurements and the minimal preparation requirements, as no separation or ionisation steps are necessary. However, in




Figure 1 Assigned high-resolution ^1H -NMR spectrum of a human plasma (600 MHz) and human urine (400 MHz) sample.

comparison to MS, the analytical sensitivity of NMR is relatively low, even if stronger magnetic fields are used to increase the analytical sensitivity.

Mass spectrometry

MS is a powerful tool for investigating molecular structure as well as for detecting and quantifying metabolites (Lenz & Wilson 2007). The first step in MS consists of the ionisation of the analyte, which is followed by the separation of the ions according to their mass-to-charge (m/z) ratio using an analyser with an electromagnetic field. In metabolomics, MS is often combined with other suitable methods for the analytical separation of compounds, including gas chromatography (GC) or liquid chromatography (LC), to achieve detection of distinct metabolite classes (Buscher *et al.* 2009). One study comparing GC, LC and capillary electrophoresis (CE) revealed that LC was the most robust method and enabled the broadest coverage of metabolite classes (Buscher *et al.* 2009). However, both GC–MS and LC–MS demonstrate high separation efficiency and are excellent tools for metabolic profiling. Exemplary MS spectrum for human urine is shown in Fig. 2. The application of MS in the metabolomics field ranges from studies of microorganisms and plants to biomarker detection.

The greatest advantage of the MS methods is their high level of sensitivity, although disadvantages arise from the destruction of the sample and the long sample preparation

time required. Detailed descriptions of MS methods have been provided elsewhere (de Hoffmann & Stroobant 2007).

Due to the limited overlap for metabolite detection and thus the complementary nature of MS and NMR spectroscopy, studies using a multiplatform approach may provide a more comprehensive understanding of metabolic alterations than studies using only one of these tools (Williams *et al.* 2006a).

Experimental studies applied to glucose metabolism and diabetes

Dietary-induced obesity and insulin resistance

A series of experimental studies have been conducted using dietary-induced obesity or insulin resistance rodent models to investigate the metabolic profiles of urine (Kim *et al.* 2009), blood (Shearer *et al.* 2008, Li *et al.* 2010a) or tissue (Li *et al.* 2010a, Lin *et al.* 2011), and these results have led to new insights into the development of diabetes. Dietary-induced obesity models have the advantage of being more similar to the development of human obesity in comparison to genetic models, and, as a result, these models mirror the progression of insulin resistance and diabetes after a prolonged period of development (Fearnside *et al.* 2008). High-fat-fed C57BL/6J mice become obese and insulin resistant and demonstrate different serum ^1H -NMR-based metabolite concentrations in comparison with chow-fed mice (Shearer *et al.* 2008). Whereas the citrate concentration was higher in high-fat-fed mice than in chow-fed mice, the concentrations of glycine, lysine, suberate, acetate, leucine, valine or trimethylamine-*N*-oxide were significantly lower in high-fat-fed mice. Furthermore, dietary-induced insulin resistance could be predicted according to various metabolite levels, specifically those of lysine, glycine, citrate, leucine, suberate and acetate, and these metabolite levels could also be used to discriminate between chow- and high-fat-fed animals. Additionally, the urinary metabolic profiles of high-fat-fed rats were significantly different from those of normal-diet-fed rats (Kim *et al.* 2009). ^1H -NMR spectroscopy also revealed diet-related variations in the levels of betaine, taurine, acetone/ acetoacetate, phenylacetylglutamine, pyruvate, lactate and citrate. MS-based studies can also detect additional diet-induced changes in metabolites (Li *et al.* 2010a, Lin *et al.* 2011); one study investigated the diet-induced inhibition of insulin in the liver tissue and plasma of wild-type and glycerol-3-phosphate acyltransferase 1-deficient mice, which remain insulin sensitive independent of their diet (Li *et al.* 2010a). Assuming that metabolic changes identified in insulin-resistant WT mice are specifically related to hepatic insulin resistance and may therefore indicate a causative pathway, these authors demonstrated alterations in the concentrations of 43 liver and 19 plasma metabolites. The identified increases in urea cycle-related metabolites, such as citrulline, aspartate or *N*-acetylglutamate, were indicative of




Figure 2 GC–MS analysis of organic acids in human urine. Above – total ion current (TIC) chromatogram: (1) glycolic, (2) p-cresol, (3) 3-hydroxyisovaleric, (4) urea, (5) succinate, (6) hydroxyphenylacetic, (7) aconitic, (8) citric, and (9) 4-(3-hydroxyprop-1-enyl)phenol. Below – mass spectrum at retention time 44.727 min.

early up-regulation of the urea cycle, whereas the altered levels of liver metabolites suggested the existence of variations in glucose metabolism (1,5-anhydroglucitol decrease), bile acid metabolism (taurocholate decrease) and pyrimidine metabolism (2'-deoxyuridine increase). Moreover, the increase in pyrimidine metabolites and the decreases in bradykinin, kynurenine and α -ketoglutarate concentrations were also confirmed in the plasma. A separate MS study extended this approach of diet-induced insulin resistance to include a metabolic oral glucose tolerance test (OGTT) and additionally examined liver, brain and skeletal tissues (Lin *et al.* 2011). These MS data enabled the authors to discriminate between both the 120- and 0-min time points for both standard-fed (SD) and high-fructose-fed (HFRD) rats, and these data also identified specific metabolic effects in insulin-resistant rats. As expected, insulin administration-related up-regulation of lysophosphatidylcholines (Lin *et al.* 2011) was observed in SD rats but not in HFRD rats. However, the levels of the branched-chain amino acids (BCAA) proline, tryptophan and methionine were decreased in HFRD rats at 120 min but were unchanged in SD rats, and opposite effects were observed for the amino acids leucine and isoleucine, which had previously been shown to be related to insulin sensitivity (Shaham *et al.* 2008) and were present at lower levels in HFRD rats. By comparing SD and HFRD rats at the 0-min time point, differences were identified for various compounds, including phospholipids, amino acids, bile acids, fatty acids and metabolites. Moreover, regarding purine metabolism and the Krebs cycle, a complex metabolic perturbation in HFRD rats was observed. Increased levels of phospholipids and fatty acid were also found in high-fat-fed mice in combination with lower levels of betaine, carnitine and acylcarnitines, which are metabolites involved in lipid metabolism (Kim *et al.* 2011). In liver and skeletal muscle tissue, a high-fructose diet leads to oxidative stress, elevated levels of amino acids and alterations in fatty acid biosynthesis, whereas this type of diet is related to decreased amino acid levels and the up-regulation of purine biosynthesis in the cerebral cortex and hippocampus (Lin *et al.* 2011).

In general, the distinction between diet-related effects and obesity-related effects represents a common problem in dietary-induced diabetes models. One recent study on mice investigated the long- and short-term consequences of various types of diets and aimed to distinguish the specific effects of each diet from those of obesity in general (Duggan *et al.* 2011). The results revealed that diet has a major impact on the metabolic profiles measured by ^1H -NMR; whereas diet influenced metabolites related to energy and glucose metabolism, obesity mainly caused alterations in amino acids and large non-polar molecules.

Genetic rodent models of diabetes

Genetic T2DM models have several advantages over diet-induced models, including a short generation time, heritable traits and lower cost, although the 'natural' development of

diabetes over a prolonged period of time is lacking in these animals. The two most popular obesity/T2DM models include the *db/db* model and the obese Zucker (*fa/fa*) model, both are characterised by an autosomal recessive defect in the leptin receptor gene that results in obesity and subsequent insulin resistance (Chen & Wang 2005). Based on these models, differences in urinary (Williams *et al.* 2005a, 2006b, Salek *et al.* 2007, Gipson *et al.* 2008, Connor *et al.* 2010, Zhao *et al.* 2010a, Patterson *et al.* 2011), plasma (Major *et al.* 2006, Williams *et al.* 2006a) and tissue (Xu *et al.* 2009, Connor *et al.* 2010) metabolic profiles have been reported between affected and control rodents. Several studies have demonstrated profound alterations in metabolites involved in the tricarboxylic acid (TCA) cycle. The TCA cycle is an amphibolic pathway that occurs in the inner mitochondrial membrane and plays an important role in energy metabolism. The final products of fatty acid degradation and glycolysis are included in the TCA cycle, and TCA cycle intermediates are involved in amino acid synthesis and degradation as well as gluconeogenesis. Whereas Zucker rats (Williams *et al.* 2005a, 2006b, Zhao *et al.* 2010a) typically have decreased urinary concentrations of TCA metabolites, such as citrate, malate, fumarate, 2-ketoglutarate or succinate, the *db/db* mouse (Salek *et al.* 2007, Connor *et al.* 2010) has exhibited increased levels of TCA metabolites, and these changes are indicative of the down- and up-regulation of the TCA cycle, respectively. In a study with rhesus macaques, animals with T2DM demonstrated twofold higher levels of citrate compared with normal animals (Patterson *et al.* 2011). Additionally, Sprague-Dawley rats with T1DM induced by streptozotocin demonstrated higher levels of pyruvate, succinate and fumarate (Zhang *et al.* 2008). This study further showed strong associations between TCA cycle intermediates and components of glucose metabolism in normal rats, specifically between pyruvate, as the end product of glycolysis, and 2-oxoglutarate, fumarate or citrate. In diabetic rats, there was evidence for a disturbed balance between the TCA cycle and glucose metabolism, as the glucose levels were not associated with those of lactate or various TCA cycle intermediates. Thus, the concentrations of succinate were not correlated with those of 2-oxoglutarate or citrate (Zhang *et al.* 2008). Beside disturbances in intermediate correlations, the metabolite composition demonstrated a strong age-dependent variation. Williams *et al.* (2006b) reported that the urinary ratios of α -ketoglutarate and succinate to citrate were in favour of citrate at an age of 4 weeks in Zucker rats, although this finding was no longer apparent at 20 weeks. In *db/db* mice, a decrease in the concentration of both TCA and non-TCA metabolites was also reported with age (Salek *et al.* 2007). Regarding taurine, the urinary levels in control and Zucker rats were comparable at 8 weeks, whereas taurine was absent in 50% of the Zucker animals at 20 weeks (Williams *et al.* 2006b). Other metabolites involved in amino acid metabolism have been shown to be associated with T2DM. For example, amino acids, such as phenylalanine, valine, tryptophan, lysine and glutamic acid, and amino

acid metabolites, such as 2-hydroxyisobutyrate, 2-hydroxyisovalerate and kynurenic acid, were present at higher concentrations in the urine of Zucker rats (Salek *et al.* 2007, Gipson *et al.* 2008, Connor *et al.* 2010), *db/db* mice (Gipson *et al.* 2008, Connor *et al.* 2010) and monkeys (Patterson *et al.* 2011) than in control animals, although lower concentrations of these compounds have also been reported (Williams *et al.* 2006b, Salek *et al.* 2007). These results provide evidence for the complex perturbation of amino acid metabolism in diabetic disease. The largest portion of amino acid metabolism takes place in the liver, and although a broad range of amino acids are glucogenic and are used for hepatic gluconeogenesis, a smaller number of amino acids are ketogenic and are converted to ketone bodies. An additional rodent study (Mochida *et al.* 2011) on T1DM using the *Akita* mouse model confirmed these previous urine findings and demonstrated higher amino acid, BCAA, alanine, citrulline and proline levels in the plasma of T1DM rats indicative of disease state-dependent changes. As hyperglycaemia progressed, the differences regarding the mentioned amino acids and BCAA became more pronounced. Furthermore, these authors demonstrated a relation between increased blood glucose levels and increases in plasma levels of valine, leucine, isoleucine, BCAA and alanine (Mochida *et al.* 2011). In addition to insulin resistance, BCAA supplementation to the high-fat diet leads to chronic increased activation of mTOR in rats (Newgard *et al.* 2009), and overactivation of the mTOR/S6K1 pathway has been linked to the development of insulin resistance via β -cell adaptation to hyperglycaemia (Um *et al.* 2004, Khamzina *et al.* 2005, Fraenkel *et al.* 2008). Further details concerning the relation between amino acids and diabetes are provided in the human studies section of this review.

A series of rodent studies found increased urinary levels of fatty acids in Zucker rats (Williams *et al.* 2006b, Salek *et al.* 2007). Furthermore, higher levels of carnitine, an essential compound for the transport of fatty acids from the cytosol into the mitochondria and one that is related to T2DM (De Palo *et al.* 1981), have been observed in diabetic rodents (Williams *et al.* 2006b, Connor *et al.* 2010). A multiplatform study confirmed the enriched fatty acid metabolism of the *db/db* mouse by revealing increased transcript levels of fatty acid metabolism-associated carnitine palmitoyltransferase in the liver and higher urinary carnitine levels, measured using LC–MS (Gipson *et al.* 2008). Furthermore, increased carnitine levels with age were observed in *db/db* but not in control mice, and similar age-dependent increases in carnitine were also reported in non-diabetic rats (Williams *et al.* 2005b). Increased fatty acid metabolism results from a higher rate of lipolysis in adipose tissue and might exacerbate insulin resistance in liver and muscle tissue (Delarue & Magnan 2007). High fatty acid levels subsequently lead to a higher oxidation rate and therefore to the induction of ketogenesis. Thus, diabetic animals typically exhibit higher levels of ketone bodies, including β -hydroxybutyrate or acetone, compared with controls (Salek *et al.* 2007, Zhao *et al.* 2010a). Similar to changes in amino acid levels, increased levels of ketone bodies

have been associated with the diseased state as well as age (Salek *et al.* 2007). Taken together, the shift from euglycaemia towards hyperglycaemia is likely linked to pronounced metabolic perturbations and mitochondrial metabolic dysfunction. Beside the previously mentioned metabolites, a wide range of additional urine or tissue metabolites, e.g. choline, allantoin, glycine or betaine, has also been related to obesity or diabetes in different studies (Williams *et al.* 2005a, 2006b, Salek *et al.* 2007, Gipson *et al.* 2008, Zhang *et al.* 2008, Zhao *et al.* 2010a, Patterson *et al.* 2011).

Several studies have also detected differences in metabolites known to originate from the gut microflora. For example, hippurate, which is mainly produced via gut microbial metabolism (Nicholson *et al.* 2005), was shown to be elevated in *db/db* mice (Salek *et al.* 2007, Connor *et al.* 2010) but decreased in Zucker rats (Williams *et al.* 2006b, Salek *et al.* 2007, Zhao *et al.* 2010a), whereas microbiota-derived methylamines such as dimethylamine or trimethylamine-N-oxide were shown to be increased in both types of rodents (Salek *et al.* 2007, Gipson *et al.* 2008, Zhang *et al.* 2008, Connor *et al.* 2010, Zhao *et al.* 2010a). Reasons for this discrepancy may be related to differences in the composition of the intestinal microflora.

Human studies applied to glucose metabolism and diabetes

Glucose ingestion

The OGTT measures the body's ability to metabolise glucose and clear excess glucose from the bloodstream. Since the 1970s, the OGTT has been a standard diagnostic tool in diabetology. A 2 h OGTT is routinely performed in fasting patients; patients drink a beverage containing a specific amount of glucose according to their body weight and 2 h after the glucose load, the blood glucose concentration is measured and provides information on glucose tolerance. According to the WHO, in healthy individuals, the venous plasma glucose level should be below 7.8 mmol/l; values greater than this can be used to diagnose impaired glucose tolerance (7.8–11.1 mmol/l) or diabetes mellitus (≥ 11.1 mmol/l) (World Health Organization 2006).

Several studies have investigated changes in the metabolic profile in relation to glucose ingestion (Shaham *et al.* 2008, Zhao *et al.* 2009, Spiegel *et al.* 2010, Matysik *et al.* 2011). An investigation (Shaham *et al.* 2008) among participants in the Metabolic Abnormalities in College Students (MACS) study, who demonstrated normal glucose tolerance, revealed significant kinetic alterations in 21 out of 191 plasma metabolites, measured by LC–MS, in response to an OGTT. Eighteen of these metabolites were independently identified in subjects from the Framingham Offspring Study, and several metabolites, including glucose, lactate, hippurate and glycerol, had also been previously related to glucose metabolism (Pelkonen *et al.* 1967). However, this study demonstrated novel changes in the levels of bile acids

following the OGTT; the levels of three bile acids, glycocholic acid, glycochenodeoxycholic acid (GCDCA) and taurochenodeoxycholic acid, were increased within the first 30 min following glucose ingestion and remained stable thereafter. Another study (Zhao *et al.* 2009) examining healthy individuals revealed similar results concerning these bile acids and reported as much as a sixfold increase in these levels after 30 min (followed by a subsequent decrease). Both findings were confirmed by a third study (Matysik *et al.* 2011) that investigated bile acid signalling during the course of an OGTT in relation to 15 conjugated and unconjugated bile acids. In normal subjects, the levels of GCDCA, the bile acid with the highest plasma levels and chenodeoxycholic acid (CDCA) increased within the first 60 min of the OGTT. Furthermore, this study found that in response to oral glucose ingestion, the levels of all of the examined glycine- and taurine-conjugated bile acids were increased at 60 min and declined slightly over the following 60 min, whereas the levels of unconjugated bile acids, e.g. cholic acid, lithocholic acid and ursodeoxycholic acid, declined throughout the course of the OGTT. Bile acids are produced in the liver by the oxidation of cholesterol and are stored in the gall bladder. On food intake, bile acids are released into the duodenum and small intestine and facilitate the intestinal absorption of nutrients, particularly dietary fat, drugs and steroids. The majority of excreted bile acids are reabsorbed in the terminal ileum and return to the liver via the enterohepatic circulation, and very low levels of bile acids are found in the systemic circulation. Beside their major role in dietary lipid absorption, bile acids are metabolic factors that play regulatory roles in fat, glucose and energy metabolism (Houten *et al.* 2006, Lefebvre *et al.* 2009). The reported increase in bile acids in response to glucose ingestion is in concordance with a threefold increase in the levels of bile acids in human serum following a standard liquid meal (De Barros *et al.* 1982). Furthermore, oral glucose ingestion leads to increased levels of cholecystokinin, a hormone that stimulates the production of hepatic bile and gall bladder contractions (Liddle *et al.* 1985). The link between bile acids and glucose homeostasis was further confirmed by the demonstration of enhanced *Cyp7a1* mRNA expression following insulin injection or oral glucose administration in fasting mice (Li *et al.* 2012). Additionally, in primary human hepatocytes, insulin and glucose were shown to stimulate *CYP7A1* mRNA expression (Li *et al.* 2006, 2010b), which suggests the existence of glucose/insulin-regulated gene transcription in the liver. The *CYP7A1* gene encodes the enzyme cholesterol 7 α -hydroxylase, which is the rate-limiting enzyme in the cholesterol catabolic pathway and in the conversion of cholesterol to bile acids and therefore represents a major point of regulation during bile acid synthesis. The direct glucose/insulin-stimulated expression of *CYP7A1* leads to an increased bile acid pool size. Taken together, these findings indicate an important connection between bile acid metabolism and glucose homeostasis. Hence, it is not surprising that bile acid metabolism is altered

in patients with diabetes (Prawitt *et al.* 2011). In addition, a metabolomic study revealed higher plasma levels of GCDCA in subjects with impaired glucose tolerance compared with subjects with normal glucose tolerance (Zhao *et al.* 2010b). Another study detected differences in the composition of the bile acid pool between T2DM patients and controls (Brufau *et al.* 2010). Whereas the size of the total bile acid pool was not different, T2DM subjects demonstrated increased deoxycholic acid input rates and cholic acid synthesis rates but exhibited a lower proportion of CDCA. Furthermore, therapy with bile acid sequestrants leads to the expected reductions in both total cholesterol and low-density lipoprotein cholesterol as well as improvements in glycaemic control in T2DM patients (Garg & Grundy 1994, Suzuki *et al.* 2006, Zieve *et al.* 2007, Kondo & Kadowaki 2010). Compared with patients who received control treatments or placebos, T2DM patients given bile acid sequestrants demonstrated greater reductions in the levels of plasma glucose and HbA1c.

In regards to the OGTT studies, metabolites beside bile acids were also altered during the OGTT in healthy subjects. In fact, increases in lysophosphatidylcholine (Zhao *et al.* 2009) and decreases in amino acids (Table 1; Shaham *et al.* 2008, Spegel *et al.* 2010), acylcarnitines (Zhao *et al.* 2009) and fatty acids (Table 2; Zhao *et al.* 2009, Spegel *et al.* 2010) were reported. The study by Zhao *et al.* (2009) provided a more systematic overview of fatty acid plasma changes during an OGTT; although the levels of fatty acids declined during an OGTT, the levels of saturated (SFA) and monounsaturated fatty acids (MUFA) were more significantly decreased than those of polyunsaturated fatty acids (PUFA). Moreover, a substantial reduction in the SFA/MUFA ratio was observed, consisting of a shift from MUFA towards SFA. These findings indicate a change in fatty acid composition following an OGTT. A more detailed discussion on this topic is presented in the next chapter. Overall, metabolic studies have revealed alterations in metabolites related to pathways involved in the action of insulin, including lipolysis, ketogenesis, proteolysis and glucose metabolism. These results indicate a change from β -oxidation to glycolysis and fat storage in response to glucose ingestion.

Patient investigations

Comparisons between the levels of various metabolites in diabetic patients and healthy controls have confirmed many of the findings from animal studies as well as studies investigating metabolic changes during an OGTT. These findings from human patients can be summarised as follows:

- i) As expected, diabetic patients exhibited differences in glucose metabolism. Patients with T1DM under insulin deprivation (Lanza *et al.* 2010) or in T2DM (Li *et al.* 2009, Suhre *et al.* 2010) demonstrated elevated glucose or mannose levels compared with healthy controls. Furthermore, in both T1DM and T2DM

Table 1 List of altered amino acids in patients with diabetes or during an OGTT. Pathways were listed according to the Human Metabolome Database

Amino acids	Essential (E) non-essential (NE)	Alanine metabolism	Glutathione metabolism	Glycine and serine metabolism	Valine, leucine/ isoleucine degradation	Glucose-alanine cycle	Urea cycle	Bile acid biosynthesis	Methionine metabolism	Histidine metabolism	Further pathways	Reference patient studies	Reference OGTT studies
Alanine	NE	X		X		X	X				Selenoamino acid metabolism	Nicholson <i>et al.</i> (1984), Messana <i>et al.</i> (1998), Zuppi <i>et al.</i> (2002) and Oresic <i>et al.</i> (2008)	
Arginine	E	X		X		X	X					Shaham <i>et al.</i> (2008)	
Citrulline	NA					X					Arginine and proline metabolism, aspartate metabolism	Lanza <i>et al.</i> (2010)	
Glutamate	NE	X	X	X		X	X			X	Arginine and proline metabolism, amino sugar metabolism, cysteine metabolism, folate metabolism	Messana <i>et al.</i> (1998) and Lanza <i>et al.</i> (2010)	Spegel <i>et al.</i> (2010)
Glycine	NE	X	X	X				X	X		Carnitine synthesis, porphyrin metabolism	Shaham <i>et al.</i> (2008)	Suhre <i>et al.</i> (2010)
Histidine	E									X			Lanza <i>et al.</i> (2010) and Wang <i>et al.</i> (2011)
Homocitrulline	NS											Shaham <i>et al.</i> (2008) and Spegel <i>et al.</i> (2010)	Lanza <i>et al.</i> (2010) and Wang <i>et al.</i> (2011)
Isoleucine	E				X							Shaham <i>et al.</i> (2008)	Shaham <i>et al.</i> (2008)
Leucine	E				X							Shaham <i>et al.</i> (2008) and Spegel <i>et al.</i> (2010)	Shaham <i>et al.</i> (2008) and Spegel <i>et al.</i> (2010)
Lysine	E										Carnitine synthesis, biotin metabolism	Lanza <i>et al.</i> (2010), Suhre <i>et al.</i> (2010) and Wang <i>et al.</i> (2011)	Shaham <i>et al.</i> (2008), Zhao <i>et al.</i> (2009) and Spegel <i>et al.</i> (2010)
Methionine	E		X						X		Spermidine/spermine biosynthesis, betaine metabolism	Spegel <i>et al.</i> (2010)	Lanza <i>et al.</i> (2010)

(continued)

Table 1 Continued

Amino acids	Essential (E) non-essential (NE)	Alanine metabolism	Glutathione metabolism	Glycine and serine metabolism	Valine, leucine/ isoleucine degradation	Glucose-alanine cycle	Urea cycle	Bile acid biosynthesis	Methionine metabolism	Histidine metabolism	Further pathways	Reference patient studies	Reference OGTT studies
Ornithine	NE	X		X			X				Spermidine/spermine biosynthesis	Spegel <i>et al.</i> (2010)	Oresic <i>et al.</i> (2008), Zhao <i>et al.</i> (2010b) and Wang <i>et al.</i> (2011)
Phenylalanine	E										Phenylalanine and tyrosine metabolism	Zhao <i>et al.</i> (2009)	Lanza <i>et al.</i> (2010) and Wang <i>et al.</i> (2011)
Pyroglutamate	NA		X									Shaham <i>et al.</i> (2008)	Nicholson <i>et al.</i> (1984), Lanza <i>et al.</i> (2010) and Wang <i>et al.</i> (2011)
Threonine	E			X							Threonine and 2-oxobutanoate degradation Tryptophan metabolism	Shaham <i>et al.</i> (2008) and Spegel <i>et al.</i> (2010)	
Tryptophan	E											Nicholson <i>et al.</i> (1984), Messana <i>et al.</i> (1998), Zuppi <i>et al.</i> (2002) and Oresic <i>et al.</i> (2008)	
Tyrosine	NE										Phenylalanine and tyrosine metabolism, catecholamine biosynthesis	Shaham <i>et al.</i> (2008)	Shaham <i>et al.</i> (2008)
Valine	E				X						Propanoate metabolism	Lanza <i>et al.</i> (2010)	

NS, not specified, NA, not applicable.

Table 2 List of altered fatty acids in patients with diabetes or during an OGTT. Pathways were listed according to the Human Metabolome Database

Pathways		Reference patient studies	Reference OGTT studies
Saturated fatty acid (SFA)	C6:0	Caproic acid	Suhre <i>et al.</i> (2010)
	C7:0	Heptanoic acid	Suhre <i>et al.</i> (2010)
	C9:0	Pelargonic acid	Suhre <i>et al.</i> (2010)
C12:0	Lauric acid	β-Oxidation of very long-chain FA, FA biosynthesis, mitochondrial β-oxidation of short-chain saturated FA	Spegel <i>et al.</i> (2010)
C14:0	Myristic acid	NS	Yi <i>et al.</i> (2007, 2008)
C15:0	Pentadecanoic acid	NS	Yi <i>et al.</i> (2007, 2008)
C16:0	Palmitic acid	β-Oxidation of very long-chain FA, FA biosynthesis, mitochondrial β-oxidation of medium-chain saturated FA	Yi <i>et al.</i> (2006, 2007, 2008), Li <i>et al.</i> (2009) and Zhao <i>et al.</i> (2010b)
C18:0	Stearic acid	FA biosynthesis, FA elongation in mitochondria, FA metabolism, glycerolipid metabolism	Yi <i>et al.</i> (2006, 2007, 2008) and Zhao <i>et al.</i> (2010b)
C20:0	Arachidic acid	Mitochondrial β-oxidation of long-chain saturated FA	Yi <i>et al.</i> (2007, 2008)
C24:0	Lignoceric acid	NS	Yi <i>et al.</i> (2006, 2007, 2008)
Monounsaturated fatty acid (MUFA)	C14:1, C20:1	β-Oxidation of very long-chain FA	Yi <i>et al.</i> (2006, 2007, 2008) and Zhao <i>et al.</i> (2010b)
C16:1	C16:1n-7: palmitoleic acid	NS	Spegel <i>et al.</i> (2010)
C18:1	C16:1n-9: hexadecenoic acid	NS	Zhao <i>et al.</i> (2009) and Spegel <i>et al.</i> (2010)
	C18:1n-7: <i>cis</i> -vaccenic acid	NS	Yi <i>et al.</i> (2007, 2008), Li <i>et al.</i> (2009) and Zhao <i>et al.</i> (2010b)
	C18:1n-9: oleic acid	NS	Zhao <i>et al.</i> (2009) and Spegel <i>et al.</i> (2010)
Polyunsaturated fatty acid (PUFA)	C18:2	α-Linolenic acid/linoleic acid metabolism	Zhao <i>et al.</i> (2010b)
C18:3	C18:2n-6: linoleic acid	α-Linolenic acid/linoleic acid metabolism	Yi <i>et al.</i> (2007, 2008) and Zhao <i>et al.</i> (2010b)
C20:2	C18:3n-3: α-linolenic acid	NS	Zhao <i>et al.</i> (2009)
C20:3, C22:4	C18:3n-6: γ-linolenic acid	NS	Yi <i>et al.</i> (2007, 2008)
C20:4	–	NS	Yi <i>et al.</i> (2008) and Zhao <i>et al.</i> (2010b)
C20:5, C22:6	C20:4n-6: arachidonic acid	NS	Zhao <i>et al.</i> (2010b)
C22:5	–	NS	
Carnitine	–	Oxidation of branched-chain FA, carnitine synthesis, β-oxidation of very long-chain FA, mitochondrial β-oxidation of long/short-chain saturated FA	Mihalik <i>et al.</i> (2010)
	–		Mihalik <i>et al.</i> (2010)
Acylcarnitines	C3	Oxidation of branched-chain FA	Mihalik <i>et al.</i> (2010)
C4, C4OH	Propionylcarnitine	NS	Mihalik <i>et al.</i> (2010)
	(Hydroxy-)butyrylcarnitine		

(continued)

Table 2 Continued

	Pathways	Reference patient studies	Reference OGTT studies
C5, C5OH	(Hydroxy-)valeryl/carnitine	Mihalik <i>et al.</i> (2010)	Zhao <i>et al.</i> (2009)
C6, C6OH	(Hydroxy-)hexenyl/carnitine	Zhao <i>et al.</i> (2009)	Mihalik <i>et al.</i> (2010)
C8	Octanoyl/carnitine	Zhao <i>et al.</i> (2009)	Mihalik <i>et al.</i> (2010)
C10, C10:1	Decanoyl/carnitine	Mitochondrial β -oxidation of short-chain saturated FA	Mihalik <i>et al.</i> (2010)
C12	Dodecanoyl/carnitine		
C14:1, C14OH	(Hydroxy-)tetradecanoyl/carnitine		
C16, C16OH	(Hydroxy-)hexadecanoyl/carnitine		
C18, C18:1	Octadecanoyl/carnitine		
		Suhre <i>et al.</i> (2010)	
		Suhre <i>et al.</i> (2010)	
		Suhre <i>et al.</i> (2010)	
		Spegel <i>et al.</i> (2010)	Yi <i>et al.</i> (2007, 2008)

NS, not specified; FA, fatty acids.

patients (Messana *et al.* 1998, Zuppi *et al.* 2002) increased lactate levels were observed, at which these levels increased with the grade of glucosuria in T2DM patients (Messana *et al.* 1998).

ii) With respect to TCA cycle metabolites (and similar to *db/db* mice), patients with T1DM (Zuppi *et al.* 2002) and T2DM (Messana *et al.* 1998) had higher levels of citrate, and these levels were also associated with increasing glucosuria (Messana *et al.* 1998). An additional study investigated the metabolic profiles of children who later progressed to T1DM and found decreased succinate and citrate levels at the time of birth (Oresic *et al.* 2008).

iii) The above-mentioned results were further confirmed in regards to the levels of ketone bodies during diabetes. Higher levels of acetone, acetoacetate and β -hydroxybutyrate were observed in insulin-deprived T1DM (Lanza *et al.* 2010) and T2DM (Nicholson *et al.* 1984, Messana *et al.* 1998, Suhre *et al.* 2010) subjects, indicating ketoacidotic metabolic decompensation.

iv) Additionally, alterations in intestinal microflora-associated metabolites have been detected. Insulin-deprived T1DM and T2DM patients exhibited elevated levels of hippurate, dimethylamine or trimethylamine-*N*-oxide (Messana *et al.* 1998, Zuppi *et al.* 2002), although a separate study observed lower levels of hippurate and 3-hydroxyhippurate among pre-diabetic individuals with impaired glucose tolerance (Zhao *et al.* 2010b). The human gut microbiota has an important role in health, which has been comprehensively discussed by several authors (Nicholson *et al.* 2005, Fujimura *et al.* 2010, Prakash *et al.* 2011) and is outside the focus of the present review.

Fatty acid alterations in patients with diabetes have been extensively examined. In accordance with genetic rat models (Williams *et al.* 2006b, Salek *et al.* 2007), increased fatty acid levels were detected in T2DM patients (Yi *et al.* 2006, 2007, 2008, Li *et al.* 2009, Suhre *et al.* 2010) as well as in subjects with impaired glucose tolerance (Zhao *et al.* 2010b). Changes in as many as 18 fatty acids, including SFA, MUFA and PUFA, were found in one study. Furthermore, the metabolic profile of plasma acylcarnitines revealed higher fasting levels of long-chain saturated and monounsaturated acylcarnitines in obese and T2DM subjects compared with lean controls (Mihalik *et al.* 2010). Moreover, the levels of free carnitine were increased in both groups, although differences between groups were observed for short- and medium-chain acylcarnitine species as well as hydroxyacylcarnitines, where higher levels were observed in T2DM patients. Similar to an OGTT-induced reduction in acylcarnitines (Zhao *et al.* 2009), an insulin-stimulated euglycaemic clamp led to a decrease in all acylcarnitine species for all three of the investigated groups, although this reduction was blunted in patients with T2DM (Mihalik *et al.* 2010). All the investigated fatty acids and acylcarnitines are listed in Table 2. Many

studies have demonstrated that obese subjects often exhibit elevated fatty acid levels due to the enlarged volume of adipose tissue (Opie & Walfish 1963, Jensen *et al.* 1989, Newgard *et al.* 2009). Furthermore, higher levels of fatty acids are related to a greater risk for diabetes (Paolisso *et al.* 1995, Charles *et al.* 1997, Pankow *et al.* 2004), although the underlying mechanisms are not completely understood. However, elevated levels of free fatty acids induce insulin resistance in muscle and liver tissue by decreasing insulin-stimulated glucose uptake and glycogenesis (Griffin *et al.* 1999, Boden 2003, Wilding 2007). Moreover, the improvement in insulin sensitivity caused by a reduction in fatty acid levels supports these findings (Boden *et al.* 1998, Santomauro *et al.* 1999, Cusi *et al.* 2007). At present, there are several hypotheses as to how free fatty acids interfere with insulin signalling; these are related to oxidative stress or inflammatory lipid pathways and have been reviewed by Boden (2011).

An additional observation from patient studies is the change in amino acid levels in diabetic patients. A broad range of amino acids, including leucine, isoleucine, valine, phenylalanine, tyrosine, alanine, tryptophan and homocitulline, has been shown to be substantially increased in T1DM (Lanza *et al.* 2010) or T2DM patients (Messana *et al.* 1998, Suhre *et al.* 2010) as well as among subjects with obesity (Newgard *et al.* 2009) or impaired glucose tolerance (Zhao *et al.* 2010b). In addition, these findings have been confirmed by studies that revealed positive associations between amino acid levels and the homeostasis model assessment index (Newgard *et al.* 2009) and insulin resistance of obese subjects, according to Bergman's minimal model (Huffman *et al.* 2009). However, lower levels of glycine, glutamate and threonine have been observed in diabetic patients and obese subjects (Messana *et al.* 1998, Newgard *et al.* 2009, Lanza *et al.* 2010). A recent investigation (Wang *et al.* 2011) of the Framingham Offspring Study first examined the predictive ability of fasting plasma metabolite levels for incident T2DM and showed that the amino acids isoleucine, leucine, valine, tyrosine and phenylalanine were elevated 12 years before the onset of diabetes and were also linked to a higher diabetes risk. **In fact, the combination of increased fasting isoleucine, tyrosine and phenylalanine levels at baseline was related to a greater than fivefold higher risk of incident diabetes.** Moreover, these results were independently replicated in the Malmo Diet and Cancer Study. These studies have highlighted the impact of amino acids on the actions of insulin and consequently glucose metabolism. In the 1940s, Luetscher (1942) already reported higher amino acid levels among diabetic patients, and this finding was later confirmed by the demonstration of the positive correlation between amino acid levels and insulin (Felig *et al.* 1969). Furthermore, the i.v. administration of amino acids leads to the stimulation of insulin secretion. This insulinotropic effect differs depending on the specific amino acids in question (Floyd *et al.* 1966, 1968). However, the underlying mechanisms are complex and are related to the inhibition of glucose transport and gluconeogenesis (Patti *et al.* 1998, Krebs *et al.* 2002, Langenberg & Savage 2011).

Conclusion and challenges for the future

The application of metabolomics in diabetes studies has rapidly evolved during the last decade and provides researchers the opportunity to gain new insights into metabolic profiling and pathophysiological mechanisms. Thus, several metabolites were identified to be related to diabetes or insulin resistance and represent the basis for the identification of novel diabetes biomarkers. Some findings were newly discovered altered metabolites, e.g. bile acids, whereas other metabolic variations were already known, e.g. fatty acids or amino acids. However, the results often led to a revalue of knowledge. **Langenberg and colleague (Langenberg & Savage 2011) discussed the potential of an amino acid profile as a predictor of T2DM and highlighted the fact that the addition of amino acids to established risk factors only minimally improved the risk predication. This general problem is also apparent for genetic variants and other clinical novel biomarkers of diabetes whose power to add considerable improvement in risk assessment is limited (Lyssenko *et al.* 2008, Meigs *et al.* 2008, Salomaa *et al.* 2010).** Nevertheless, metabolomics increases the knowledge of disease progression and provides approaches for therapy.

Declaration of interest

The author declares that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review.

Funding

The author receives the Käthe-Kluth-Scholarship funded by the Ernst-Moritz-Arndt University Greifswald.

References

- American Association of Clinical Endocrinologists 2007 State of diabetes complications in America: a comprehensive report issued by the American Association of Clinical Endocrinologists. http://multivu.prnewswire.com/mnr/AACE/2007/docs/Diabetes_Complications_Report.pdf. Accessed May 23, 2012.
- Bell JD, Brown JC & Sadler PJ 1989 NMR studies of body fluids. *NMR in Biomedicine* **2** 246–256. (doi:10.1002/nbm.1940020513)
- Bictash M, Ebbls TM, Chan Q, Loo RL, Yap IK, Brown IJ, de Iorio M, Daviglus ML, Holmes E, Stamler J *et al.* 2010 Opening up the "Black Box": metabolic phenotyping and metabolome-wide association studies in epidemiology. *Journal of Clinical Epidemiology* **63** 970–979. (doi:10.1016/j.jclinepi.2009.10.001)
- Blümich B 2005 *Essential NMR: for Scientists and Engineers*. Berlin: Springer.
- Boden G 2003 Effects of free fatty acids (FFA) on glucose metabolism: significance for insulin resistance and type 2 diabetes. *Experimental and Clinical Endocrinology & Diabetes* **111** 121–124. (doi:10.1055/s-2003-39781)
- Boden G 2011 Obesity, insulin resistance and free fatty acids. *Current Opinion in Endocrinology, Diabetes, and Obesity* **18** 139–143. (doi:10.1097/MED.0b013e3283444b09)
- Boden G, Chen X & Iqbal N 1998 Acute lowering of plasma fatty acids lowers basal insulin secretion in diabetic and nondiabetic subjects. *Diabetes* **47** 1609–1612. (doi:10.2337/diabetes.47.10.1609)

- Brufau G, Stellaard F, Prado K, Bloks VW, Jonkers E, Boverhof R, Kuipers F & Murphy EJ 2010 Improved glycemic control with colessevelam treatment in patients with type 2 diabetes is not directly associated with changes in bile acid metabolism. *Hepatology* **52** 1455–1464. (doi:10.1002/hep.23831)
- Buscher JM, Czernik D, Ewald JC, Sauer U & Zamboni N 2009 Cross-platform comparison of methods for quantitative metabolomics of primary metabolism. *Analytical Chemistry* **81** 2135–2143. (doi:10.1021/ac8022857)
- Charles MA, Eschwege E, Thibault N, Claude JR, Warnet JM, Rosselin GE, Girard J & Balkau B 1997 The role of non-esterified fatty acids in the deterioration of glucose tolerance in Caucasian subjects: results of the Paris Prospective Study. *Diabetologia* **40** 1101–1106. (doi:10.1007/s001250050793)
- Chen D & Wang MW 2005 Development and application of rodent models for type 2 diabetes. *Diabetes, Obesity and Metabolism* **7** 307–317. (doi:10.1111/j.1463-1326.2004.00392.x)
- Connor SC, Hansen MK, Corner A, Smith RF & Ryan TE 2010 Integration of metabolomics and transcriptomics data to aid biomarker discovery in type 2 diabetes. *Molecular BioSystems* **6** 909–921. (doi:10.1039/b914182k)
- Cusi K, Kashyap S, Gastaldelli A, Bajaj M & Cersosimo E 2007 Effects on insulin secretion and insulin action of a 48-h reduction of plasma free fatty acids with acipimox in nondiabetic subjects genetically predisposed to type 2 diabetes. *American Journal of Physiology. Endocrinology and Metabolism* **292** E1775–E1781. (doi:10.1152/ajpendo.00624.2006)
- De Barros SG, Balistreri WF, Soloway RD, Weiss SK, Miller PC & Soper K 1982 Response of total and individual serum bile acids to endogenous and exogenous bile acid input to the enterohepatic circulation. *Gastroenterology* **82** 647–652.
- Delarue J & Magnan C 2007 Free fatty acids and insulin resistance. *Current Opinion in Clinical Nutrition and Metabolic Care* **10** 142–148. (doi:10.1097/MCO.0b013e328042ba90)
- De Palo E, Gatti R, Siculo N, Padovan D, Vettor R & Federspil G 1981 Plasma and urine free L-carnitine in human diabetes mellitus. *Acta Diabetologica Latina* **18** 91–95. (doi:10.1007/BF02056110)
- Duggan GE, Hittel DS, Hughey CC, Weljie A, Vogel HJ & Shearer J 2011 Differentiating short- and long-term effects of diet in the obese mouse using (1) H-nuclear magnetic resonance metabolomics. *Diabetes, Obesity and Metabolism* **13** 859–862. (doi:10.1111/j.1463-1326.2011.01410.x)
- Dunn WB, Bailey NJ & Johnson HE 2005 Measuring the metabolome: current analytical technologies. *Analyst* **130** 606–625. (doi:10.1039/b418288j)
- Fearnside JF, Dumas ME, Rothwell AR, Wilder SP, Cloarec O, Tøye A, Blancher C, Holmes E, Tatoud R, Barton RH *et al.* 2008 Phylometabonomic patterns of adaptation to high fat diet feeding in inbred mice. *PLoS ONE* **3** e1668. (doi:10.1371/journal.pone.0001668)
- Felig P, Marliss E & Cahill GF Jr 1969 Plasma amino acid levels and insulin secretion in obesity. *New England Journal of Medicine* **281** 811–816. (doi:10.1056/NEJM196910092811503)
- Floyd JC Jr, Fajans SS, Conn JW, Knopf RF & Rull J 1966 Stimulation of insulin secretion by amino acids. *Journal of Clinical Investigation* **45** 1487–1502. (doi:10.1172/JCI105456)
- Floyd JC Jr, Fajans SS, Conn JW, Thiffault C, Knopf RF & Guntzsch E 1968 Secretion of insulin induced by amino acids and glucose in diabetes mellitus. *Journal of Clinical Endocrinology and Metabolism* **28** 266–276. (doi:10.1210/jcem-28-2-266)
- Fraenkel M, Ketzinel-Gilad M, Ariav Y, Pappo O, Karaca M, Castel J, Berthault MF, Magnan C, Cerasi E, Kaiser N *et al.* 2008 mTOR inhibition by rapamycin prevents beta-cell adaptation to hyperglycemia and exacerbates the metabolic state in type 2 diabetes. *Diabetes* **57** 945–957. (doi:10.2337/db07-0922)
- Fujimura KE, Slusher NA, Cabana MD & Lynch SV 2010 Role of the gut microbiota in defining human health. *Expert Review of Anti-Infective Therapy* **8** 435–454. (doi:10.1586/eri.10.14)
- Garg A & Grundy SM 1994 Cholestyramine therapy for dyslipidemia in non-insulin-dependent diabetes mellitus. A short-term, double-blind, crossover trial. *Annals of Internal Medicine* **121** 416–422. (doi:10.1059/0003-4819-121-6-199409150-00004)
- Gipson GT, Tatsuoaka KS, Ball RJ, Sokhansanj BA, Hansen MK, Ryan TE, Hodson MP, Sweatman BC & Connor SC 2008 Multi-platform investigation of the metabolome in a leptin receptor defective murine model of type 2 diabetes. *Molecular BioSystems* **4** 1015–1023. (doi:10.1039/b807332e)
- Griffin ME, Marcucci MJ, Cline GW, Bell K, Barucci N, Lee D, Goodyear LJ, Kraegen EW, White MF & Shulman GI 1999 Free fatty acid-induced insulin resistance is associated with activation of protein kinase C theta and alterations in the insulin signaling cascade. *Diabetes* **48** 1270–1274. (doi:10.2337/diabetes.48.6.1270)
- Griffin JL, Atherton H, Shockcor J & Atzori L 2011 Metabolomics as a tool for cardiac research. *Nature Reviews. Cardiology* **8** 630–643. (doi:10.1038/nrcardio.2011.138)
- Herder C & Roden M 2011 Genetics of type 2 diabetes: pathophysiologic and clinical relevance. *European Journal of Clinical Investigation* **41** 679–692. (doi:10.1111/j.1365-2362.2010.02454.x)
- de Hoffmann & Stroobant 2007 *Mass Spectrometry: Principles and Applications*. New York: John Wiley & Sons.
- Hollywood K, Brison DR & Goodacre R 2006 Metabolomics: current technologies and future trends. *Proteomics* **6** 4716–4723. (doi:10.1002/pmic.200600106)
- Houten SM, Watanabe M & Auwerx J 2006 Endocrine functions of bile acids. *EMBO Journal* **25** 1419–1425. (doi:10.1038/sj.emboj.7601049)
- Huffman KM, Shah SH, Stevens RD, Bain JR, Muehlbauer M, Slentz CA, Tanner CJ, Kuchibhatla M, Houmard JA, Newgard CB *et al.* 2009 Relationships between circulating metabolic intermediates and insulin action in overweight to obese, inactive men and women. *Diabetes Care* **32** 1678–1683. (doi:10.2337/dc08-2075)
- Iles RA, Hind AJ & Chalmers RA 1985 Use of proton nuclear magnetic resonance spectroscopy in detection and study of organic acidurias. *Clinical Chemistry* **31** 1795–1801.
- Jensen MD, Haymond MW, Rizza RA, Cryer PE & Miles JM 1989 Influence of body fat distribution on free fatty acid metabolism in obesity. *Journal of Clinical Investigation* **83** 1168–1173. (doi:10.1172/JCI113997)
- Khamzina L, Veilleux A, Bergeron S & Marette A 2005 Increased activation of the mammalian target of rapamycin pathway in liver and skeletal muscle of obese rats: possible involvement in obesity-linked insulin resistance. *Endocrinology* **146** 1473–1481. (doi:10.1210/en.2004-0921)
- Kim SH, Yang SO, Kim HS, Kim Y, Park T & Choi HK 2009 ¹H-nuclear magnetic resonance spectroscopy-based metabolic assessment in a rat model of obesity induced by a high-fat diet. *Analytical and Bioanalytical Chemistry* **395** 1117–1124. (doi:10.1007/s00216-009-3054-8)
- Kim HJ, Kim JH, Noh S, Hur HJ, Sung MJ, Hwang JT, Park JH, Yang HJ, Kim MS, Kwon DY *et al.* 2011 Metabolomic analysis of livers and serum from high-fat diet induced obese mice. *Journal of Proteome Research* **10** 722–731. (doi:10.1021/pr100892r)
- Kondo K & Kadowaki T 2010 Colestilan monotherapy significantly improves glycaemic control and LDL cholesterol levels in patients with type 2 diabetes: a randomized double-blind placebo-controlled study. *Diabetes, Obesity and Metabolism* **12** 246–251. (doi:10.1111/j.1463-1326.2009.01159.x)
- Krebs M, Krssak M, Bernroider E, Anderwald C, Brehm A, Meyerspeer M, Nowotny P, Roth E, Waldhausl W & Roden M 2002 Mechanism of amino acid-induced skeletal muscle insulin resistance in humans. *Diabetes* **51** 599–605. (doi:10.2337/diabetes.51.3.599)
- Langenberg C & Savage DB 2011 An amino acid profile to predict diabetes? *Nature Medicine* **17** 418–420. (doi:10.1038/nm0411-418)
- Lanza IR, Zhang S, Ward LE, Karakelides H, Rafferty D & Nair KS 2010 Quantitative metabolomics by H-NMR and LC-MS/MS confirms altered metabolic pathways in diabetes. *PLoS ONE* **5** e10538. (doi:10.1371/journal.pone.0010538)
- Lefebvre P, Cariou B, Lien F, Kuipers F & Staels B 2009 Role of bile acids and bile acid receptors in metabolic regulation. *Physiological Reviews* **89** 147–191. (doi:10.1152/physrev.00010.2008)
- Lenz EM & Wilson ID 2007 Analytical strategies in metabolomics. *Journal of Proteome Research* **6** 443–458. (doi:10.1021/pr0605217)
- Li T, Kong X, Owsley E, Ellis E, Strom S & Chiang JY 2006 Insulin regulation of cholesterol 7alpha-hydroxylase expression in human hepatocytes: roles of

- forkhead box O1 and sterol regulatory element-binding protein 1c. *Journal of Biological Chemistry* **281** 28745–28754. (doi:10.1074/jbc.M605815200)
- Li X, Xu Z, Lu X, Yang X, Yin P, Kong H, Yu Y & Xu G 2009 Comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry for metabolomics: biomarker discovery for diabetes mellitus. *Analytica Chimica Acta* **633** 257–262. (doi:10.1016/j.aca.2008.11.058)
- Li LO, Hu YF, Wang L, Mitchell M, Berger A & Coleman RA 2010a Early hepatic insulin resistance in mice: a metabolomics analysis. *Molecular Endocrinology* **24** 657–666. (doi:10.1210/me.2009-0152)
- Li T, Chanda D, Zhang Y, Choi HS & Chiang JY 2010b Glucose stimulates cholesterol 7 α -hydroxylase gene transcription in human hepatocytes. *Journal of Lipid Research* **51** 832–842. (doi:10.1194/jlr.M002782)
- Li T, Francel JM, Boehme S, Ochoa A, Zhang Y, Klaassen CD, Erickson SK & Chiang JY 2012 Glucose and insulin induction of bile acid synthesis: mechanisms and implication in diabetes and obesity. *Journal of Biological Chemistry* **287** 1861–1873. (doi:10.1074/jbc.M111.305789)
- Liddle RA, Goldfine ID, Rosen MS, Taplitz RA & Williams JA 1985 Cholecystokinin bioactivity in human plasma. Molecular forms, responses to feeding, and relationship to gallbladder contraction. *Journal of Clinical Investigation* **75** 1144–1152. (doi:10.1172/JCI111809)
- Lin S, Yang Z, Liu H, Tang L & Cai Z 2011 Beyond glucose: metabolic shifts in responses to the effects of the oral glucose tolerance test and the high-fructose diet in rats. *Molecular BioSystems* **7** 1537–1548. (doi:10.1039/c0mb00246a)
- Lindon JC, Holmes E & Nicholson JK 2007 Metabolomics in pharmaceutical R&D. *FEBS Journal* **274** 1140–1151. (doi:10.1111/j.1742-4658.2007.05673.x)
- Lutscher JA 1942 The metabolism of amino acids in diabetes mellitus. *Journal of Clinical Investigation* **21** 275–279. (doi:10.1172/JCI101300)
- Lyssenko V, Jonsson A, Almgren P, Pulizzi N, Isomaa B, Tuomi T, Berglund G, Altschuler D, Nilsson P & Groop L 2008 Clinical risk factors, DNA variants, and the development of type 2 diabetes. *New England Journal of Medicine* **359** 2220–2232. (doi:10.1056/NEJMoa0801869)
- Major HJ, Williams R, Wilson AJ & Wilson ID 2006 A metabolomic analysis of plasma from Zucker rat strains using gas chromatography/mass spectrometry and pattern recognition. *Rapid Communications in Mass Spectrometry* **20** 3295–3302. (doi:10.1002/rcm.2732)
- Matysik S, Martin J, Bala M, Scherer M, Schaffler A & Schmitz G 2011 Bile acid signaling after an oral glucose tolerance test. *Chemistry and Physics of Lipids* **164** 525–529. (doi:10.1016/j.chemphyslip.2011.05.003)
- Meigs JB, Shrader P, Sullivan LM, McAteer JB, Fox CS, Dupuis J, Manning AK, Florez JC, Wilson PW, D'Agostino RB Sr *et al.* 2008 Genotype score in addition to common risk factors for prediction of type 2 diabetes. *New England Journal of Medicine* **359** 2208–2219. (doi:10.1056/NEJMoa0804742)
- Messana I, Forni F, Ferrari F, Rossi C, Giardina B & Zuppi C 1998 Proton nuclear magnetic resonance spectral profiles of urine in type II diabetic patients. *Clinical Chemistry* **44** 1529–1534.
- Mihalik SJ, Goodpaster BH, Kelley DE, Chace DH, Vockley J, Toledo FG & DeLany JP 2010 Increased levels of plasma acylcarnitines in obesity and type 2 diabetes and identification of a marker of glucolipotoxicity. *Obesity* **18** 1695–1700. (doi:10.1038/oby.2009.510)
- Mochida T, Tanaka T, Shiraki Y, Tajiri H, Matsumoto S, Shimbo K, Ando T, Nakamura K, Okamoto M & Endo F 2011 Time-dependent changes in the plasma amino acid concentration in diabetes mellitus. *Molecular Genetics and Metabolism* **103** 406–409. (doi:10.1016/j.ymgme.2011.05.002)
- Newgard CB, An J, Bain JR, Muehlbauer MJ, Stevens RD, Lien LF, Haqq AM, Shah SH, Arlotto M, Slentz CA *et al.* 2009 A branched-chain amino acid-related metabolic signature that differentiates obese and lean humans and contributes to insulin resistance. *Cell Metabolism* **9** 311–326. (doi:10.1016/j.cmet.2009.02.002)
- Nicholson JK, O'Flynn MP, Sadler PJ, Macleod AF, Juul SM & Sonksen PH 1984 Proton-nuclear-magnetic-resonance studies of serum, plasma and urine from fasting normal and diabetic subjects. *Biochemical Journal* **217** 365–375.
- Nicholson JK, Lindon JC & Holmes E 1999 'Metabolomics': understanding the metabolic responses of living systems to pathophysiological stimuli via multivariate statistical analysis of biological NMR spectroscopic data. *Xenobiotica* **29** 1181–1189. (doi:10.1080/004982599238047)
- Nicholson JK, Holmes E & Wilson ID 2005 Gut microorganisms, mammalian metabolism and personalized health care. *Nature Reviews. Microbiology* **3** 431–438. (doi:10.1038/nrmicro1152)
- Noble JA & Erlich HA 2012 Genetics of type 1 diabetes. *Cold Spring Harbor Perspectives in Medicine* **2** a007732.
- Oliver SG, Winson MK, Kell DB & Baganz F 1998 Systematic functional analysis of the yeast genome. *Trends in Biotechnology* **16** 373–378. (doi:10.1016/S0167-7799(98)01214-1)
- Opie LH & Walfish PG 1963 Plasma free fatty acid concentrations in obesity. *New England Journal of Medicine* **268** 757–760. (doi:10.1056/NEJM196304042681404)
- Oresic M, Simell S, Sysi-Aho M, Nanto-Salonen K, Seppanen-Laakso T, Parikka V, Katajamaa M, Hekkala A, Mattila I, Keskinen P *et al.* 2008 Dysregulation of lipid and amino acid metabolism precedes islet autoimmunity in children who later progress to type 1 diabetes. *Journal of Experimental Medicine* **205** 2975–2984. (doi:10.1084/jem.20081800)
- Pankow JS, Duncan BB, Schmidt MI, Ballantyne CM, Couper DJ, Hoogeveen RC & Golden SH 2004 Fasting plasma free fatty acids and risk of type 2 diabetes: the atherosclerosis risk in communities study. *Diabetes Care* **27** 77–82. (doi:10.2337/diacare.27.1.77)
- Paolisso G, Tataranni PA, Foley JE, Bogardus C, Howard BV & Ravussin E 1995 A high concentration of fasting plasma non-esterified fatty acids is a risk factor for the development of NIDDM. *Diabetologia* **38** 1213–1217. (doi:10.1007/BF00422371)
- Patterson AD, Bonzo JA, Li F, Krausz KW, Eichler GS, Aslam S, Tigno X, Weinstein JN, Hansen BC, Idle JR *et al.* 2011 Metabolomics reveals attenuation of the SLC6A20 kidney transporter in nonhuman primate and mouse models of type 2 diabetes mellitus. *Journal of Biological Chemistry* **286** 19511–19522. (doi:10.1074/jbc.M111.221739)
- Patti ME, Brambilla E, Luzi L, Landaker EJ & Kahn CR 1998 Bidirectional modulation of insulin action by amino acids. *Journal of Clinical Investigation* **101** 1519–1529. (doi:10.1172/JCI1326)
- Pelkonen R, Nikkila EA & Kekki M 1967 Metabolism of glycerol in diabetes mellitus. *Diabetologia* **3** 1–8. (doi:10.1007/BF01269904)
- Polychronakos C & Li Q 2011 Understanding type 1 diabetes through genetics: advances and prospects. *Nature Reviews. Genetics* **12** 781–792. (doi:10.1038/nrg3069)
- Prakash S, Tomaro-Duchesneau C, Saha S & Cantor A 2011 The gut microbiota and human health with an emphasis on the use of microencapsulated bacterial cells. *Journal of Biomedicine & Biotechnology* **2011** 981214.
- Prawitt J, Caron S & Staels B 2011 Bile acid metabolism and the pathogenesis of type 2 diabetes. *Current Diabetes Reports* **11** 160–166. (doi:10.1007/s11892-011-0187-x)
- Rhee EP & Gerszten RE 2012 Metabolomics and cardiovascular biomarker discovery. *Clinical Chemistry* **58** 139–147. (doi:10.1373/clinchem.2011.169573)
- Salek RM, Maguire ML, Bentley E, Rubtsov DV, Hough T, Cheeseman M, Nunez D, Sweatman BC, Haselden JN, Cox RD *et al.* 2007 A metabolomic comparison of urinary changes in type 2 diabetes in mouse, rat, and human. *Physiological Genomics* **29** 99–108. (doi:10.1152/physiolgenomics.00194.2006)
- Salomaa V, Havulinna A, Saarela O, Zeller T, Jousilahti P, Julia A, Muenzel T, Aromaa A, Evans A, Kuulasmaa K *et al.* 2010 Thirty-one novel biomarkers as predictors for clinically identified diabetes. *PLoS ONE* **5** e10100. (doi:10.1371/journal.pone.0010100)
- Santomauro AT, Boden G, Silva ME, Rocha DM, Santos RF, Ursich MJ, Strassmann PG & Wajchenberg BL 1999 Overnight lowering of free fatty acids with Acipimox improves insulin resistance and glucose tolerance in obese diabetic and nondiabetic subjects. *Diabetes* **48** 1836–1841. (doi:10.2337/diabetes.48.9.1836)

- Shaham O, Wei R, Wang TJ, Ricciardi C, Lewis GD, Vasan RS, Carr SA, Thadhani R, Gerszten RE & Mootha VK 2008 Metabolic profiling of the human response to a glucose challenge reveals distinct axes of insulin sensitivity. *Molecular Systems Biology* **4** 214. (doi:10.1038/msb.2008.50)
- Shearer J, Duggan G, Weljie A, Hittel DS, Wasserman DH & Vogel HJ 2008 Metabolomic profiling of dietary-induced insulin resistance in the high fat-fed C57BL/6J mouse. *Diabetes, Obesity and Metabolism* **10** 950–958. (doi:10.1111/j.1463-1326.2007.00837.x)
- Spegel P, Danielsson APH, Bacos K, Nagorny CLF, Moritz T, Mulder H & Filipsson K 2010 Metabolomic analysis of a human oral glucose tolerance test reveals fatty acids as reliable indicators of regulated metabolism. *Metabolomics* **6** 56–66. (doi:10.1007/s11306-009-0177-z)
- Suhre K, Meisinger C, Doring A, Altmair E, Belcredi P, Gieger C, Chang D, Milburn MV, Gall WE, Weinberger KM *et al.* 2010 Metabolic footprint of diabetes: a multiplatform metabolomics study in an epidemiological setting. *PLoS ONE* **5** e13953. (doi:10.1371/journal.pone.0013953)
- Suzuki T, Oba K, Futami S, Suzuki K, Ouchi M, Igari Y, Matsumura N, Watanabe K, Kigawa Y & Nakano H 2006 Blood glucose-lowering activity of colestimide in patients with type 2 diabetes and hypercholesterolemia: a case–control study comparing colestimide with acarbose. *Journal of Nihon Medical School* **73** 277–284. (doi:10.1272/jnms.73.277)
- Tweddale H, Notley-McRobb L & Ferenci T 1998 Effect of slow growth on metabolism of *Escherichia coli*, as revealed by global metabolite pool ("metabolome") analysis. *Journal of Bacteriology* **180** 5109–5116.
- Um SH, Frigerio F, Watanabe M, Picard F, Joaquin M, Sticker M, Fumagalli S, Allegrini PR, Kozma SC, Auwerx J *et al.* 2004 Absence of S6K1 protects against age- and diet-induced obesity while enhancing insulin sensitivity. *Nature* **431** 200–205. (doi:10.1038/nature02866)
- Wang TJ, Larson MG, Vasan RS, Cheng S, Rhee EP, McCabe E, Lewis GD, Fox CS, Jacques PF, Fernandez C *et al.* 2011 Metabolite profiles and the risk of developing diabetes. *Nature Medicine* **17** 448–453. (doi:10.1038/nm.2307)
- Wild S, Roglic G, Green A, Sicree R & King H 2004 Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* **27** 1047–1053. (doi:10.2337/diacare.27.5.1047)
- Wilding JP 2007 The importance of free fatty acids in the development of type 2 diabetes. *Diabetic Medicine* **24** 934–945. (doi:10.1111/j.1464-5491.2007.02186.x)
- Williams RE, Lenz EM, Evans JA, Wilson ID, Granger JH, Plumb RS & Stumpf CL 2005a A combined (1)H NMR and HPLC-MS-based metabolomic study of urine from obese (fa/fa) Zucker and normal Wistar-derived rats. *Journal of Pharmaceutical and Biomedical Analysis* **38** 465–471. (doi:10.1016/j.jpba.2005.01.013)
- Williams RE, Lenz EM, Lowden JS, Rantalainen M & Wilson ID 2005b The metabolomics of aging and development in the rat: an investigation into the effect of age on the profile of endogenous metabolites in the urine of male rats using ¹H NMR and HPLC-TOF MS. *Molecular BioSystems* **1** 166–175. (doi:10.1039/b500852b)
- Williams R, Lenz EM, Wilson AJ, Granger J, Wilson ID, Major H, Stumpf C & Plumb R 2006a A multi-analytical platform approach to the metabolomic analysis of plasma from normal and Zucker (fa/fa) obese rats. *Molecular BioSystems* **2** 174–183. (doi:10.1039/b516356k)
- Williams RE, Lenz EM, Rantalainen M & Wilson ID 2006b The comparative metabolomics of age-related changes in the urinary composition of male Wistar-derived and Zucker (fa/fa) obese rats. *Molecular BioSystems* **2** 193–202. (doi:10.1039/b517195d)
- World Health Organization 2006 Definition and diagnosis of diabetes mellitus and intermediate hyperglycemia. Report of a WHO consultation. Geneva.
- Xu J, Zhang J, Cai S, Dong J, Yang JY & Chen Z 2009 Metabonomics studies of intact hepatic and renal cortical tissues from diabetic db/db mice using high-resolution magic-angle spinning ¹H NMR spectroscopy. *Analytical and Bioanalytical Chemistry* **393** 1657–1668. (doi:10.1007/s00216-009-2623-1)
- Yi LZ, He J, Liang YZ, Yuan DL & Chau FT 2006 Plasma fatty acid metabolic profiling and biomarkers of type 2 diabetes mellitus based on GC/MS and PLS-LDA. *FEBS Letters* **580** 6837–6845. (doi:10.1016/j.febslet.2006.11.043)
- Yi L, He J, Liang Y, Yuan D, Gao H & Zhou H 2007 Simultaneously quantitative measurement of comprehensive profiles of esterified and non-esterified fatty acid in plasma of type 2 diabetic patients. *Chemistry and Physics of Lipids* **150** 204–216. (doi:10.1016/j.chemphyslip.2007.08.002)
- Yi L, Yuan D, Che Z, Liang Y, Zhou Z, Gao H & Wang Y 2008 Plasma fatty acid metabolic profile coupled with uncorrelated linear discriminant analysis to diagnose and biomarker screening of type 2 diabetes and type 2 diabetic coronary heart diseases. *Metabolomics* **4** 30–38. (doi:10.1007/s11306-007-0098-7)
- Zhang S, Nagana Gowda GA, Asiago V, Shanaiah N, Barbas C & Raftery D 2008 Correlative and quantitative ¹H NMR-based metabolomics reveals specific metabolic pathway disturbances in diabetic rats. *Analytical Biochemistry* **383** 76–84. (doi:10.1016/j.ab.2008.07.041)
- Zhao X, Peter A, Fritsche J, Elcnrova M, Fritsche A, Haring HU, Schleicher ED, Xu G & Lehmann R 2009 Changes of the plasma metabolome during an oral glucose tolerance test: is there more than glucose to look at? *American Journal of Physiology. Endocrinology and Metabolism* **296** E384–E393. (doi:10.1152/ajpendo.90748.2008)
- Zhao LC, Zhang XD, Liao SX, Gao HC, Wang HY & Lin DH 2010a A metabolomic comparison of urinary changes in Zucker and GK rats. *Journal of Biomedicine & Biotechnology* **2010** 431894.
- Zhao X, Fritsche J, Wang J, Chen J, Rittig K, Schmitt-Kopplin P, Fritsche A, Haring HU, Schleicher ED, Xu G *et al.* 2010b Metabolomic fingerprints of fasting plasma and spot urine reveal human pre-diabetic metabolic traits. *Metabolomics* **6** 362–374. (doi:10.1007/s11306-010-0203-1)
- Zieve FJ, Kalin MF, Schwartz SL, Jones MR & Bailey WL 2007 Results of the glucose-lowering effect of WelChol study (GLOWS): a randomized, double-blind, placebo-controlled pilot study evaluating the effect of colesevelam hydrochloride on glycemic control in subjects with type 2 diabetes. *Clinical Therapeutics* **29** 74–83. (doi:10.1016/j.clinthera.2007.01.003)
- Zuppi C, Messina I, Tapanainen P, Knip M, Vincenzoni F, Giardina B & Nuutinen M 2002 Proton nuclear magnetic resonance spectral profiles of urine from children and adolescents with type 1 diabetes. *Clinical Chemistry* **48** 660–662.

Received in final form 1 June 2012

Accepted 20 June 2012

Made available online as an Accepted Preprint
20 June 2012