

Contents

1	Recollections	1
1.1	The L -group	1
1.2	Pure and rigid inner forms	2
1.3	Weyl denominators	2
1.4	Double covers of tori and L -embeddings	3
1.5	Essentially square-integrable representations	5
1.6	Endoscopic groups and double covers	6
2	Whittaker data and generic representations	9
3	Construction of L-packet and internal structure	10
3.1	Factorization of a parameter	11
3.2	Construction of the L -packet	12
3.3	Internal structure of the compound packet	12
3.4	The case of a cover of G	13
3.5	Dependence on the choice of Whittaker datum	13
4	Endoscopic character identities	13
4.1	Statement of the main theorem	13
4.2	Reduction to the elliptic set	14
4.3	The left hand side	14
4.4	The right hand side: covers	15
4.5	The right hand side: classical set-up	17
	References	19

1 RECOLLECTIONS

1.1 The L -group

We review the L -group of a connected reductive group following [Vog93, §2].

Let F be a field. Assume first that F is separably closed. Let G be a connected reductive F -group. Given a Borel pair (T, B) of G one has the based root datum $\text{brd}(T, B, G) = (X^*(T), \Delta, X_*(T), \Delta^\vee)$, where $\Delta \subset X^*(T)$ is the set of B -simple roots for the adjoint action of T on $\text{Lie}(G)$, and $\Delta^\vee \subset X_*(T)$ are the corresponding coroots. For a second Borel pair (T', B') , there is a unique element of $T'(F) \backslash G(F) / T(F)$ that conjugates (T, B) to (T', B') . This element provides an isomorphism $\text{brd}(T, B, G) \rightarrow \text{brd}(T', B', G)$. This procedure leads to a system of based root data and isomorphisms, indexed by the set of Borel pairs of G . The limit of that system is the based root datum $\text{brd}(G)$ of G .

One can formalize the notion of a based root datum: we refer the reader to [Spr09, §7.4] for the formal notion of a root datum, to which one has to add a set of simple roots to obtain the formal notion of a based root datum. Based root data can be placed into a category, in which all morphisms are isomorphisms,

for the evident notion of isomorphism of based root data. The classification of connected reductive F -groups [Spr09, Theorem 9.6.2, Theorem 10.1.1] can be stated as saying that $G \mapsto \text{brd}(G)$ is a full essentially surjective functor from the category of connected reductive F -groups and isomorphisms to the category of based root data and isomorphisms. Moreover, two morphisms lie in the same fiber of this functor if and only if they differ by an inner automorphism.

Consider now a general field F , let F^s a separable closure, $\Gamma = \text{Gal}(F^s/F)$ the Galois group. Given a connected reductive F -group G , there is a natural action of Γ on the set of Borel pairs of G_{F^s} , and this leads to a natural action of Γ on $\text{brd}(G_{F^s})$. We denote by $\text{brd}(G)$ the based root datum $\text{brd}(G_{F^s})$ equipped with this Γ -action. Given two connected reductive F -groups G_1, G_2 , an isomorphism $\xi : G_{1,F^s} \rightarrow G_{2,F^s}$ is called an *inner twist*, if $\xi^{-1} \circ \sigma \circ \xi \circ \sigma^{-1}$ is an inner automorphism of G_{1,F^s} for all $\sigma \in \Gamma$. The two groups G_1, G_2 are then called inner forms of each other. The functor $G \mapsto \text{brd}(G)$ from the category of connected reductive F -groups to the category of based root data over F and isomorphisms is again essentially surjective. It maps inner twists to isomorphisms, and two inner twists map to the same isomorphism if they differ by an inner automorphism. The fiber over a given based root datum over F consists of all reductive groups that are inner forms of each other.

Given a based root datum $(X, \Delta, Y, \Delta^\vee)$ over F , its dual $(Y, \Delta^\vee, X, \Delta)$ is also a based root datum over F . If G is a connected reductive F -group with based root datum $(X, \Delta, Y, \Delta^\vee)$, its dual \widehat{G} is the unique split connected reductive group defined over a chosen base field (we will work with \mathbb{C}) with based root datum $(Y, \Delta^\vee, X, \Delta)$. Thus, given a Borel pair $(\widehat{T}, \widehat{B})$ of \widehat{G} and a Borel pair (T, B) of G_{F^s} , one is given an identification $X_*(\widehat{T}) = X^*(T)$ that identifies the Weyl chambers associated to \widehat{B} and B .

To form the L -group, one chooses a pinning $(\widehat{T}, \widehat{B}, \{Y_\alpha\})$ of \widehat{G} . The group of automorphisms of \widehat{G} that preserve this pinning is in natural isomorphism with the group of automorphisms of $\text{brd}(\widehat{G})$, hence with that of $\text{brd}(G)$. The Γ -action on $\text{brd}(G)$ then lifts to an action on \widehat{G} by algebraic automorphisms, and ${}^L G = \widehat{G} \rtimes \Gamma$.

1.2 Pure and rigid inner forms

Let G_0 be a quasi-split reductive \mathbb{R} -group. Following Vogan [Vog93], a pure inner twist of G_0 is a triple (G, ξ, z) , where G is a connected reductive \mathbb{R} -group, $\xi : G_{0,\mathbb{C}} \rightarrow G_{\mathbb{C}}$ is an isomorphism and $z \in Z^1(\Gamma, G_0)$, subject to $\xi^{-1} \sigma(\xi) = \text{Ad}(\bar{z}_\sigma)$. An isomorphism of pure inner twists $(G_1, \xi_1, z_1) \rightarrow (G_2, \xi_2, z_2)$ is a pair (f, g) consisting of an isomorphism $f : G_1 \rightarrow G_2$ of \mathbb{R} -groups and $g \in G_0(\mathbb{C})$ such that

finish

1.3 Weyl denominators

Let G be a connected reductive \mathbb{R} -group and $T \subset G$ a maximal \mathbb{R} -torus. One can consider the function $T(\mathbb{R}) \rightarrow \mathbb{R}$ defined as

$$D_G(t) = \prod_{\alpha \in R(T, G)} (1 - \alpha(t)).$$

In this paper we will normalize orbital integrals and characters by multiplying them by $|D_G(t)|^{1/2}$. Thus, for $t \in T(\mathbb{R})$ strongly regular and $f \in \mathcal{C}_c^\infty(G(\mathbb{R}))$ we set

$$J(t, f) = |D_G(t)| \int_{G(\mathbb{R})/T(\mathbb{R})} f(gt g^{-1}) dg,$$

while for π admissible representation of $G(\mathbb{R})$ we have

$$J(t, \pi) = |D_G(t)| \Theta_\pi(t),$$

where Θ_π is the character function of π . This has the advantage that the resulting functions remain bounded as t approaches singular elements in $T(\mathbb{R})$.

A key role in this paper will be played by a function D_B which has the property that $|D_B| = |D_G|^{1/2}$. To see this function, let us interpret D_G as an element of the group ring $\mathbb{Z}[Q]$, where $Q \subset X^*(T) \otimes \mathbb{Q}$ is the root lattice, we write the group operation on Q multiplicatively. Given a Borel \mathbb{C} -subgroup $B \subset G$ containing T we write $\alpha > 0$ when α is a B -positive root, and define

$$D'_B = \prod_{\alpha > 0} (1 - \alpha^{-1}) \in \mathbb{Q}[Q], \quad D_B = \prod_{\alpha > 0} (\alpha^{1/2} - \alpha^{-1/2}) \in \mathbb{Q}[Q]. \quad (1.1)$$

In $\mathbb{Q}[Q]$ we have the identity

$$D_B = \rho \cdot D'_B, \quad (1.2)$$

where $\rho = \prod_{\alpha > 0} \alpha^{1/2} \in Q$. This implies

$$D_G = D'_B \cdot D'_B = D_B \cdot D_{\bar{B}},$$

where \bar{B} is the Borel subgroup opposite to B . Moreover, for $w \in \Omega(T, G)$ we have

$$w D_B = D_{w^{-1} B w} = \text{sgn}(w) D_B. \quad (1.3)$$

In particular, $|D_B|$ is independent of the choice of B and hence $|D_G|^{1/2} = |D_B|$, provided we can interpret D_B as a function on $T(\mathbb{R})$.

It is clear that D'_B is a function on $T(\mathbb{R})$. If we want to interpret D_B as a function of $T(\mathbb{R})$, the occurrence of $\alpha^{1/2}$ in the formula causes a problem. From (1.2) we see that D_B will be a function of $T(\mathbb{R})$ if and only if ρ is, which is equivalent to the element ρ lying in $X^*(T)$. This is always the case when G is semi-simple and simply connected, but can fail in general. To remedy this situation, one can introduce a double cover of $T(\mathbb{R})$, which will be discussed in the next section.

1.4 Double covers of tori and L -embeddings

Let G be a connected reductive \mathbb{R} -group and let $T \subset G$ be a maximal torus. An obstruction to the element D_B defining a function on $T(\mathbb{R})$ is the fact that $\rho \in \frac{1}{2} X^*(T)$ may not lie in $X^*(T)$. To remedy this, Adams–Vogan introduce in [AV92], [AV16] the ρ -double cover $T(\mathbb{R})_\rho$ as the pull-back of the diagram

$$T(\mathbb{R}) \xrightarrow{\rho^2} \mathbb{C}^\times \xleftarrow{(-)^2} \mathbb{C}^\times,$$

which comes equipped with a natural character $\rho : T(\mathbb{R})_\rho \rightarrow \mathbb{C}^\times$, namely the projection onto the right factor \mathbb{C}^\times . By construction we have an exact sequence

$$1 \rightarrow \{\pm 1\} \rightarrow T(\mathbb{R})_\rho \rightarrow T(\mathbb{R}) \rightarrow 1$$

and ρ is a genuine character, i.e. $\rho(-x) = -\rho(x)$ for $x \in T(\mathbb{R})_\rho$, where $-x$ denote the product of x and the element -1 .

While the double cover $T(\mathbb{R})_\rho$ appears to depend on ρ , this is actually not so. Indeed, for any other Borel \mathbb{C} -subgroup B' we have $\rho'/\rho \in X^*(T)$, which allows us to define the genuine character $\rho' : T(\mathbb{R})_\rho \rightarrow \mathbb{C}^\times$ as $\rho \cdot (\rho'/\rho)$. Combining this character with the natural projection $T(\mathbb{R})_\rho \rightarrow T(\mathbb{R})$ gives a map from $T(\mathbb{R})_\rho$ to the diagram defining $T(\mathbb{R})_{\rho'}$, hence an isomorphism $T(\mathbb{R})_\rho \rightarrow T(\mathbb{R})_{\rho'}$.

To emphasize the independence of $T(\mathbb{R})_\rho$ on ρ , and emphasize the dependence on the ambient group G , we will write $T(\mathbb{R})_G$ for this cover. For each Borel \mathbb{C} -subgroup B we have the genuine character $\rho_B : T(\mathbb{R})_G \rightarrow \mathbb{C}^\times$.

In this paper we will be particularly interested in the case when T is elliptic. Then there is a different way to obtain the ρ -cover that generalizes to all local fields, as discussed in [Kal19]. One first defines the “big cover” of $T(\mathbb{R})$ as follows. Each root provides a homomorphism $\alpha : T(\mathbb{R}) \rightarrow \mathbb{S}^1$. Combining these homomorphisms for a pair $A = \{\alpha, -\alpha\}$ provides a homomorphism $A : T(\mathbb{R}) \rightarrow \mathbb{S}^1_-$, where $\mathbb{S}^1_- \subset \mathbb{S}^1 \times \mathbb{S}^1$ is the kernel of the product map $\mathbb{S}^1 \times \mathbb{S}^1 \rightarrow \mathbb{S}^1$. Define the “big cover” as the pull-back of

$$T(\mathbb{R}) \xrightarrow{(\alpha)} \prod \mathbb{S}^1_- \xrightarrow{z/\bar{z}} \prod (\mathbb{C}^\times/\mathbb{R}_{>0})_- \quad (1.4)$$

where the products run over the set of pairs $A = \{\alpha, -\alpha\}$ consisting of a root and its negative, and $(\mathbb{C}^\times/\mathbb{R}_{>0})_-$ denotes analogously the anti-diagonal in $(\mathbb{C}^\times/\mathbb{R}_{>0}) \times (\mathbb{C}^\times/\mathbb{R}_{>0})$. The result is an extension

$$1 \rightarrow \prod \{\pm 1\} \rightarrow T(\mathbb{R})_{GG} \rightarrow T(\mathbb{R}) \rightarrow 1.$$

Under the isomorphism $\mathbb{S}^1 \rightarrow \mathbb{C}^\times/\mathbb{R}_{>0}$ the map z/\bar{z} becomes the squaring map, and we see that $T(\mathbb{R})_{GG}$ is equipped with a character $\alpha^{1/2} : T(\mathbb{R})_{GG} \rightarrow \mathbb{S}^1$ for each root α , and that $\beta^{1/2} = (\alpha^{1/2})^{-1}$ whenever $\beta = \alpha^{-1}$. There is an obvious surjective homomorphism $T(\mathbb{R})_{GG} \rightarrow T(\mathbb{R})_G$ whose kernel is the kernel of the multiplication map $\prod \{\pm 1\} \rightarrow \{\pm 1\}$. The function $\alpha^{1/2} - \alpha^{-1/2}$ is well-defined on the big cover, while for any choice Borel \mathbb{C} -subgroup V the function

$$D_B := \prod_{\alpha > 0} (\alpha^{1/2} - \alpha^{-1/2})$$

descends to the double cover $T(\mathbb{R})_G$.

The action of the Weyl group $\Omega_G(T)(\mathbb{R})$ lifts naturally to an action on $T(\mathbb{R})_G$, even on $T(\mathbb{R})_{GG}$, because $\Omega_G(T)(\mathbb{R})$ acts naturally on each term in (1.4). The identity (1.3) holds for this function.

What makes the double cover $T(\mathbb{R})_G$ very useful in the setting of the Langlands program is the fact that there is an associated L -group ${}^L T_G$, as well as a canonical \widehat{G} -conjugacy class of L -embeddings ${}^L T_G \rightarrow {}^L G$, cf. [Kal19, §4.1]. The property of the L -group ${}^L T_G$ is that the set of \widehat{T} -conjugacy classes of L -homomorphisms $W_{\mathbb{R}} \rightarrow {}^L T_G$ is in natural bijection with the set of genuine characters of $T(\mathbb{R})_G$. Therefore, any L -parameter for G that factors through the image of the embedding of ${}^L T_G$ provides in a canonical way an $\Omega_G(T)(\mathbb{R})$ -orbit of genuine characters of $T(\mathbb{R})_G$.

In contrast to ${}^L T_G$, there is generally no canonical L -embedding ${}^L T \rightarrow {}^L G$. In fact, if the Galois form of ${}^L T$ is used, there is generally no L -embedding

${}^L T \rightarrow {}^L G$ at all, let alone a canonical one. If the Weil form of ${}^L T$ is used, then there always do exist L -embeddings ${}^L T \rightarrow {}^L G$, but there is generally no canonical choice. If one chooses a genuine character of $T(\mathbb{R})_G$, then the pointwise product of its L -parameter $W_{\mathbb{R}} \rightarrow {}^L T_G$ with the natural inclusion $\widehat{T} \rightarrow {}^L T_G$ does lead to an L -isomorphism ${}^L T \rightarrow {}^L T_G$ between the Weil forms of the L -groups for $T(\mathbb{R})$ and $T(\mathbb{R})_G$. Composing this isomorphism with the canonical L -embedding ${}^L T_G \rightarrow {}^L G$ provides an L -embedding ${}^L T \rightarrow {}^L G$, and every L -embedding arises from this construction. A convenient choice for a genuine character on $T(\mathbb{R})_G$ is the character ρ associated to some Borel \mathbb{C} -subgroup of G containing T .

It is worth pointing out that all L -embeddings ${}^L T \rightarrow {}^L G$, as well as all L -embeddings ${}^L T_G \rightarrow {}^L G$, that extend a fixed embedding $\widehat{j} : \widehat{T} \rightarrow \widehat{G}$, have the same image, namely

$$\{x \in N_{LG}(\widehat{T}) \mid x \cdot \widehat{j}(t) \cdot x^{-1} = \widehat{j}(\sigma_x(t)) \ \forall t \in \widehat{T}\}, \quad (1.5)$$

where $\sigma_x \in \Gamma$ is the image of x under the natural projection ${}^L G \rightarrow \Gamma$.

1.5 Essentially square-integrable representations

Let G be a connected reductive \mathbb{R} -group. An essentially square integrable representation of $G(\mathbb{R})$ is one which, after possibly tensoring with a continuous character $G(\mathbb{R}) \rightarrow \mathbb{C}^\times$, has a unitary central character, and such that every matrix coefficient is square-integrable on $G(\mathbb{R})/Z_G(\mathbb{R})$ (since the central character is unitary, the absolute value of a matrix coefficient is trivial on $Z_G(\mathbb{R})$).

Harish-Chandra has shown that the set of isomorphism classes of essentially square-integrable representations is in bijection with the set of $G(\mathbb{R})$ -conjugacy classes of pairs (S, τ) , where $S \subset G$ is an elliptic maximal torus, and τ is a genuine character of the double cover $S(\mathbb{R})_G$ whose differential is regular. This bijection is characterized by the fact that the character function of the representation corresponding to (S, τ) , evaluated at a regular element $\delta \in S(\mathbb{R})$, is given by

$$(-1)^{q(G)} \sum_{w \in N(S, G)(\mathbb{R})/S(\mathbb{R})} \frac{\tau}{d_\tau}(w\delta) = (-1)^{q(G)} \sum_{w \in N(S, G)(\mathbb{R})/S(\mathbb{R})} \frac{\tau'}{d'_\tau}(w\delta). \quad (1.6)$$

We explain the notation. Pull back τ to a character of $S_{\text{sc}}(\mathbb{R})_G$, where S_{sc} is the preimage of S in the universal cover G_{sc} of the derived subgroup of G . The cover $S_{\text{sc}}(\mathbb{R})_G$ splits canonically, because ρ is divisible by 2 in $X^*(S_{\text{sc}})$. Therefore τ provides a character τ_{sc} of $S_{\text{sc}}(\mathbb{R})$. This being a compact torus, τ_{sc} is an algebraic character, i.e. an element of $X^*(S_{\text{sc}})$, and coincides with its differential, which is still regular. Thus τ specifies a choice of positive roots, i.e. a Borel \mathbb{C} -subgroup B containing S . Write D_τ in place of D_B for the Weyl denominator (1.1). Since our convention (cf. §1.3) is to normalize orbital integrals and characters by the absolute value of this denominator, we will only need $d_\tau = \arg D_\tau$. Both τ and d_τ are genuine functions of $S(\mathbb{R})_G$, so their quotient $\Theta := \tau/d_\tau$ descends to $S(\mathbb{R})$.

In the second sum we have set $\tau' = \tau \cdot \rho_B^{-1}$, and $d'_\tau = d_\tau \cdot \rho_B^{-1}$, cf. (1.2). In this way, both numerator and denominator are functions of $S(\mathbb{R})$. Note that ρ_B takes values in \mathbb{S}^1 because S is elliptic.

1.6 Endoscopic groups and double covers

The notion of endoscopic data is introduced in [LS87, §1.2], and is a variation of the notion of endoscopic pairs or endoscopic triples discussed in [Kot84] and [Kot86].

It can be described equivalently as follows. An endoscopic datum for G is a tuple $(H, s, \mathcal{H}, \eta)$ consisting of

- (1) a quasi-split connected reductive group H ,
- (2) an extension $1 \rightarrow \widehat{H} \rightarrow \mathcal{H} \rightarrow \Gamma \rightarrow 1$ of topological groups,
- (3) a semi-simple element $s \in Z(\widehat{H})$, and
- (4) an L -embedding $\mathcal{H} \rightarrow {}^L G$.

It is required that

- (a) the extension \mathcal{H} admits a splitting by a continuous group homomorphism $\Gamma \rightarrow \mathcal{H}$,
- (b) the homomorphism $\Gamma \rightarrow \text{Out}(\widehat{H})$ provided by \mathcal{H} coincides with the one provided by the extension ${}^L H$,
- (c) η identifies \widehat{H} with the identity component of the centralizer of $\eta(s)$ in \widehat{G} ,
- (d) there exists $z \in Z(\widehat{G})$ such that $s\eta^{-1}(z) \in Z(\widehat{H})^\Gamma$.

The map η produces a Γ -equivariant embedding $Z(\widehat{G}) \rightarrow Z(\widehat{H})$, that we will use without explicit notation.

An isomorphism $(H_1, s_1, \mathcal{H}_1, \eta_1) \rightarrow (H_2, s_2, \mathcal{H}_2, \eta_2)$ is an element $g \in \widehat{G}$ that satisfies the following properties. First, $\text{Ad}(g)\eta_1(\mathcal{H}_1) = \eta_2(\mathcal{H}_2)$. In particular, $\eta_2^{-1} \circ \text{Ad}(g) \circ \eta_1$ is an L -isomorphism $\mathcal{H}_1 \rightarrow \mathcal{H}_2$, and restricts to a Γ -equivariant isomorphism $Z(\widehat{H}_1) \rightarrow Z(\widehat{H}_2)$. The second condition is that the resulting isomorphism $\pi_0(Z(\widehat{H}_1)/Z(\widehat{G})) \rightarrow \pi_0(Z(\widehat{H}_2)/Z(\widehat{G}))$ maps the coset of s_1 to the coset of s_2 .

In this paper we are working with pure (resp. rigid) inner twists, and this necessitates a slight refinement of the notion of endoscopic datum. A *pure refined* endoscopic datum is one in which it is required $s \in Z(\widehat{H})^\Gamma$ in point (3), and this eliminates the need for condition (d). An isomorphism of such data is required to map the coset of s_1 to the coset of s_2 under $\pi_0(Z(\widehat{H}_1)^\Gamma) \rightarrow \pi_0(Z(\widehat{H}_2)^\Gamma)$, without dividing by $Z(\widehat{G})$. A *rigid refined* endoscopic datum replaces $s \in Z(\widehat{H})^\Gamma$ by $\dot{s} \in Z(\widehat{H})^+$. An isomorphism of such data is required to map the coset of \dot{s}_1 to the coset of \dot{s}_2 under $\pi_0(Z(\widehat{H}_1)^+) \rightarrow \pi_0(Z(\widehat{H}_2)^+)$.

Given an L -parameter $\varphi : W_{\mathbb{R}} \rightarrow {}^L G$ and a semi-simple element $s \in S_\varphi$, where $S_\varphi = \text{Cent}(\varphi, \widehat{G})$, one obtains a pure refined endoscopic datum as follows. Set $\widehat{H} = \text{Cent}(s, \widehat{G})^\circ$. The homomorphism $\varphi : W_{\mathbb{R}} \rightarrow \text{Cent}(s, \widehat{G}) \rightarrow \text{Out}(\widehat{H})$ factors through the projection $W_{\mathbb{R}} \rightarrow \Gamma$. There is a unique (up to isomorphism) quasi-split connected reductive \mathbb{R} -group H with dual group \widehat{H} such that the homomorphism $\Gamma \rightarrow \text{Out}(H) = \text{Out}(\widehat{H})$ induced by the \mathbb{R} -structure of H matches

the one induced by φ . Set $\mathcal{H} = \widehat{H} \cdot \varphi(W_{\mathbb{R}})$, and let η be the tautological inclusion $\mathcal{H} \rightarrow {}^L G$. In the rigid setting, the same construction works starting with $\dot{s} \in S_{\varphi}^+$, where S_{φ}^+ is the preimage in \widehat{G} of S_{φ} .

By construction the parameter φ takes values in \mathcal{H} . However, the extensions \mathcal{H} and ${}^L H$ of Γ by \widehat{H} need not be isomorphic, and even if they are, there is no natural isomorphism between them. Therefore, φ is *not* a parameter for H in any natural way. There are two ways to remedy this situation.

The classical approach is to choose a z -extension $H_1 \rightarrow H$ and an L -embedding $\mathcal{H} \rightarrow {}^L H_1$ that extends the natural embedding $\widehat{H} \rightarrow \widehat{H}_1$. These choices (which always exist, cf. [KS99, §2.2]) are called a z -pair. They provide a parameter φ_1 for H_1 .

An approach introduced in [Kal22] is to extend the theory of double covers of tori from [Kal19] to the setting of quasi-split connected reductive groups. The datum \mathcal{H} then leads to a *canonical* double cover $H(F)_{\pm}$ of $H(F)$ and a *canonical* isomorphism ${}^L H_{\pm} \rightarrow \mathcal{H}$. In this way, φ naturally becomes a parameter for $H(F)_{\pm}$.

The transfer of orbital integrals and characters between G and H is governed by the transfer factor. In the classical case, it is a function

$$\Delta : H_1(\mathbb{R})^{\text{rs}} \times G(\mathbb{R})^{\text{rs}} \rightarrow \mathbb{C}$$

that depends on the z -pair datum, while in the setting of covers it is a function

$$\Delta : H(R)_{\pm}^{\text{rs}} \times G(\mathbb{R})^{\text{rs}} \rightarrow \mathbb{C}$$

that is genuine in the first argument. In the classical case, it is given as the product

$$\epsilon \cdot \Delta_I^{-1} \Delta_{II} \Delta_{III_1}^{-1} \Delta_{III_2}.$$

The individual factors are defined in [LS87], except for ϵ , which is defined in the more general twisted setting in [KS99, §5.3], and Δ_{III_1} , whose relative definition is given in [LS87], but whose absolute definition is given in [Kal11] in the setting of pure inner forms, and in [Kal16] in the setting of rigid inner forms. The inverses appear due to the conventions of [KS, (1.0.4)], which we will use in this paper. The term Δ_{IV} is missing because we have normalized orbital integrals and characters by the Weyl denominator. We will not review the construction of the individual pieces here, as it has been reviewed in various other places, such as [Kal, §3.5, §4.2, §4.3]. The individual factors depend on auxiliary data, known as a -data and χ -data. The total factor depends on a choice of Whittaker datum, and z -datum.

In the case of covers, the transfer factor becomes the product

$$\epsilon \cdot \Delta_I^{-1} \cdot \Delta_{III}.$$

The terms Δ_I and Δ_{III} are slightly different from the original ones, and are defined in [Kal22, §4.3]. Neither of them depends on auxiliary data, although they are defined on certain covers of tori, and one could argue that the elements of those covers count as auxiliary data.

We now state the theorem asserting transfer of orbital integrals. It is a fundamental result of Shelstad, [She82], [She08a]. We state two versions, one using the cover $H(\mathbb{R})_{\pm}$ and one using a z -pair (H_1, η_1) .

Theorem 1.6.1. Let $f \in \mathcal{C}_c^\infty(G(\mathbb{R}))$.

1. There exists a genuine function $f^{H^\pm} \in \mathcal{C}_c^\infty(H(\mathbb{R})_\pm)$ such that for all $\dot{\gamma} \in H(F)_\pm^{rs}$

$$SO_{\dot{\gamma}}(f^{H^\pm}) = \sum_{\delta} \Delta(\dot{\gamma}, \delta) O_{\delta}(f).$$

2. Assume chosen a z -pair (H_1, η_1) . There exists a genuine function $f^{H_1} \in \mathcal{C}_c^\infty(H_1(\mathbb{R}))$ such that for all $\gamma_1 \in H_1(F)^{rs}$

$$SO_{\gamma_1}(f^{H_1}) = \sum_{\delta} \Delta(\gamma_1, \delta) O_{\delta}(f).$$

In both cases δ runs over the set of $G(\mathbb{R})$ -conjugacy classes in $G(\mathbb{R})^{rs}$.

Definition 1.6.2. The functions f and f^{H^\pm} are called *matching*. The functions f and f^{H_1} are called *matching*.

For the computations of this paper it would be useful to review the factor ϵ and compute it in the case of the base field \mathbb{R} , where the computation is rather straightforward. Consider the universal maximal torus T_0^G of G and T_0^H of H . The complexified character modules $V_G := X^*(T_0^G) \otimes_{\mathbb{Z}} \mathbb{C}$ and $V_H := X^*(T_0^H) \otimes_{\mathbb{Z}} \mathbb{C}$ are self-dual Artin representations of the same dimension. Choose a pinning of the quasi-split form G_0 of G and a non-trivial additive character $\Lambda : \mathbb{R} \rightarrow \mathbb{C}$, which combine to the Whittaker datum fixed for G_0 . Then

$$\epsilon = \epsilon(1/2, V_G - V_H, \Lambda),$$

where we have use Langlands' convention [Tat79, (3.6.4)] for the ϵ -factor. The pinning is used in the construction of G_0 . [May be better to review the construction of Δ_I and Δ_{III} here.]

Lemma 1.6.3. Let $\Lambda(x) = e^{irx}$ with $r > 0$. Then

$$\epsilon = (-1)^{q(H) - q(G_0)} i^{r_G/2 - r_H/2},$$

where r_G is the number of roots in the absolute root system of G , and r_H is the analogous number for H .

Proof. Let $A_0^G \subset T_0^G$ be the maximal split torus ($X_*(A_0^G) = X_*(T_0^G)^\Gamma$), and $S_0^G \subset T_0^G$ the maximal anisotropic torus ($X^*(S_0^G) = X^*(T_0^G)/X^*(T_0^G)^\Gamma$). Then $X^*(T_0^G)_{\mathbb{C}} = X^*(A_0^G)_{\mathbb{C}} \oplus X^*(S_0^G)_{\mathbb{C}}$, where we have abbreviated $\otimes_{\mathbb{Z}} \mathbb{C}$ by the subscript \mathbb{C} . One has $\epsilon(1/2, \mathbf{1}, \Lambda) = 1$ and $\epsilon(1/2, \text{sgn}, \Lambda) = i$ according to [Tat79, (3.2.4)], hence

$$\epsilon(1/2, V_G, \Lambda) = i^{d - \dim(A_0^G)},$$

where $d = \dim(T_0^G)$. We use the same computation for H and conclude

$$\epsilon = \frac{\epsilon(1/2, X^*(T_0^G)_{\mathbb{C}}, \Lambda)}{\epsilon(1/2, X^*(T_0^H)_{\mathbb{C}}, \Lambda)} = \frac{i^{d - \dim(A_0^G)}}{i^{d - \dim(A_0^H)}} = i^{\dim(A_0^H) - \dim(A_0^G)}.$$

The Iwasawa decomposition $\text{Lie}(G_0) = \mathfrak{a} \oplus \mathfrak{n} \oplus \mathfrak{k}$ shows $2q(G_0) = \dim(\mathfrak{a}) + \dim(\mathfrak{n}) = \dim(A_0^G) + r_G/2$. We note that $q(G_0)$ is an integer, because G_0 has an elliptic maximal torus, and $q(G_0)$ equals the number of positive non-compact roots with respect to any Weyl chamber. Therefore

$$\dim(A_0^H) - \dim(A_0^G) = 2(q(H) - q(G_0)) + (r_G/2 - r_H/2). \quad \square$$

This section was written to state all of the facts needed. It should be cleaned up, and the terminology/notation needs to be coordinate with the rest of the paper.

By a *Whittaker datum* we mean a $G(\mathbb{R})$ -conjugacy class of pairs $(B(\mathbb{R}), \eta)$ where $B(\mathbb{R})$ is a Borel subgroup of $G(\mathbb{R})$, and η is a non-degenerate character of the nilpotent radical $N(\mathbb{R})$ of $B(\mathbb{R})$. Non-degenerate means: non-trivial on each simple root space.

Let B be a Borel subgroup of G . Let N be the nilradical of B , and \bar{N} the opposite nilpotent subgroup. Let $\mathfrak{n}, \bar{\mathfrak{n}}$ be the Lie algebras of N, \bar{N} . If B is defined over \mathbb{R} we consider $N(\mathbb{R}), \mathfrak{n}(\mathbb{R}), \bar{\mathfrak{n}}(\mathbb{R})$, etc. Let $\kappa(\cdot, \cdot)$ be any bilinear form on \mathfrak{g} , restricting to the Killing form on the derived algebra.

For $X \in i\bar{\mathfrak{n}}(\mathbb{R})$ define a unitary character ψ_X of $N(\mathbb{R})$ by:

$$\psi_X(\exp(Y)) = e^{\kappa(X, Y)} \quad (Y \in \mathfrak{n}(\mathbb{R})).$$

Note that this definition requires a choice of $\sqrt{-1}$, which is equivalent to choosing the non-trivial additive character $e^{2\pi i x}$ of \mathbb{R} . The map $X \rightarrow \psi_X$ is an isomorphism between $i\bar{\mathfrak{n}}(\mathbb{R})$ and the unitary characters of $N(\mathbb{R})$.

Let $\mathcal{O}_p \subset \mathfrak{g}$ be the set of regular nilpotent element of \mathfrak{g} . This is a single G -orbit. Recall $\mathcal{O}_p \cap i\mathfrak{g}(\mathbb{R})$ is the union of finitely many $G(\mathbb{R})$ -orbits, and this has a simply transitive action of the finite group

$$\begin{aligned} Q(\mathbb{R}) &= \ker(H^1(\Gamma, Z) \rightarrow H^1(\Gamma, G)) \\ &\simeq G_{\text{ad}}(\mathbb{R})/G(\mathbb{R})_{\text{ad}}. \end{aligned}$$

For X nilpotent it is easy to see ψ_X is non-degenerate if and only if $X \in \mathcal{O}_p$, in which case X is contained in a unique Borel subalgebra. Suppose $X \in \mathcal{O}_p \cap i\mathfrak{g}(\mathbb{R})$. Let B be the opposite Borel subgroup, so ψ_X is a unitary character of $N(\mathbb{R})$. It follows that $X \mapsto (B(\mathbb{R}), \psi_X)$ is a bijection from $(\mathcal{O}_p \cap i\mathfrak{g}(\mathbb{R}))/G(\mathbb{R})$ to the set of Whittaker data. We write this bijection as:

$$X \mapsto \mathfrak{w}_X = (B(\mathbb{R}), \psi_X) \quad (X \in \mathcal{O}_p \cap i\mathfrak{g}(\mathbb{R})).$$

Suppose $\mathcal{O}_{i\mathbb{R}}$ is a $G(\mathbb{R})$ -orbit in $\mathcal{O}_p \cap i\mathfrak{g}(\mathbb{R})$. A *Kostant section* for $\mathcal{O}_{i\mathbb{R}}$ is obtained as follows. Choose an $SL(2)$ triple $(H_{\mathbb{R}}, iE_{\mathbb{R}}, iF_{\mathbb{R}})$ with $iE_{\mathbb{R}} \in \mathcal{O}_{i\mathbb{R}}, iF_{\mathbb{R}} \in i\mathfrak{g}(\mathbb{R})$ ($\Rightarrow H_{\theta} \in \mathfrak{g}(\mathbb{R})$), and set $V = iF_{\mathbb{R}} + \text{Cent}_{\mathfrak{g}}(iE_{\mathbb{R}})$ (we've multiplied the usual definition by i). Then V meets every regular orbit \mathcal{O} in a single point X , and if $\mathcal{O} \cap i\mathfrak{g}(\mathbb{R}) \neq \emptyset$ then $X \in i\mathfrak{g}(\mathbb{R})$.

There are two choices here. We choose $iE_{\mathbb{R}} \in \mathcal{O}_{i\mathbb{R}}$; obviously any two such choices are $G(\mathbb{R})$ -conjugate. Given $iE_{\mathbb{R}}$, we choose a triple $\{H_{\mathbb{R}}, iE_{\mathbb{R}}, iF_{\mathbb{R}}\}$; any two such triples are conjugate by $\text{Cent}_G(iE_{\mathbb{R}})$. It follows that the $G(\mathbb{R})$ -conjugacy class of V is unique up to conjugation by $G(\mathbb{R})$.

Lemma 2.0.1. *This defines a bijection between $(\mathcal{O}_p \cap i\mathfrak{g}(\mathbb{R}))/G(\mathbb{R})$ and the set of $G(\mathbb{R})$ -conjugacy class of Kostant sections. Composing with the map $X \rightarrow \mathfrak{w}_X$ gives a canonical bijection $(\mathcal{O}_p \cap i\mathfrak{g}(\mathbb{R}))/G(\mathbb{R})$ with the set of Whittaker data.*

Let σ be the Galois action defining $\mathfrak{g}(\mathbb{R})$, and choose a compatible Cartan involution θ . Write $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ where $\mathfrak{k}, \mathfrak{p}$ are the $1, -1$ eigenspaces of θ , respectively.

Note: this is one piece of the following result. We probably don't need all of this, but some of it might prove useful.

Proposition 2.0.2. *There are canonical bijections between the following sets.*

1. Large discrete series representations in Π
2. Whittaker data for $G(\mathbb{R})$
3. $(\mathcal{O}_p \cap \mathfrak{p})/K$
4. $(\mathcal{O}_p \cap i\mathfrak{g}(\mathbb{R}))/G(\mathbb{R})$
5. The set of $G(\mathbb{R})$ -conjugacy classes of Kostant sections.

The group $Q(\mathbb{R})$ acts simply transitively on each of these sets, and the bijections commute with this action.

Suppose $\pi = \pi(S, \tau)$ is a generic discrete series representation. Here S is a compact Cartan subgroup, with Lie algebra \mathfrak{s} . We identify $d\tau$ with an element $H = H_\pi \in i\mathfrak{s}(\mathbb{R})$ using κ . Choose an $SL(2)$ triple $\{iH, E, F\}$ with $E, F \in \mathfrak{p}$ (note that $iH \in \mathfrak{g}(\mathbb{R})$). After conjugating by K we may assume $\sigma(E) = F$. Set $F_\pi = \frac{1}{2}(E - F - iH) \in i\mathfrak{g}(\mathbb{R})$.

Lemma 2.0.3. *Suppose $\pi = \pi(S, \tau)$ is a generic discrete series representation. Then F_π is a regular nilpotent element of $i\mathfrak{g}(\mathbb{R})$, and the wave-front set of π is $G(\mathbb{R})F_\pi$. In particular the $G(\mathbb{R})$ orbit of F_π only depends on π .*

Proposition 2.0.4. *Suppose $\pi = \pi(S, \tau)$ is a generic discrete series representation. Let \mathfrak{w} be the unique Whittaker datum for which π has a \mathfrak{w} -Whittaker model. Write $\mathfrak{w} = \mathfrak{w}_X$ for some regular nilpotent element $X \in i\mathfrak{g}(\mathbb{R})$.*

Suppose $\gamma^\vee \in i\mathfrak{s}(\mathbb{R})$ is regular and dominant for the positive roots defined by $d\tau$. Then γ^\vee is $G(\mathbb{R})$ -conjugate to an element of the Kostant section of X .

The previous statement:

Proposition 2.0.5. *Let $S \subset G_0$ be an elliptic maximal torus, ρ a Weyl chamber in $X^*(S/Z_{G_0})$, and τ_0 a character of $S(\mathbb{R})$ whose differential is ρ -dominant and ρ -integral. Let $(T_0, B_0, \{X_\alpha\})$ be a pinning of G_0 and $\Lambda(x) = e^{2\pi i x}$. If the discrete series representation associated to (S, ρ, τ_0) is generic with respect to the Whittaker datum associated to the pinning and Λ , then the element $\rho^\vee(-i) \in \text{Lie}(S_{\text{sc}})(\mathbb{R})$ is $G_0(\mathbb{R})$ -conjugate to the Kostant section associated to the pinning.*

Proof. Note that $X = \rho^\vee(-i) \in \text{Lie}(S_{\text{sc}})$ is Galois-fixed and thus lies in $\text{Lie}(S_{\text{sc}})(\mathbb{R})$.
TODO □

3 CONSTRUCTION OF L -PACKET AND INTERNAL STRUCTURE

Let G_0 be a quasi-split connected reductive \mathbb{R} -group with dual group \widehat{G} and L -group ${}^L G$. Let $\varphi : W_{\mathbb{R}} \rightarrow {}^L G$ be a discrete Langlands parameter, given up to conjugation by \widehat{G} .

3.1 Factorization of a parameter

We choose a Γ -invariant Borel pair (\hat{T}, \hat{B}) of \hat{G} . Conjugating by \hat{G} we arrange that $\varphi(z) \in \hat{T}$ for all $z \in \mathbb{C}^\times$. Thus, $\varphi|_{\mathbb{C}^\times}$ is a continuous group homomorphism $\mathbb{C}^\times \rightarrow \hat{T}$. Every continuous group homomorphism $\mathbb{C}^\times \rightarrow \mathbb{C}^\times$ is of the form $z^a \bar{z}^b = |z|^{a+b} \arg(z)^{a-b}$ for some $a, b \in \mathbb{C}$ with $a - b \in \mathbb{Z}$. Thus there exist $\lambda, \mu \in X_*(\hat{T}) \otimes_{\mathbb{Z}} \mathbb{C}$ with $\lambda - \mu \in X_*(\hat{T})$ such that

$$\varphi(z) = \lambda(z) \cdot \mu(\bar{z}), \quad \forall z \in \mathbb{C}^\times.$$

Lemma 3.1.1. *Assume that G_0 is semi-simple and simply connected.*

1. $\lambda, \mu \in X_*(\hat{T})$
2. $\langle \lambda, \alpha \rangle \neq 0$ for all $\alpha \in R(\hat{T}, \hat{G})$.
3. $\mu = -\lambda$ in $X_*(\hat{T}/Z(\hat{G}))$

Proof. [TODO]

□

We can apply Lemma 3.1.1 to the composition of φ with the projection ${}^L G \rightarrow {}^L G/\hat{Z}$, where \hat{Z} is the center of \hat{G} , noting that \hat{G}/\hat{Z} is the dual group of G_{sc} . It implies that the centralizer of $\varphi|_{\mathbb{C}^\times}$ in \hat{G} equals \hat{T} . Since \mathbb{C}^\times is normal in $W_{\mathbb{R}}$, the image of φ lies in $N(\hat{T}, \hat{G})$. Its projection to $\Omega(\hat{T}, \hat{G})$ factors through a homomorphism $\xi : \Gamma \rightarrow \Omega(\hat{T}, \hat{G}) \rtimes \Gamma$. Let \hat{S} denote the Γ -module with underlying abelian group \hat{T} and Γ -structure given by $\text{Ad} \circ \xi$. Let S be the \mathbb{R} -torus whose dual is \hat{S} , i.e. the \mathbb{R} -torus determined by $X^*(S) = X_*(\hat{S})$ as Γ -modules.

By construction we have $R(\hat{T}, \hat{G}) \subset X^*(\hat{T}) = X^*(\hat{S}) = X_*(S)$, and we write $R^\vee(S, G)$ for this set. Analogously we have a subset $R(S, G) \subset X^*(S)$. Both of these subsets are Γ -stable. Moreover $\mu = \sigma\lambda$ in $X_*(\hat{S})$ and according to Lemma 3.1.1 the action of σ on $R(S, G)$ is by negation.

Let $S(\mathbb{R})_G$ be the double cover of $S(\mathbb{R})$ reviewed in §1.4, associated to the subset $R(S, G) \subset X^*(S)$. As discussed there, there is a canonical \hat{G} -conjugacy class of L -embeddings ${}^L S_G \rightarrow {}^L G$. Inside of this class, there is a unique \hat{S} -conjugacy class, call it ${}^L j$, whose restriction to \hat{S} is the tautological embedding $\hat{S} \rightarrow \hat{G}$. The image of this L -embedding is described in (1.5), and contains the image of φ by construction. Thus $\varphi = {}^L j \circ \varphi_S$ for a unique \hat{S} -conjugacy class of L -homomorphisms $\varphi_S : W_{\mathbb{R}} \rightarrow {}^L S_G$. According to [Kal19, Theorem 3.15], φ_S corresponds to a genuine character $\tau : S(\mathbb{R})_G \rightarrow \mathbb{C}^\times$.

The construction of (S, τ) depended on the choice of \hat{T} .

Lemma 3.1.2. *If (S_1, τ_1) and (S_2, τ_2) are two pairs obtained from two different choices of \hat{T} , there exists $g \in \hat{G}^\Gamma$ such that $\text{Ad}(g)\hat{S}_1 = \hat{S}_2$, $\text{Ad}(g) : \hat{S}_1 \rightarrow \hat{S}_2$ is Γ -equivariant, and its dual isomorphism $S_2 \rightarrow S_1$ identifies τ_2 with τ_1 .*

Proof. [TODO]

□

3.2 Construction of the L -packet

Construction 3.2.1. Let (G, ξ, z) be a pure (or rigid) inner form of G_0 . We construct a natural stable class J_ξ of embeddings $j : S \rightarrow G$. [TODO]

Consider $j \in J_\xi$ and use it to identify S with its image, an elliptic maximal torus of G . As discussed in §1.5, there exists a unique essentially discrete series representation π_j of $G(\mathbb{R})$ associated to the pair (S, τ) , transported to G via j .

Definition 3.2.2.

$$\Pi_\varphi((G, \xi, z)) = \{\pi_j | j \in J_\xi\}.$$

We define the pure (resp rigid) compound L -packet

$$\Pi_\varphi = \{(G, \xi, z, \pi) | (G, \xi, z) \in \mathcal{I}, \pi \in \Pi(G, \xi, z)\},$$

where \mathcal{I} is the category of pure (resp. rigid) inner forms of G_0 .

Lemma 3.2.3. *The set of representations $\Pi_\varphi((G, \xi, z))$ is independent of z and coincides with the set $\Pi_\varphi(G)$ constructed by Langlands in [Lan89, §3].*

Proof. [TODO] □

3.3 Internal structure of the compound packet

Let \mathcal{J} be the category of quadruples (G, ξ, z, j) , where (G, ξ, z) is a pure (resp. rigid) inner form of G_0 and $j \in J_\xi$. By construction we have a functor $\mathcal{J} \rightarrow \Pi_\varphi$.

Lemma 3.3.1. *The functor $\mathcal{J} \rightarrow \Pi_\varphi$ is an equivalence.*

Proof. [TODO] □

Recall from [ref] that the abelian group $H^1(\Gamma, S)$ in the pure case (resp. $H^1(u \rightarrow W, Z(G_0) \rightarrow S)$ in the rigid case) acts simply transitively on the set of isomorphism classes of \mathcal{J} , hence according to Lemma 3.3.1 also on the set of isomorphism classes of Π_φ . At the same time, Lemma 3.1.1 provides an identification $S_\varphi = \widehat{S}^\Gamma$, hence by Tate-Nakayama duality $\pi_0(S_\varphi)^* = \pi_0(\widehat{S}^\Gamma)^* = H^1(\Gamma, S)$. Analogously, in the rigid setting we obtain $\pi_0(S_\varphi^+)^* = H^1(u \rightarrow W, Z(G_0) \rightarrow S)$. This provides a simply transitive action of the abelian group $\pi_0(S_\varphi)^*$ in the pure setting, and $\pi_0(S_\varphi^+)^*$ in the rigid setting, on the set of isomorphism classes in Π_φ .

Lemma 3.3.2. *The set $\Pi((G_0, 1, 1))$ contains a unique \mathfrak{w} -generic member.*

Proof. [TODO] □

Taking the unique \mathfrak{w} -generic member of $\Pi((G_0, 1, 1)) \subset \Pi_\varphi / \sim$, provided by Lemma 3.3.2, as a base-point, the simply-transitive action turns into the desired bijection from $\pi_0(S_\varphi)^*$ (resp. $\pi_0(S_\varphi^+)^*$) to Π_φ / \sim .

3.4 The case of a cover of G

3.5 Dependence on the choice of Whittaker datum

[TODO]

4 ENDOSCOPIC CHARACTER IDENTITIES

Let $\varphi : W_{\mathbb{R}} \rightarrow {}^L G$ be a discrete parameter. Let $s \in S_{\varphi}$ (resp. $s \in S_{\varphi}^+$) be a semi-simple element. Let $(H, s, \mathcal{H}, \eta)$ be the (pure or rigid) refined endoscopic datum associated to the pair (φ, s) , whose construction was reviewed in §1.6.

4.1 Statement of the main theorem

As discussed in §3.4 there is an associated compound L -packet Π_{φ_H} . We will be only interested in the contribution of the trivial twist $(H, 1, 1)$, and we write $\Pi_{\varphi}(H) \subset \Pi_{\varphi_H}$ for it. Consider the virtual character

$$S\Theta_{\varphi_H} := \sum_{\sigma \in \Pi_{\varphi}(H)} \langle \sigma, s \rangle \Theta_{\sigma} = \sum_{\sigma \in \Pi_{\varphi}(H)} \langle \sigma, 1 \rangle \Theta_{\sigma} = \sum_{\sigma \in \Pi_{\varphi}(H)} \Theta_{\sigma}$$

on $H(\mathbb{R})_{\pm}$, where $\langle \sigma, - \rangle$ is the character of the irreducible representation of $\pi_0(S_{\varphi})$ (resp. $\pi_0(S_{\varphi}^+)$) associated to σ by the bijection of §3.3. Let us argue the two equalities. Since $Z(\widehat{H})^{\Gamma}$ (resp. $Z(\widehat{H})^+$) acts trivially on this irreducible representation, and s belongs by construction to this group, we see $\langle \sigma, s \rangle = \langle \sigma, 1 \rangle$, hence the first equality. The second comes from the fact that S_{φ} is abelian, because it lies in \widehat{S} (and S_{φ}^+ lies in \widehat{S}), where \widehat{S} is the torus involved in the construction of the L -packet on H . Note that, while the bijection of §3.3 depends on the choice of a Whittaker datum, the argument of §3.5 shows that the value $\langle \sigma, 1 \rangle$ does not depend on this choice.

Let (G, ξ, z) be a pure (resp. rigid) inner twist of G_0 . We have the virtual character on $G(\mathbb{R})$ given by

$$\Theta_{\varphi}^{\mathfrak{w}, s} := e(G) \sum_{\pi \in \Pi_{\varphi}((G, \xi, z))} \langle \pi, s \rangle \Theta_{\pi}.$$

This virtual character does depend on \mathfrak{w} .

The following is the main theorem of this article. It is a fundamental result of Shelstad [She82], [She10], [She08b].

Theorem 4.1.1. *Let $f \in \mathcal{C}_c^{\infty}(G(\mathbb{R}))$ be a test function.*

1. *If $f^{H_{\pm}} \in \mathcal{C}_c^{\infty}(H(\mathbb{R})_{\pm})$ matches f as in Definition 1.6.2, then*

$$\Theta_{\varphi}^{\mathfrak{w}, s}(f) = S\Theta_{\varphi_H}(f^{H_{\pm}}).$$

2. *If $f^{H_1} \in \mathcal{C}_c^{\infty}(H_1(\mathbb{R}))$ matches f as in Definition 1.6.2, then*

$$\Theta_{\varphi}^{\mathfrak{w}, s}(f) = S\Theta_{\varphi_{H_1}}(f^{H_1}).$$

4.2 Reduction to the elliptic set

Theorem 4.2.1. 1. For every strongly regular semi-simple element $\delta \in G(\mathbb{R})$ the following identity holds

$$\Theta_{\varphi}^{\mathfrak{w},s}(\delta) = \sum_{\gamma \in H(\mathbb{R})/st} \Delta[\mathfrak{w}, \mathfrak{e}, z](\dot{\gamma}, \delta) S\Theta_{\varphi_H}(\dot{\gamma}).$$

2. For every strongly regular semi-simple element $\delta \in G(\mathbb{R})$ the following identity holds

$$\Theta_{\varphi}^{\mathfrak{w},s}(\delta) = \sum_{\gamma \in H(\mathbb{R})/st} \Delta[\mathfrak{w}, \mathfrak{e}, \mathfrak{z}, z](\gamma_1, \delta) S\Theta_{\varphi_{H_1}}(\gamma_1).$$

Lemma 4.2.2. Theorem 4.1.1 is equivalent to Theorem 4.2.1.

Proof. [TODO] □

Lemma 4.2.3. If Theorem 4.2.1 holds for all elliptic δ , then it holds for all δ .

Proof. [TODO] □

4.3 The left hand side

In this subsection we will provide a formula for the left hand side of the identity in Theorem 4.2.1, i.e. $\Theta_{\varphi,s}^{\mathfrak{w}}(\delta)$, for strongly regular semi-simple elliptic $\delta \in G(\mathbb{R})$. The end result is (4.1).

The members of $\Pi_{\varphi}((G, \xi, z))$ are parameterized by the set of $G(\mathbb{R})$ -conjugacy classes of admissible embedding $j : S \rightarrow G$. Given such an embedding let π_j be the corresponding representation. Let $j_{\mathfrak{w}} : S \rightarrow G_0$ be the unique embedding for which $\pi_{j_{\mathfrak{w}}}$ is the unique \mathfrak{w} -generic member of $\Pi_{\varphi}((G_0, 1, 1))$. Then $\text{inv}(j_{\mathfrak{w}}, j) \in H^1(\Gamma, S) = \pi_0(\widehat{S}^{\Gamma})^* = \pi_0(S_{\varphi})^*$ equals ρ_{π_j} . Therefore the left hand side becomes

$$\Theta_{\varphi,s}^{\mathfrak{w}}(\delta) = e(G) \sum_j \langle s, \text{inv}(j_{\mathfrak{w}}, j) \rangle \Theta_{\pi_j}(\delta),$$

where j runs over (a set of representatives for) the set of $G(\mathbb{R})$ -conjugacy classes in J_{ξ} . Harish-Chandra's character formula (1.6) states

$$\Theta_{\pi_j}(\delta) = (-1)^{q(G)} \sum_{w \in W_{\mathbb{R}}(G, jS)} \frac{\tau'}{d'_{\tau}}(j^{-1}w^{-1}\delta),$$

where we have conjugated δ within $G(\mathbb{R})$ to land in $jS(\mathbb{R})$. Combining the two formulas and using $e(G) = (-1)^{q(G_0)-q(G)}$, we obtain

$$\Theta_{\varphi,s}^{\mathfrak{w}}(\delta) = (-1)^{q(G_0)} \sum_j \langle s, \text{inv}(j_{\mathfrak{w}}, j) \rangle \sum_{w \in W_{\mathbb{R}}(G, jS)} \frac{\tau'}{d'_{\tau}}(j^{-1}w^{-1}\delta).$$

Instead of conjugating δ to land in jS , we can conjugate j by $G(\mathbb{R})$ to achieve this, without changing π_j . With this shift in point of view we can combine the two sums and arrive at

$$\Theta_{\varphi,s}^{\mathfrak{w}}(\delta) = (-1)^{q(G_0)} \sum_j \langle s, \text{inv}(j_{\mathfrak{w}}, j) \rangle \frac{\tau'}{d'_{\tau}}(j^{-1}w^{-1}\delta),$$

where now the sum runs over the set of those $j \in J_\xi$ whose image contains δ .

As j runs over this set, $j_{\mathfrak{w}} j^{-1}(\delta)$ runs over the set of elements $\delta_0 \in S_{\mathfrak{w}}(\mathbb{R})$ that are stably conjugate to δ , where $S_{\mathfrak{w}} \subset G_0$ is the image of $j_{\mathfrak{w}}$, an elliptic maximal torus of G_0 . Moreover, $j_{\mathfrak{w}}$ transports $\text{inv}(j_{\mathfrak{w}}, j) \in H^1(\mathbb{R}, S)$ to $\text{inv}(\delta_0, \delta) \in H^1(\mathbb{R}, S_{\mathfrak{w}})$. So we arrive at

$$\Theta_{\varphi, s}^{\mathfrak{w}}(\delta) = (-1)^{q(G_0)} \sum_{\delta_0} \langle s_{\mathfrak{w}}, \text{inv}(\delta_0, \delta) \rangle \frac{\tau'_{\mathfrak{w}}}{d'_{\mathfrak{w}}}(\delta_0), \quad (4.1)$$

where the sum runs over the set of elements $\delta_0 \in S_{\mathfrak{w}}(\mathbb{R})$ that are stably conjugate to δ , and we have used the subscript \mathfrak{w} to indicate various transports under $j_{\mathfrak{w}} : S \rightarrow S_{\mathfrak{w}}$.

4.4 The right hand side: covers

In this subsection we will show that the right hand side of the identity of Theorem 4.2.1(1) is also equal to (4.1).

We begin by applying (4.1) to the group H , the parameter φ_H , and the trivial endoscopic element, and obtain

$$S\Theta_{\varphi_H}(\dot{\gamma}) = \sum_{\dot{\gamma}_0} \frac{\tau_H}{d_H}(\dot{\gamma}_0),$$

where we have fixed an embedding $j_H : S \rightarrow H$ for which the corresponding discrete series representation of $H(\mathbb{R})_{\pm}$ is generic with respect to some Whittaker datum and denote by subscript H the various transports under j_H , $\dot{\gamma}_0$ runs over the elements of $S_H(\mathbb{R})_{\pm}$ that are H -stably conjugate to $\dot{\gamma}$, and we have used $\tau_H/d_H = \tau'_H/d'_H$.

The right hand side of Theorem 4.2.1(1) then becomes

$$\sum_{\gamma \in H(\mathbb{R})/\text{st}} \Delta[\mathfrak{w}, \mathfrak{e}, z](\dot{\gamma}, \delta) \sum_{\dot{\gamma}_0} \frac{\tau_H}{d_H}(\dot{\gamma}_0).$$

The first sum runs over elements of $H(\mathbb{R})$ that are related to δ , up to stable conjugacy under H . Each such stable conjugacy class consists of regular semi-simple elliptic elements (because δ is such), and hence intersects $S_H(\mathbb{R})$. The second sum runs over elements $\dot{\gamma}_0 \in S_H(\mathbb{R})_{\pm}$ that lie in the H -stable class of the lift $\dot{\gamma} \in S_H(\mathbb{R})_{\pm}$ of γ . Since Δ is H -stably invariant in the first factor, its values at $\dot{\gamma}$ and $\dot{\gamma}_0$ are the same. We can combine the two sums together and obtain

$$\sum_{\gamma_0 \in S_H(\mathbb{R})} \Delta[\mathfrak{w}, \mathfrak{e}, z](\dot{\gamma}_0, \delta) \frac{\tau_H}{d_H}(\dot{\gamma}_0),$$

where now γ_0 runs over all elements of $S_H(\mathbb{R})$, equivalently all those that are related to δ , since the transfer factor vanishes for the others.

Having fixed the embeddings $j_{\mathfrak{w}}$ and j_H , they provide an isomorphism $S_{\mathfrak{w}} \rightarrow S_H$, and this isomorphism induces a bijection

$$\delta_0 \leftrightarrow \gamma_0$$

between the set of elements of $S_{\mathfrak{w}}(\mathbb{R})$ that are stably conjugate to δ and the set of elements of $S_H(\mathbb{R})$ related to δ . Using the basic property of transfer factors [ref](#) we obtain

$$\sum_{\delta_0 \in S_{\mathfrak{w}}(\mathbb{R})} \Delta[\mathfrak{w}, \mathfrak{e}, z](\dot{\gamma}_0, \delta_0) \langle s_{\mathfrak{w}}, \text{inv}(\delta_0, \delta) \rangle \frac{\tau_H}{d_H}(\dot{\gamma}_0), \quad (4.2)$$

where δ_0 runs over the elements of $S_{\mathfrak{w}}(\mathbb{R})$ that are stably conjugate to δ , $\gamma_0 \in S_H(\mathbb{R})$ denotes the element corresponding to δ_0 under above bijection, and $\dot{\gamma}_0 \in S_H(\mathbb{R})_{\pm}$ is an arbitrary lift of γ_0 .

We now unpack the transfer factor. It is given as

$$\Delta(\dot{\gamma}_0, \delta_0) = \epsilon \Delta_I^{-1}(\dot{\gamma}_0, \dot{\delta}_0) \Delta_{III}(\dot{\gamma}_0, \dot{\delta}_0),$$

where we recall that the term Δ_{IV} is missing because we are working with normalized characters and orbital integrals, and $\dot{\delta}_0 \in S_{\mathfrak{w}}(\mathbb{R})_{G/H}$ is an arbitrary lift of δ_0 .

By construction, [\[ref\]](#)

$$\Delta_{III}(\dot{\gamma}_0, \dot{\delta}_0) = \frac{\tau_{\mathfrak{w}}}{\tau_H}(\dot{\delta}_0).$$

The following lemma completes the proof of Theorem 4.2.1(1).

Lemma 4.4.1. *For any $\dot{\delta}_0 \in S_{\mathfrak{w}}(\mathbb{R})_{G/H}$, the following identity holds*

$$\Delta_I(\dot{\gamma}_0, \dot{\delta}_0) = \epsilon \cdot (-1)^{q(G_0) - q(H)} \cdot \prod_{\alpha \in R(S, G/H)^+} \arg(\alpha^{1/2}(\dot{\delta}_0) - \alpha^{-1/2}(\dot{\delta}_0)),$$

where $\dot{\gamma}_0 \in S_H(\mathbb{R})_{G/H}$ is the image of $\dot{\delta}_0$ under the isomorphism $j_H \circ j_{\mathfrak{w}}^{-1}$.

Proof. We first investigate how both sides vary as functions of $\dot{\delta}_0$. Consider another strongly regular $\dot{\delta}_1$. Replacing $\dot{\delta}_0$ with $\dot{\delta}_1$ in the right hand side results in multiplication by $\prod_{\alpha} \arg(b_{\alpha})$, where the product runs again over $R(S, G/H)^+$ and $b_{\alpha} = (\dot{\delta}_{1,\alpha} - \dot{\delta}_{1,-\alpha})/(\dot{\delta}_{0,\alpha} - \dot{\delta}_{0,-\alpha})$. By construction $\dot{\delta}_{0,-\alpha} = \sigma(\dot{\delta}_{0,\alpha})$, and the same holds for $\dot{\delta}_1$, from which follows $b_{\alpha} \in \mathbb{R}^{\times}$, and hence $\arg(b_{\alpha}) = \text{sgn}(b_{\alpha})$.

We now look at the left hand side. Replacing $\dot{\delta}_0$ by $\dot{\delta}_1$ multiplies $\text{inv}(\dot{\delta}_0, \text{pin})$ by $\prod_{\alpha > 0, \sigma\alpha < 0} \alpha^{\vee}(b_{\alpha})$, and hence Δ_I by the Tate-Nakayama pairing of this 1-cocycle with the endoscopic element $s_{\mathfrak{w}}$. Since the torus S is elliptic, the conditions $\alpha > 0$ and $\sigma\alpha < 0$ are equivalent, and the value of the 1-cocycle at σ equals $\prod_{\alpha > 0} \alpha^{\vee}(b_{\alpha})$. This is the product over $\alpha > 0$ of the images of the 1-cocycles $b_{\alpha} \in Z^1(\Gamma, R_{\mathbb{C}/R}^1 \mathbb{G}_m)$ under the homomorphisms $\alpha^{\vee} : R_{\mathbb{C}/R}^1 \mathbb{G}_m \rightarrow S$, so the change in $\Delta_{I,\pm}$ is given by $\prod_{\alpha > 0} \langle b_{\alpha}, s_{\alpha} \rangle$, where s_{α} is the image of $s \in \widehat{S}$ under $\widehat{\alpha} : \widehat{S} \rightarrow \mathbb{C}^{\times}$. This is a Galois-equivariant homomorphism, with σ acting as inversion on \mathbb{C}^{\times} . Since s is σ -fixed, so is s_{α} , i.e. $s_{\alpha} \in \{\pm 1\} \subset \mathbb{C}^{\times}$. By construction of the endoscopic group H , we have $s_{\alpha} = 1$ precisely for $\alpha \in R(S, H)$. On the other hand, when $s_{\alpha} = -1$, then $\langle b_{\alpha}, s_{\alpha} \rangle = \text{sgn}(b_{\alpha})$.

We have thus shown that both sides of the identity multiply by the same factor upon replacing $\dot{\delta}_0$ by a different element $\dot{\delta}_1$. To establish the identity we may thus evaluate at an arbitrary element $\dot{\delta}_0$. For this we note that both sides descend to functions on $S_{\text{ad}}(\mathbb{R})_{G/H}$, so we may assume that G is adjoint. Let $\rho^{\vee} \in X_*(S_{\mathfrak{w}})$ denote half the sum of the coroots that pair positively with $d\tau_{\mathfrak{w}}$. Since complex conjugation acts on $X_*(S_{\mathfrak{w}})$ by multiplication by -1 , we have

$X = \rho^\vee(-ir) \in \text{Lie}(S_{\mathfrak{w}})(\mathbb{R})$ for any $r \in \mathbb{R}$. We will choose $r > 0$ small enough and set $\check{\delta} = \exp(X) \in S_{\mathfrak{w}}(\mathbb{R})_{\pm\pm}$, where we are using the exponential map $\text{Lie}(S_{\mathfrak{w}})(\mathbb{R}) \rightarrow S_{\mathfrak{w}}(\mathbb{R})_{\pm\pm}$ discussed in [Kal19, §3.7].

Considering the right hand side, we have for each $\alpha \in R(S, G)$

$$\alpha^{1/2}(\dot{\delta}_0) = \dot{\delta}_{0,\alpha} = \exp(d\alpha(X)/2) = e^{-ir\langle d\alpha, \rho^\vee \rangle/2}.$$

Then

$$\alpha^{1/2}(\dot{\delta}_0) - \alpha^{-1/2}(\dot{\delta}_0) = -2i \sin(r\langle d\alpha, \rho^\vee \rangle/2).$$

Choosing $r > 0$ so that $r\langle d\alpha, \rho^\vee \rangle/2 < \pi$ for all $\alpha \in R(S, G)^+$ we obtain

$$\arg(\alpha^{1/2}(\dot{\delta}_0) - \alpha^{-1/2}(\dot{\delta}_0)) = (-i)^{\#R(S, G/H)^+}.$$

Choose a pinning $(T_0, B_0, \{X_\alpha\})$, which, together with $\Lambda(x) = e^{2\pi i x}$, produces the chosen Whittaker datum \mathfrak{w} . Proposition 2.0.5 then shows that $\rho^\vee(-i)$ is $G_0(\mathbb{R})$ -conjugate lies in the Kostant section of that pinning. Thus X is $G_0(\mathbb{R})$ -conjugate to the Kostant section of the pinning $(T_0, B_0, \{rX_\alpha\})$. If we construct $\Delta_I(\dot{\gamma}_0, \dot{\delta}_0)$ with respect to this rescaled pinning, then [Kal22, Lemma 4.1.4] show that $\Delta_I(\dot{\gamma}_0, \dot{\delta}_0) = 1$. On the other hand, the rescaled pinning together with the character $\Lambda_r(x) = e^{r2\pi i x}$ also produces the Whittaker datum \mathfrak{w} . According to Lemma 1.6.3 the right hand side equals 1. \square

4.5 The right hand side: classical set-up

In this subsection we will show that the right hand side of the identity of Theorem 4.2.1(2) is also equal to (4.1). The initial arguments of §4.4, which applied to the right hand side of the identity in Theorem 4.2.1(1), have direct analogs in the setting of Theorem 4.2.1(2) and show that the right hand side of that identity equals the analog of the expression (4.2), which is given by

$$\sum_{\delta_0 \in S_{\mathfrak{w}}(\mathbb{R})} \Delta[\mathfrak{w}, \mathfrak{e}, z](\gamma_1, \delta_0) \langle s_{\mathfrak{w}}, \text{inv}(\delta_0, \delta) \rangle \frac{\tau'_{H_1}}{d'_{H_1}}(\gamma_1), \quad (4.3)$$

where we have used the parameter φ_{H_1} of the z -extension H_1 to obtain the character τ'_{H_1} of the torus $S_{H_1}(\mathbb{R})$.

It is the handling of the transfer factor that is slightly different. Indeed, in this setting without covers the transfer factor is given by

$$\Delta = \epsilon \Delta_I^{-1} \Delta_{II} \Delta_{III_2}.$$


The construction of the pieces involves a choice of an admissible isomorphism $S^H \rightarrow S_{\mathfrak{w}}$, which we take to be $j_{\mathfrak{w}} \circ j_H^{-1}$, as we did in §4.4. It further involves choices of χ -data and a -data for S . We take ρ -based χ -data, and $(-\rho)$ -based a -data, so that $\chi_\alpha(x) = \arg(x)$ when $\alpha > 0$ and $a_\alpha = i$ when $\alpha < 0$.

We claim that

$$\Delta_{III_2}(\gamma_1, \delta_0) = \frac{\tau'_{\mathfrak{w}}(\delta_0)}{\tau'_{H_1}(\gamma_1)}.$$

We will explain this under the assumption that the z -pair is trivial, i.e. there exists an L -isomorphism ${}^L\eta : {}^LH \rightarrow \mathcal{H}$, the general case being entirely analogous by requiring more cumbersome notation. We then have the commutative

diagram



where the horizontal arrows are the L -embeddings obtained via ρ -based and ρ^H -based χ -data, respectively. We have $\Delta_{II_2}(\gamma_1, \delta_0) = \langle a, \delta_0 \rangle$, where $a \in Z^1(W_{\mathbb{R}}, \widehat{S})$ is the 1-cocycle that makes the above diagram commute, the pairing is the Langlands pairing, and $\delta_0 \in S(\mathbb{R})$ is the image of $\gamma_0 \in S^H(\mathbb{R})$ to $S(\mathbb{R})$ under the chosen fixed admissible isomorphism. The claim now follows from the above commutative diagram and the fact that τ and τ^H are the characters with parameters φ_S and φ_S^H .

Next we consider

$$\frac{\Delta_{II}(\gamma_0, \delta)}{d'_H(\gamma_0)}.$$

By definition,

$$\Delta_{II}(\gamma_0, \delta) = \frac{\Delta_{II}^G(\gamma_0, \delta)}{\Delta_{II}^H(\gamma_0, \delta)}.$$

With the chosen a -data and χ -data we have

$$\Delta_{II}^H(\gamma_0) = \prod_{\substack{\alpha \in R(S^H, H) \\ \langle \alpha, \rho^H \rangle > 0}} \arg\left(\frac{\alpha(\gamma_0) - 1}{-i}\right) = i^{\#R(S^H, H)/2} \cdot \prod_{\substack{\alpha \in R(S^H, H) \\ \langle \alpha, \rho^H \rangle > 0}} \arg(\alpha(\gamma_0) - 1).$$

On the other hand,

$$(\alpha(\gamma_0) - 1)(1 - \alpha(\gamma_0)^{-1}) = \alpha(\gamma_0) + \alpha(\gamma_0)^{-1} - 2 = 2(\operatorname{Re}(\alpha(\gamma_0)) - 1) < 0.$$

Hence

$$\Delta_{II}^H(\gamma_0, \delta) d'_H(\gamma_0) = (-i)^{\#R(S^H, H)/2}.$$

In the same way one shows

$$\Delta_{II}^G(\gamma_0, \delta) d'_{G_0}(\delta_0) = (-i)^{\#R(S, G_0)/2}.$$

and we conclude

$$\frac{\Delta_{II}(\gamma_0, \delta)}{d'_H(\gamma_0)} = \frac{i^{\#R(S^H, H)/2 - \#R(S, G_0)/2}}{d'_{G_0}(\delta_0)}.$$

With this (4.3) becomes

$$(-1)^{q(H)} i^{\#R(S^H, H)/2 - \#R(S, G_0)/2} \epsilon \sum_{\delta_0 \in S_{\mathfrak{w}}(\mathbb{R})} \Delta_I(\gamma_0, \delta)^{-1} \langle s_{\mathfrak{w}}, \operatorname{inv}(\delta_0, \delta) \rangle \cdot \frac{\tau_{\mathfrak{w}}(\delta_0)}{d'_{G_0}(\delta_0)}.$$

We have

$$\Delta_I(\gamma_0, \delta)^{-1} = \langle s, \lambda \rangle^{-1}$$

where $\lambda \in H^1(\Gamma, S)$ is the splitting invariant of S relative to the chosen a -data. Using Lemma 1.6.3 we obtain

$$(-1)^{q(G_0)} \langle s, \lambda \rangle^{-1} \sum_{\delta_0} \langle s, \text{inv}(\delta_0, \delta) \rangle \cdot \frac{\tau_{\text{w}}(\delta_0)}{d'_{G_0}(\delta_0)}.$$

The following lemma completes the proof.

Lemma 4.5.1. *The splitting invariant λ of S computed in terms of $(-\rho)$ -based a -data is trivial.*

Proof. Let $X = \rho^\vee(-i) \in \text{Lie}(S_{\text{sc}})$. This element is Galois-fixed and thus lies in $\text{Lie}(S_{\text{sc}})(\mathbb{R})$. For every $\alpha > 0$ the complex number $d\alpha(X)$ is a positive real multiple of $-i$. Therefore we can replace the $(-\rho)$ -based a -data with the a -data $d\alpha(X)$ without changing the splitting invariant. According to [Kot99, Theorem 5.1] and Proposition 2.0.5, λ is trivial. \square

REFERENCES

- [AV92] Jeffrey Adams and David A. Vogan, Jr., *L-groups, projective representations, and the Langlands classification*, Amer. J. Math. **114** (1992), no. 1, 45–138. MR 1147719 (93c:22021)
- [AV16] ———, *Contragredient representations and characterizing the local Langlands correspondence*, Amer. J. Math. **138** (2016), no. 3, 657–682. MR 3506381
- [Kal] Tasho Kaletha, *Lectures on the stable trace formula, with emphasis on SL_2* , available on author’s website.
- [Kal11] ———, *Endoscopic character identities for depth-zero supercuspidal L -packets*, Duke Math. J. **158** (2011), no. 2, 161–224. MR 2805068 (2012f:22031)
- [Kal16] ———, *Rigid inner forms of real and p -adic groups*, Ann. of Math. (2) **184** (2016), no. 2, 559–632. MR 3548533
- [Kal19] ———, *On L -embeddings and double covers of tori over local fields*, arXiv:1907.05173 (2019).
- [Kal22] ———, *Covers of reductive groups and functoriality*, arXiv:2209.14357 (2022).
- [Kot84] Robert E. Kottwitz, *Stable trace formula: cuspidal tempered terms*, Duke Math. J. **51** (1984), no. 3, 611–650. MR 757954 (85m:11080)
- [Kot86] ———, *Stable trace formula: elliptic singular terms*, Math. Ann. **275** (1986), no. 3, 365–399. MR 858284 (88d:22027)
- [Kot99] ———, *Transfer factors for Lie algebras*, Represent. Theory **3** (1999), 127–138 (electronic). MR 1703328 (2000g:22028)
- [KS] Robert E. Kottwitz and Diana Shelstad, *On splitting invariants and sign conventions in endoscopic transfer*, arXiv:1201.5658.
- [KS99] ———, *Foundations of twisted endoscopy*, Astérisque (1999), no. 255, vi+190. MR 1687096 (2000k:22024)

- [Lan89] R. P. Langlands, *On the classification of irreducible representations of real algebraic groups*, Representation theory and harmonic analysis on semisimple Lie groups, Math. Surveys Monogr., vol. 31, Amer. Math. Soc., Providence, RI, 1989, pp. 101–170. MR 1011897 (91e:22017)
- [LS87] R. P. Langlands and D. Shelstad, *On the definition of transfer factors*, Math. Ann. **278** (1987), no. 1-4, 219–271. MR 909227 (89c:11172)
- [She82] D. Shelstad, *L-indistinguishability for real groups*, Math. Ann. **259** (1982), no. 3, 385–430. MR 661206 (84c:22017)
- [She08a] ———, *Tempered endoscopy for real groups. I. Geometric transfer with canonical factors*, Representation theory of real reductive Lie groups, Contemp. Math., vol. 472, Amer. Math. Soc., Providence, RI, 2008, pp. 215–246. MR 2454336 (2011d:22013)
- [She08b] ———, *Tempered endoscopy for real groups. III. Inversion of transfer and L-packet structure*, Represent. Theory **12** (2008), 369–402. MR 2448289 (2010c:22016)
- [She10] ———, *Tempered endoscopy for real groups. II. Spectral transfer factors*, Automorphic forms and the Langlands program, Adv. Lect. Math. (ALM), vol. 9, Int. Press, Somerville, MA, 2010, pp. 236–276. MR 2581952
- [Spr09] T. A. Springer, *Linear algebraic groups*, second ed., Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2009. MR 2458469
- [Tat79] J. Tate, *Number theoretic background*, Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 3–26. MR 546607 (80m:12009)
- [Vog93] David A. Vogan, Jr., *The local Langlands conjecture*, Representation theory of groups and algebras, Contemp. Math., vol. 145, Amer. Math. Soc., Providence, RI, 1993, pp. 305–379. MR 1216197 (94e:22031)