

# Funnel web spiders are a menace

Jeffrey O. Hanson\*<sup>1</sup>

<sup>1</sup>School of Biological Sciences, The University of Queensland, Saint Lucia, QLD, Australia  
\*email jeffrey.hanson@uqconnect.edu.au

16 March 2021

## MANUSCRIPT INFORMATION

Format: Journal of Applied Ecology (article)

Abstract word count: ? / 350

Main text word count: ? / 7,000 (including captions and references)

Number of references: ?

Number of figures: 1 / 6

Keywords: spiders, spiders, spiders, spiders, spiders, spiders, spiders, spiders

Spelling: American English (en-US)

## 13 ABSTRACT

14 Funnel web spiders are ferocious beasts that terrorize humans across Australia. In particular, they  
15 wreak havoc on the local tourism industry of Fraser Island. Here we examine the population of  
16 funnel web spiders found along a walking path from a camp site in Fraser Island to Lake Boomanjin.  
17 To achieve this, we subdivided the walking path into multiple transects and surveyed each transect  
18 multiple times over eight days. We then used open  $n$ -mixture models to estimate population size  
19 within each transect. Our models indicate that over 800 spiders could occur along this walking track.  
20 Fortunately, our models suggest that few spiders occur near the camp site – meaning that tourists  
21 may be relatively safe as long as they do not stray too far from this camp site. In conclusion, the  
22 dense infestation of spiders poses a serious threat to the safety of tourists and we recommend for  
23 their immediate eradication.

## 1 | INTRODUCTION

Fraser island is a popular tourist attraction. It is home to a great many interesting flora and fauna. In addition, it has several famous sites. The most famous site, perhaps, is Lake Boomanjin. This magnificent water feature is visited by scores of individuals throughout the year. However, the walking track to Lake Boomanjin is home by a population of funnel web spiders. This population may pose a serious safety risk to tourists. Here, we aimed to map the distribution of these hairy beasts along the walking track. Additionally – using the size of the spider holes as a proxy for age – we also aimed to understand the age-class structure of the population.

## 2 | MATERIALS AND METHODS

### 2.1 | Study system

Our study system was Fraser Island, Australia. Google it.

### 2.2 | Data collection

The Boomanjin Walking track was divided into 16 transects (each approximately 370 m in length; Figure 1). Each transect was surveyed for spider holes once a day over eight days (11–14/07/2016, 16–20/07/2016, see Table S1 for details). These surveys were conducted by a team consisting of two to three expert spider hunters. The size of each hole was measured to the nearest millimeter. The start and end times for each survey was recorded to assess search effort.

*Insert Figure 1 here.*

## 2.3 | Statistical analysis

Open  $n$ -mixture models were used to estimate the population size (Royle 2004). These models estimate the probability of detecting a spider in a given transect independent of the total number of spiders in the transect. Models were fit using the *R2jags* *R* package (Su & Masanao Yajima 2015) (100 total iterations; 50 burnin iterations; thinned by 2 iterations; 2). Model convergence was assessed using  $\hat{R}$  values. All analyses were conducted in *R* (version 4.0.3; R Core Team 2016).

## 3 | RESULTS

A total of 1639 spider holes were found. An average of 19.9375 (33.51 standard deviation [S.D.]) spider holes were found in each visit to a given transect. Typically, transects were surveyed for 28.32 minutes (17.98 S.D.). Based on the open  $n$ -mixture model, the probability of detecting a spider hole in a given transect was estimated to be 23.85% (0.01 S.D.). The total population size along the track was estimated to be 1341.58 (18.94 S.D.).

The size of the spider holes were bimodally distributed (Figure 2). The spider holes tended to range between 0–10 mm and 20–35 mm. These results suggest that most of the spiders surveyed were either juveniles or adults.

*Insert Figure 2 here.*

The spider holes were mostly found in a single transect (transect 11 contained 17.66 % of the population; Figure 3).

*Insert Figure 3 here.*

## 4 | DISCUSSION

Clearly, there are too many spiders. These results speak for themselves—no further discussion is necessary.

## ACKNOWLEDGEMENTS

JOH was supported by a chair while working on this manuscript. I thank all individuals involved with collecting the data.

## AUTHORS' CONTRIBUTIONS

JOH conceived the study, performed the analyses, and drafted the manuscript. All authors contributed critically to writing the manuscript and gave final approval for publication.

## DATA AVAILABILITY STATEMENT

Pending publication, code, data, and results will be made available via the Zenodo Digital Repository <https://doi.org/10.5281/zenodo.4607130> (Hanson 2021).

## ORCID

Jeffrey O. Hanson: 0000-0002-4716-6134

## REFERENCES

- Hanson, J.O. (2021). *Code and data for: Funnel web spiders are a menace*. Zenodo Digital Repository, <https://doi.org/10.5281/zenodo.4607130>.
- R Core Team. (2016). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>.
- Royle, J.A. (2004). N-mixture models for estimating population size from spatially replicated counts. *Biometrics*, **60**, 108–115.
- Su, Y.-S. & Masanao Yajima. (2015). *R2jags: Using R to Run 'JAGS'*. R package version 0.5-7, <https://CRAN.R-project.org/package=R2jags>.