

Speed Limits for Quantum Control of Local Spin Systems

Jeffrey M. Epstein^{1,3,4} and K. Birgitta Whaley^{2,3,4}

¹Department of Physics

²Department of Chemistry

³Berkeley Center for Quantum Information and Computation (BQIC)

⁴Center for Quantum Coherent Sciences (CQCS)

University of California, Berkeley

February 23, 2017

1. Motivation for bounds on optimal control
2. Brief review of Lieb-Robinson bounds
3. Bounds on state transfer and entanglement generation
4. Comparison to theory and numerics

Numerical upper bound on state transfer speed

Speed Limits for
Quantum Control
of Local Spin
Systems


Jeffrey M. Epstein
and K. Birgitta
Whaley

¹Caneva et al. arXiv:0902.4193, Murphy et al. arXiv:1004.3445

Numerical upper bound on state transfer speed


Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley



¹Caneva et al. arXiv:0902.4193, Murphy et al. arXiv: 1004.3445


Numerical upper bound on state transfer speed



$$H = -\frac{J}{2} \sum_j \vec{\sigma}_j \cdot \vec{\sigma}_{j+1} + h_j(t) \sigma_j^z$$

¹Caneva et al. arXiv:0902.4193, Murphy et al. arXiv: 1004.3445

Numerical upper bound on state transfer speed




$$H = -\frac{J}{2} \sum_j \vec{\sigma}_j \cdot \vec{\sigma}_{j+1} + h_j(t) \sigma_j^z$$

How fast can this be
achieved?

¹Caneva et al. arXiv:0902.4193, Murphy et al. arXiv: 1004.3445

Numerical upper bound on state transfer speed




Numerical
optimization¹(Krotov):

$$H = -\frac{J}{2} \sum_j \vec{\sigma}_j \cdot \vec{\sigma}_{j+1} + h_j(t) \sigma_j^z$$

How fast can this be
achieved?

¹Caneva et al. arXiv:0902.4193, Murphy et al. arXiv:1004.3445


Numerical upper bound on state transfer speed



$$H = -\frac{J}{2} \sum_j \vec{\sigma}_j \cdot \vec{\sigma}_{j+1} + h_j(t) \sigma_j^z$$


How fast can this be
achieved?

Numerical
optimization¹(Krotov):



¹Caneva et al. arXiv:0902.4193, Murphy et al. arXiv:1004.3445


Numerical upper bound on state transfer speed



$$H = -\frac{J}{2} \sum_j \vec{\sigma}_j \cdot \vec{\sigma}_{j+1} + h_j(t) \sigma_j^z$$

How fast can this be
achieved?

Numerical
optimization¹(Krotov):



$$T_{\min} \approx R/2J$$

¹Caneva et al. arXiv:0902.4193, Murphy et al. arXiv:1004.3445

Quantum Control in High and Low Dimension

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

$d = \dim(\mathcal{H}) \simeq 1$	$d \gg 1$

Quantum Control in High and Low Dimension

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

$d = \dim(\mathcal{H}) \simeq 1$	$d \gg 1$
analytical optimal control results	

Quantum Control in High and Low Dimension

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

$d = \dim(\mathcal{H}) \simeq 1$	$d \gg 1$
analytical optimal control results	numerics and lower bounds

Quantum Control in High and Low Dimension

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

$d = \dim(\mathcal{H}) \simeq 1$	$d \gg 1$
analytical optimal control results	numerics and lower bounds
“bare Hilbert space bounds”: Margolus-Levitin, Mandelstam-Tamm	

Quantum Control in High and Low Dimension

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

$d = \dim(\mathcal{H}) \simeq 1$	$d \gg 1$
analytical optimal control results	numerics and lower bounds
“bare Hilbert space bounds”: Margolus-Levitin, Mandelstam-Tamm	spatial structure enforced by experimental limitations

Quantum Control in High and Low Dimension

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

$d = \dim(\mathcal{H}) \simeq 1$	$d \gg 1$
analytical optimal control results	numerics and lower bounds
“bare Hilbert space bounds”: Margolus-Levitin, Mandelstam-Tamm	spatial structure enforced by experimental limitations → Lieb-Robinson

Quantum Control in High and Low Dimension

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

$d = \dim(\mathcal{H}) \simeq 1$	$d \gg 1$
analytical optimal control results	numerics and lower bounds
“bare Hilbert space bounds”: Margolus-Levitin, Mandelstam-Tamm	spatial structure enforced by experimental limitations → Lieb-Robinson
target specific unitary	

Quantum Control in High and Low Dimension

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

$d = \dim(\mathcal{H}) \simeq 1$	$d \gg 1$
analytical optimal control results	numerics and lower bounds
“bare Hilbert space bounds”: Margolus-Levitin, Mandelstam-Tamm	spatial structure enforced by experimental limitations → Lieb-Robinson
target specific unitary	target specific task

Quantum Control in High and Low Dimension

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

$d = \dim(\mathcal{H}) \simeq 1$	$d \gg 1$
analytical optimal control results	numerics and lower bounds
“bare Hilbert space bounds”: Margolus-Levitin, Mandelstam-Tamm	spatial structure enforced by experimental limitations → Lieb-Robinson
target specific unitary	target specific task

These are different problems that require different techniques!


Outline

1. Motivation for bounds on optimal control
2. Brief review of Lieb-Robinson bounds
3. Bounds on state transfer and entanglement generation
4. Comparison to theory and numerics

Local Spin Systems and Lieb-Robinson Bounds

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley



Local Spin Systems and Lieb-Robinson Bounds

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

The Finite Group Velocity of Quantum Spin Systems


Elliott H. Lieb*

Dept. of Mathematics, Massachusetts Institute of Technology
Cambridge, Massachusetts, USA

Derek W. Robinson**

Dept. of Physics, Univ. Aix-Marseille II, Marseille-Luminy, France

Received May 15, 1972



Local Spin Systems and Lieb-Robinson Bounds

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

The Finite Group Velocity of Quantum Spin Systems

Elliott H. Lieb*


Dept. of Mathematics, Massachusetts Institute of Technology
Cambridge, Massachusetts, USA

Derek W. Robinson**

Dept. of Physics, Univ. Aix-Marseille II, Marseille-Luminy, France

Received May 15, 1972

$$\|[A(t), B]\| \leq c_t(X, Y) \|A\| \|B\|$$



Local Spin Systems and Lieb-Robinson Bounds

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

The Finite Group Velocity of Quantum Spin Systems

Elliott H. Lieb*

Dept. of Mathematics, Massachusetts Institute of Technology
Cambridge, Massachusetts, USA

Derek W. Robinson**

Dept. of Physics, Univ. Aix-Marseille II, Marseille-Luminy, France

Received May 15, 1972

$$\|[A(t), B]\| \leq c_t(X, Y) \|A\| \|B\|$$

$$c_t(X, Y) \leq \begin{cases} c \min(|X|, |Y|) e^{-\frac{L-vt}{\xi}} \end{cases}$$

Local Spin Systems and Lieb-Robinson Bounds

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

The Finite Group Velocity of Quantum Spin Systems

Elliott H. Lieb*

Dept. of Mathematics, Massachusetts Institute of Technology
Cambridge, Massachusetts, USA

Derek W. Robinson**

Dept. of Physics, Univ. Aix-Marseille II, Marseille-Luminy, France

Received May 15, 1972


$$\|[A(t), B]\| \leq c_t(X, Y) \|A\| \|B\|$$

$$c_t(X, Y) \leq \begin{cases} c \min(|X|, |Y|) e^{-\frac{L-Yt}{\xi}} \\ 2 \sum_{n=1}^{\infty} \frac{(2Jt)^n}{n!} N(n) \end{cases}$$

Emergent Lieb-Robinson Lightcone

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley



Local Spin Systems and Lieb-Robinson Bounds


Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

Local Spin Systems and Lieb-Robinson Bounds

Speed Limits for
Quantum Control
of Local Spin
Systems


Jeffrey M. Epstein
and K. Birgitta
Whaley



Local Spin Systems and Lieb-Robinson Bounds

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley




$$c_t(X, Y) \leq 2l_L(4Jt)$$


Local Spin Systems and Lieb-Robinson Bounds

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley




$$c_t(X, Y) \leq 2l_L(4Jt)$$




Local Spin Systems and Lieb-Robinson Bounds

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley



$$c_t(X, Y) \leq 2l_L(4Jt)$$



$$V_{LR} \leq 6J$$

Outline

1. Motivation for bounds on optimal control
2. Brief review of Lieb-Robinson bounds
3. Bounds on state transfer and entanglement generation
4. Comparison to theory and numerics

Figures of Merit for State Transfer


Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley


Figures of Merit for State Transfer

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley




Figures of Merit for State Transfer



T : unitary accounting for passive dynamics + control


Figures of Merit for State Transfer



T : unitary accounting for passive dynamics + control

Goal: $\forall \rho_X \quad F(\rho_X, (T\rho T^\dagger)_Y) \approx 1$


Figures of Merit for State Transfer



T : unitary accounting for passive dynamics + control

$$\text{Goal: } \forall \rho_X \quad F(\rho_X, (T\rho T^\dagger)_Y) \approx 1$$

Figures of Merit for State Transfer




T : unitary accounting for passive dynamics + control

A : bounded operator on X accounting for state preparation

Goal: $\forall \rho_X \quad F(\rho_X, (T\rho T^\dagger)_Y) \approx 1$

Figures of Merit for State Transfer




T : unitary accounting for passive dynamics + control

A : bounded operator on X accounting for state preparation

Goal: $\forall \rho_X \quad F(\rho_X, (T\rho T^\dagger)_Y) \approx 1$

Figures of Merit for State Transfer



T : unitary accounting for passive dynamics + control

A : bounded operator on X accounting for state preparation

Goal: $\forall \rho_X \quad F(\rho_X, (T\rho T^\dagger)_Y) \approx 1$

Necessary condition: $\exists A \quad F((T\rho T^\dagger)_Y, (TA\rho A^\dagger T^\dagger)_Y) \approx 0$

Algebraic Bound on State Transfer

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

²JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Algebraic Bound on State Transfer

Let X and Y be disjoint subsystems of a system S in the initial state ρ .

²JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Algebraic Bound on State Transfer

Let X and Y be disjoint subsystems of a system S in the initial state ρ . For T a unitary on S , suppose for any A, B acting on X and Y , respectively,

²JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Algebraic Bound on State Transfer

Let X and Y be disjoint subsystems of a system S in the initial state ρ . For T a unitary on S , suppose for any A, B acting on X and Y , respectively,

$$\|[A_T, B]\| \leq c_T(X, Y) \|A\| \|B\|. \quad (1)$$

²JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Algebraic Bound on State Transfer

Let X and Y be disjoint subsystems of a system S in the initial state ρ . For T a unitary on S , suppose for any A, B acting on X and Y , respectively,

$$\|[A_T, B]\| \leq c_T(X, Y) \|A\| \|B\|. \quad (1)$$

Then the fidelity between the reduced states of subsystem Y given the overall states $T\rho T^\dagger$ and $TA\rho A^\dagger T^\dagger$ satisfies:

²JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Algebraic Bound on State Transfer

Let X and Y be disjoint subsystems of a system S in the initial state ρ . For T a unitary on S , suppose for any A, B acting on X and Y , respectively,

$$\|[A_T, B]\| \leq c_T(X, Y) \|A\| \|B\|. \quad (1)$$

Then the fidelity between the reduced states of subsystem Y given the overall states $T\rho T^\dagger$ and $TA\rho A^\dagger T^\dagger$ satisfies:

$$F \geq 1 - c_T(X, Y) \|A\|^2. \quad (2)$$


²JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Figure of Merit for Entanglement Generation

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley


Figure of Merit for Entanglement Generation



Speed Limits for
Quantum Control
of Local Spin
Systems


Jeffrey M. Epstein
and K. Birgitta
Whaley

Figure of Merit for Entanglement Generation



$$\max \{F(\rho, \psi) : \psi \text{ maximally entangled}\}$$


Figure of Merit for Entanglement Generation



$$\max \{F(\rho, \psi) : \psi \text{ maximally entangled}\}$$

Intuition: large entanglement \longrightarrow large correlations
 small entanglement \longleftarrow small correlations

Figure of Merit for Entanglement Generation



$$\max \{F(\rho, \psi) : \psi \text{ maximally entangled}\}$$

Intuition: large entanglement \longrightarrow large correlations
small entanglement \longleftarrow small correlations

connected correlator: $\langle AB \rangle_c = \langle AB \rangle - \langle A \rangle \langle B \rangle$

Algebraic Entanglement Bound

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

³JME, K. Birgitta Whaley 2016 arXiv:1612.04767 

Algebraic Entanglement Bound

Let ρ be a state of a bipartite system XY

³JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Algebraic Entanglement Bound

Let ρ be a state of a bipartite system XY such that for Hermitian operators A, B on X and Y , respectively, with $\|A\|, \|B\| \leq 1$

³JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Algebraic Entanglement Bound

Let ρ be a state of a bipartite system XY such that for Hermitian operators A, B on X and Y , respectively, with $\|A\|, \|B\| \leq 1$, the bound $|\langle AB \rangle_c| \leq f$ on the magnitude of the connected correlator $\langle AB \rangle - \langle A \rangle \langle B \rangle$ holds.

³JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Algebraic Entanglement Bound

Let ρ be a state of a bipartite system XY such that for Hermitian operators A, B on X and Y , respectively, with $\|A\|, \|B\| \leq 1$, the bound $|\langle AB \rangle_c| \leq f$ on the magnitude of the connected correlator $\langle AB \rangle - \langle A \rangle \langle B \rangle$ holds.

Then

$$F(\rho, \psi) \leq \sqrt{\frac{79}{81} + \frac{2f}{27} - \frac{f^2}{18}} \quad (3)$$

for any maximally entangled state Ψ .³

³JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Bound on Correlations

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

Bound on Correlations

Let a system S be initialized in the state ρ with the property that for any disjoint subsystems $X, Y \subset S$ and any A and B with $\|A\|, \|B\| \leq 1$ acting on X and Y , respectively,

$$|\langle AB_c \rangle| \leq f_0(X, Y).$$

⁴JME, K. Birgitta Whaley 2016 arXiv:1612.04767

Bound on Correlations

Let a system S be initialized in the state ρ with the property that for any disjoint subsystems $X, Y \subset S$ and any A and B with $\|A\|, \|B\| \leq 1$ acting on X and Y , respectively, $|\langle AB_c \rangle| \leq f_0(X, Y)$.

Then for some unitary T :

$$|\langle A_T B_T \rangle_c| \leq f_0(Z, \bar{Z}) + 2((c_T(X, \bar{Z}) + 1)(c_T(Z, Y) + 1)) - 1 \quad (4)$$

for $X \subseteq Z \subseteq S \setminus Y$.⁴

⁴JME, K. Birgitta Whaley 2016 arXiv:1612.04767


Bound on Correlations

Let a system S be initialized in the state ρ with the property that for any disjoint subsystems $X, Y \subset S$ and any A and B with $\|A\|, \|B\| \leq 1$ acting on X and Y , respectively, $|\langle AB_c \rangle| \leq f_0(X, Y)$.

Then for some unitary T :

$$|\langle A_T B_T \rangle_c| \leq f_0(Z, \bar{Z}) + 2((c_T(X, \bar{Z}) + 1)(c_T(Z, Y) + 1)) - 1 \quad (4)$$

for $X \subseteq Z \subseteq S \setminus Y$.⁴



⁴JME, K. Birgitta Whaley 2016 arXiv:1612.04767

1. Motivation for bounds on optimal control
2. Brief review of Lieb-Robinson bounds
3. Bounds on state transfer and entanglement generation
4. Comparison to theory and numerics

State Transfer v. Entanglement Generation

Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

⁵Calabrese, Cardy 2006 arXiv:0601225 ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ≡ ≡ ≡ ↺ 🔍 ↻

State Transfer v. Entanglement Generation

state transfer:

requires $c_T(X, Y) \gtrsim 0$

State Transfer v. Entanglement Generation

state transfer:

$$\text{requires } c_T(X, Y) \gtrsim 0$$

entanglement generation:

$$\text{requires } (c_T(X, \bar{Z}) + 1)(c_T(Z, Y) + 1) - 1 \gtrsim 0$$


State Transfer v. Entanglement Generation

state transfer:

$$\text{requires } c_T(X, Y) \gtrsim 0$$

entanglement generation:

$$\text{requires } (c_T(X, \bar{Z}) + 1)(c_T(Z, Y) + 1) - 1 \gtrsim 0$$




⁵Calabrese, Cardy 2006 arXiv:0601225

State Transfer v. Entanglement Generation

state transfer:

requires $c_T(X, Y) \geq 0$

entanglement generation:




“entanglement carriers”⁵

Is the Gap Real?


Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

Is the Gap Real?




Is the Gap Real?



$2J$: group velocity of 1-excitation Heisenberg chain


Is the Gap Real?



$2J$: group velocity of 1-excitation Heisenberg chain

Lieb-Robinson velocities can be significantly higher than group velocities

Is the Gap Real?



$2J$: group velocity of 1-excitation Heisenberg chain

Lieb-Robinson velocities can be significantly higher than group velocities

theoretical bound applies to arbitrary 2-local models

Thank you!



Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley



Quantum Speed Limits without Spatial Structure


Speed Limits for
Quantum Control
of Local Spin
Systems

Jeffrey M. Epstein
and K. Birgitta
Whaley

Orthogonalization/distinguishability time:


$$\theta(\psi(t), \psi) \leq \frac{1}{2} \Delta_{\max} t \longrightarrow t_{\perp} = \pi / \Delta_{\max}$$

Information Transfer:



Quantum

$$t_*^Q = \pi / \Delta_{\max}$$



Classical

$$t_*^C = \pi / 2 \Delta_{\max}$$

$$\text{Entanglement generation: } t_*^E = \frac{2}{\Delta_{\max}} \arccos d^{-1/2}$$

Proof of State Transfer Bound

X-localization of an operator: $[A]_X = \int_{U(\bar{X})} UAU^\dagger d\mu(U)$

small lemma: $\|U\rho U^\dagger - V\rho V^\dagger\|_1 \leq 2\|U - V\|$

$$\begin{aligned}
 & \left\| \text{tr}_{S \setminus Y} \left(A_T \rho_T A_T^\dagger - \rho_T \right) \right\|_1 \\
 & \leq \left\| \text{tr}_{S \setminus Y} \left(A_T \rho_T A_T^\dagger - [A_T]_{S \setminus Y} \rho_T [A_T^\dagger]_{S \setminus Y} \right) \right\|_1 \\
 & \quad + \left\| \text{tr}_{S \setminus Y} \left([A_T]_{S \setminus Y} \rho_T [A_T^\dagger]_{S \setminus Y} - \rho_T \right) \right\|_1 \\
 & \leq \left\| A_T \rho_T A_T^\dagger - [A_T]_{S \setminus Y} \rho_T [A_T^\dagger]_{S \setminus Y} \right\|_1 \leq 2 \left\| A_T - [A_T]_{S \setminus Y} \right\| \\
 & = 2 \left\| A_T - \int U A_T U^\dagger d\mu(U) \right\| = 2 \int \| [A_T, U] \| d\mu(U) \\
 & \leq 2c_T(X, Y) \|A\|
 \end{aligned}$$

Similar to analysis in ⁶

⁶Bravyi, Hastings, Verstraete 2006 arXiv:0603121

Proof of Entanglement Bound

Let $|\Psi\rangle$ be maximally entangled, $\Delta = \rho - |\Psi\rangle\langle\Psi|$.

$$\begin{aligned}\langle AB \rangle_c &= \text{Tr}(\rho AB) - \text{Tr}(\rho A) \text{Tr}(\rho B) \\ &= \langle \Psi | AB | \Psi \rangle - \langle \Psi | A | \Psi \rangle \langle \Psi | B | \Psi \rangle \\ &\quad + \text{Tr}(\Delta AB) - \text{Tr}(\Delta A) \text{Tr}(\Delta B).\end{aligned}\tag{5}$$

Rearranging and taking the modulus:

$$\begin{aligned}|\langle AB \rangle_c - \langle \Psi | AB | \Psi \rangle + \langle \Psi | A | \Psi \rangle \langle \Psi | B | \Psi \rangle| \\ \leq |\text{Tr}(\Delta AB)| + |\text{Tr}(\Delta A)| |\text{Tr}(\Delta B)| \\ \leq \|\Delta\|_1 \|A\| \|B\| + \|\Delta\|_1^2 \|A\| \|B\| \\ \leq \|\Delta\|_1 + \|\Delta\|_1^2 \leq 3 \|\Delta\|_1.\end{aligned}\tag{6}$$

Reverse triangle inequality:

$$\begin{aligned}3 \|\Delta\|_1 &\geq ||\langle \Psi | AB | \Psi \rangle - \langle \Psi | A | \Psi \rangle \langle \Psi | B | \Psi \rangle| - |\langle AB \rangle_c| \\ &= \left| |\langle AB \rangle_{c,\psi}| - |\langle AB \rangle_c| \right|\end{aligned}\tag{7}$$

Proof of Entanglement Bound

For any maximally entangled state, there are A' and B' such that $\langle A'B' \rangle_c \geq 2/3$, so that

$$3 \|\Delta\|_1 \geq \left| \frac{2}{3} - |\langle A'B' \rangle_c| \right|. \quad (8)$$

Given $|\langle A'B' \rangle_c| \leq f$ (assuming w.l.o.g. $f < 2/3$)

$$3 \|\Delta\|_1 \geq \frac{2}{3} - f \quad (9)$$

$$F \leq \sqrt{1 - \frac{1}{2} \|\Delta\|_1^2} \leq \sqrt{\frac{79}{81} + \frac{2f}{27} - \frac{f^2}{18}} \quad (10)$$