

# **Dynamic Modeling of the Human Coagulation Cascade using Reduced Order Effective Kinetic Models**

Adithya Sagar and Jeffrey D. Varner\*

School of Chemical and Biomolecular Engineering  
Cornell University, Ithaca NY 14853

**Running Title:** Reduced order modeling of blood coagulation

**To be submitted:** *Processes*

\*Corresponding author:

Jeffrey D. Varner,

Associate Professor, School of Chemical and Biomolecular Engineering,  
244 Olin Hall, Cornell University, Ithaca NY, 14853

Email: [jdv27@cornell.edu](mailto:jdv27@cornell.edu)

Phone: (607) 255 - 4258

Fax: (607) 255 - 9166

## Abstract

In this study, we present a novel modeling approach which combines ordinary differential equation (ODE) modeling with logical rules to simulate an archetype biochemical network, the human coagulation cascade. The model consisted of five differential equations augmented with several logical rules describing regulatory connections between model components, and unmodeled interactions in the network. This formulation was more than an order of magnitude smaller than current coagulation models, because many of the mechanistic details of coagulation were encoded as logical rules. We estimated an ensemble of likely model parameters ( $N = 20$ ) from *in vitro* extrinsic coagulation data sets, with and without inhibitors, by minimizing the residual between model simulations and experimental measurements using particle swarm optimization (PSO). Each parameter set in our ensemble corresponded to a unique particle in the PSO. We then validated the model ensemble using thrombin data sets that were not used during training. The ensemble predicted thrombin trajectories for conditions not used for model training, including thrombin generation for normal and hemophilic coagulation in the presence of platelets (a significant unmodeled component). We then used flux analysis to understand how the network operated in a variety of conditions, and global sensitivity analysis to identify which parameters controlled the performance of the network. Taken together, the hybrid approach produced a surprisingly predictive model given its small size, suggesting the proposed framework could also be used to dynamically model other biochemical networks, including intracellular metabolic networks, gene expression programs or potentially even cell free metabolic systems.

**Keywords:** Blood coagulation, Mathematical modeling, Systems biology

## 1 Introduction

2 Developing mathematical models of biochemical networks is a significant facet of systems  
3 biology. Modeling approaches differ in their degree of detail, where the choice of approach  
4 is often determined by prior knowledge, or model requirements [1]. Ordinary differential  
5 equation (ODE) models are common tools for modeling biochemical systems because of  
6 their ability to capture dynamics and encode mechanism. However, ODE models typi-  
7 cally come with difficult (or sometimes impossible) parameter identification problems. For  
8 example, Gadkar *et al.*, showed that even with near-perfect information, it was often im-  
9 possible to identify all the parameters in typical signal transduction models [2]. However,  
10 it is not clear whether we actually need precise estimates for all model parameters. Bailey  
11 suggested more than a decade ago, that achieving qualitative or even quantitative un-  
12 derstanding of biological systems should not require complete structural and parametric  
13 knowledge [3]. Since Bailey's complex biology with no parameters hypothesis, Sethna  
14 showed that model performance is typically sensitive to only a few parameters, a charac-  
15 teristic seemingly universal to multi-parameter models referred to as *sloppiness* [4]. Thus,  
16 reasonable predictions may be possible, despite parametric uncertainty, if a few critical  
17 parameters are well-defined. For example, Tasseff et al., showed in a model of Retinoic  
18 acid (RA) induced differentiation of HL-60 cells, that correct predictions were possible  
19 even when 75% of the parameters were known only to an order of magnitude [5]. Per-  
20 haps more importantly, ODE models require significant mechanistic information, thereby  
21 limiting their utility in poorly understood systems, or conversely explode in size when con-  
22 sidering multiple pathways or subsystems. Toward this challenge, logical modeling is an  
23 emerging paradigm that encodes causal relationships between model components using  
24 quasi-mechanistic non-linear transfer functions [6]. Logical models are highly flexible, and  
25 despite their simplicity, they have captured rich behaviors in a variety of systems impor-  
26 tant to human health [7–9]. However, modeling complex dynamics with logical models is

challenging. Thus, there is an unmet need for a third approach which combines ODEs and logical models, where ODEs could encode mechanistic information, while missing or incomplete mechanistic knowledge can be approximated using a logical approach.

In this study, we developed a hybrid approach which combined ODE modeling with logical rules to model a well studied biochemical network, the human coagulation system. Coagulation is an archetype proteolytic cascade involving both positive and negative feedback [10–12]. Coagulation is mediated by a family proteases in the circulation, called factors and a key group of blood cells, called platelets. The central process in coagulation is the conversion of prothrombin (fII), an inactive coagulation factor, to the master protease thrombin (FIIa). Thrombin generation involves three phases, initiation, amplification and termination [13, 14]. Initiation requires a trigger event, for example vessel injury, which leads to the activation of factor VII (FVIIa). Two converging pathways, the extrinsic and intrinsic cascades, then process and amplify this initial coagulation signal. The extrinsic cascade is generally believed to be the main mechanism of thrombinogenesis in the blood [15–17]. Initially, thrombin is produced upon cleavage of prothrombin by fluid phase activated factor X (FXa), which itself has been activated by TF/FVIIa [10]. Picomolar amounts of thrombin then activate the cofactors factors V and VIII (fV and fVIII) and platelets, leading to the formation of the tenase and prothrombinase complexes on activated platelets. These complexes amplify the early coagulation signal by further activating FXa, and directly converting prothrombin to thrombin. There are several control points in the cascade that inhibit thrombin formation, and eventually terminate thrombin generation. Tissue Factor Pathway Inhibitor (TFPI) inhibits FXa formation catalyzed by TF/FVIIa, while antithrombin III (ATIII) neutralizes several of the proteases generated during coagulation, including thrombin. Thrombin itself also inadvertently plays a role in its own inhibition; thrombin, through interaction with thrombomodulin, protein C and endothelial cell protein C receptor (EPCR), converts protein C to activated protein C (APC) which

attenuates the coagulation response by proteolytic cleavage of fV/FVa and fVIII/FVIIIa. Termination occurs after either prothrombin is consumed, or thrombin formation is neutralized by inhibitors such as APC or ATIII.

Previous coagulation models have typically been formulated as systems of nonlinear ordinary differential equations, using mass action or more complex kinetics, to describe the rates of biochemical conversions [18–22]. Mechanistic ODE coagulation models from our laboratory [23, 24] were built upon the earlier studies of Jones and Mann [25], Hockin *et al.* [26], and later Butenas *et al.*, [27] who developed and then subsequently refined highly mechanistic coagulation models. Other laboratories have also expanded upon Hockin *et al.*, for example by exploring the intrinsic pathway, the role of stochastic fluctuations in coagulation [28], and the dynamics of thrombin mediated clot formation [29]. Other aspects of coagulation have also been modeled, such as platelet biochemistry [30], multi-scale models of clot formation [31, 32], and transport inside clots [33]. However, these previous studies were largely based upon extensive mechanistic knowledge. This is possible because blood, while enormously complex, can be systematically interrogated. Other systems, such as intracellular signaling networks, are much more difficult to experimentally interrogate. Towards this unmet need, we formulated a hybrid modeling approach which combines ODEs and logical rules to model biochemical processes for which a complete mechanistic understanding is missing. We tested this approach by modeling the human coagulation cascade. The hybrid model consisted of only five differential equations augmented with several logical rules. Thus, the model was more than an order of magnitude smaller than comparable purely ODE models in the literature. We estimated the model parameters from *in vitro* extrinsic coagulation data sets, in the presence of ATIII, with and without the protein C pathway. We then compared the model predictions with thrombin data sets, for both normal and hemophilic coagulation, that were not used for model training. Once validated, we performed flux and sensitivity analysis on the

79 model to estimate which parameters were critical to model performance in several condi-  
80 tions. The reduced order hybrid approach produced a surprisingly predictive coagulation  
81 model, suggesting this framework could potentially be used to model other biochemical  
82 networks important to human health.

## Results

**Formulation of reduced order coagulation models.** We developed a reduced order extrinsic coagulation model to test our hybrid modeling approach (Fig. 1). The core of our model was based upon the earlier work of Ismagilov and coworkers [34–37], where we added initiation, factor dependence, and specific inhibition terms to the earlier simplified model. A trigger event initiates thrombin formation (FIIa) from prothrombin (fII) through a lumped initiation step. This step loosely represents the initial activation of thrombin by activated FXa. Once activated, thrombin catalyzes its own formation (amplification step), and inhibition via the conversion of protein C to activated protein C (APC). Antithrombin III (ATIII) inhibits amplification, while APC and tissue factor pathway inhibitor (TFPI) potentially inhibit both initiation and amplification. All initiation and inhibition processes, as well as the dependence of amplification upon other coagulation factors, was approximated using our rule-based approach (Fig. 2). Individual regulatory contributions to the activity of pathway enzymes were integrated into control coefficients ( $v$ 's) using an integration rule (min/max). These control coefficients then modified the rates of model processes at each time step. Hill-like transfer functions  $0 \leq f(\mathcal{Z}) \leq 1$  quantified the contribution of components upon a target process. Components were either individual inhibitor or activator levels or some function of levels, e.g., the product of factor levels. In this study,  $\mathcal{Z}$  corresponded to the abundance of individual inhibitors or activators, with the exception of the dependence of amplification upon specific coagulation factors (modeled as the product of factors). When a process was potentially sensitive to multiple inputs, logical integration rules were used to select which transfer functions influenced the process at any given time. In our proof of concept model, we used a winner takes all strategy; the maximum or minimum transfer function was selected at any given time step. However, other integration rules are certainly possible. Taken together, while the reduced order coagulation model encodes significant biological complexity, it is highly compact (consisting of only

five differential equations). Thus, it will serve as an excellent proof of principle example to study the reduction of a highly complex human subsystem.

**Identification of model parameters using particle swarm optimization.** A critical challenge for any dynamic model is the estimation of kinetic parameters. We estimated kinetic and control parameters simultaneously from eight *in vitro* time-series coagulation data sets with and without the protein C pathway. The residual between model simulations and experimental measurements was minimized using particle swarm optimization (PSO). A population of particles ( $N = 20$ ) was initialized with randomized kinetic and control parameters and allowed to search for parameter vectors that minimized the residual. However, not all parameters were varied simultaneously. We partitioned the parameter estimation problem into two subproblems based upon the biological organization of the training data; (i) estimation of parameters associated with thrombin formation in the absence of the protein C pathway and (ii) estimation of parameters associated with the protein C pathway. Only those parameters associated with each subproblem were varied during the optimization procedure for that subproblem, e.g., thrombin parameters were *not* varied during the protein C subproblem. The PSO procedure was run for 20 generations for each subproblem, where each generation was 1200 iterations. The best particle from each generation was used to generate the particle population for the next generation. We rotated the subproblems, starting with subproblem 1 in the first generation.

The reduced order coagulation model captured the role of initial prothrombin abundance, and the decay of the thrombin signal following from ATIII activity (Fig. 3). However, we systematically under-predicted the thrombin peak and the strength of ATIII inhibition in this training data set. On the other hand, with fixed thrombin parameters, we captured peak thrombin values and the decay of the thrombin signal (at least for the 150% fII case) in the presence of both ATIII and the protein C pathway (Fig. 4). Lastly, we were unable to capture global differences in initiation time *across* separate data sets with a single en-



semble of model parameters. These differences likely resulted from normal experimental variability. For example, different thrombin generation experiments within the training data (at the same physiological factor levels) had significantly different initiation times (data not shown). However, the inability to globally capture initiation time also highlighted a potential shortcoming of the initiation module within the model. To capture the variability in initiation time *across* training data sets, we included a constant time-delay parameter ( $T_D$ ) for each data group. The delay parameter was constant within a data set, but allowed to vary *across* training data sets. Introduction of the delay parameter allowed the model to simulate multiple training data sets using a single ensemble of model parameters. Taken together, the model identification results suggested that our hybrid approach could reproduce a panel of thrombin generation data sets in the neighborhood of physiological factor and inhibitor concentrations. However, it was unclear whether the reduced order model could predict new data, without updating the model parameters.

**Validation of the reduced order coagulation model.** We tested the predictive power of the reduced order coagulation model with validation data sets not used during model training. Two validation data sets were used, thrombin generation for various prothrombin and ATIII concentrations with the protein C pathway, and thrombin generation in normal versus hemophilic plasma in the presence of the protein C pathway. Lastly, we compared the qualitative output of the model to rFVIIa addition in the presence of hemophilia. The hemophilia case was an especially difficult test as it was taken from a different study which used a plasma-based *in vitro* assay involving platelets instead of phospholipid vesicles (PCPS). All kinetic and control parameters were fixed for the validation simulations. The only globally adjustable parameter  $T_D$ , was fixed within each validation data set but allowed to vary between data sets. The reduced order model predicted the thrombin generation profile for ratios of prothrombin and ATIII in the absence of the protein C pathway (Fig. 5). Simulations near the physiological range ( $[II,ATIII] = (100\%, 100\%)$ ) or

(125%,75%) tracked the measured thrombin values (Fig. 5B and C). On the other hand, predictions for factor levels outside of the physiological range ( $fII,ATIII$ ) = (50%, 150%) or (150%, 50%), while qualitatively consistent with measured thrombin values, did show significant deviation from the measurements (Fig. 5A and D). Likewise, simulations of thrombin generation in normal versus hemophilia (missing both  $fVIII$  and  $fIX$ ) were consistent with measured thrombin values (Fig. 6). We modeled the dependence of thrombin amplification on factor levels using a product rule ( $\mathcal{Z} = fV \times fX \times fVIII \times fIX$ ), which was then integrated using a `min` integration rule into the control variable governing amplification. Thus, in the absence of  $fVIII$  or  $fIX$ , the amplification control variable evaluated to zero, and the only thrombin produced was from initiation (Fig. 6B). However, the decay of the thrombin signal was underpredicted in the normal case (Fig. 6A), while the activated thrombin level was overpredicted in hemophilia simulations, although thrombin generation was far less than normal (Fig. 6B). Taken together, the reduced order model performed well in the physiological range of factors, even with unmodeled components such as platelet activation in the hemophilia data set.

The model ensemble predicted a direct correlation between thrombin generation and  $rFVIIa$  addition in hemophilia (Fig. 7). In the current model, we cannot distinguish between different initiation sources, e.g.,  $TF/FVIIa$  versus  $rFVIIa$ , as we have only a single lumped initiation source (trigger). Thus, we simulated the addition of  $rFVIIa$  in hemophilia by removing  $fVIII$  and  $fIX$  from the model, and modulating the initial level of trigger. Simulations with a baseline level of trigger were consistent with the previous hemophilia simulations, where the only thrombin produced was from initiation (Fig. 7A,  $1 \times$  trigger). However, as we increased the trigger strength, the thrombin profile began to approximate normal coagulation, showing a pronounced peak albeit with a slower peak time (Fig. 7B,  $t^{**} > t^*$ ). Further increases in trigger strength resulted in decreased thrombin peak time and increased maximum thrombin values (Fig. 7A,  $50 \times$  trigger). Thus, for large trigger

values ( $200\times$ trigger), the hemophilic thrombin profile approximated normal coagulation, where peak thrombin was achieved shortly after administration and 95% of the thrombin was gone by 20 min after initiation. We performed flux analysis to understand how the reduced order coagulation model balanced initiation, amplification and termination of thrombin generation for normal and hemophilic coagulation. Analysis of the reaction flux through the reduced order network for thrombin generation in normal, hemophilia and rFVIIa-treated hemophilia identified three distinct operational modes (Fig. 8). We calculated the flux through four lumped reactions, initiation, amplification, thrombin-induced APC generation and total thrombin inhibition (including both APC and ATIII action). Directly after the addition of a trigger (e.g., TF/FVIIa or rFVIIa), the lumped initiation flux was the largest for all three cases. However, within a few minutes enough thrombin was generated by the initiation mechanism to induce the amplification stage. During amplification, thrombin catalyzes its own formation and inhibition by generating activated protein C (APC), a potent inhibitor of the coagulation cascade. For normal coagulation, amplification and thrombin inhibition were the dominate reactions by 6 min after initiation (Fig. 8, left). After 10 min, the dominate reaction had shifted to thrombin inhibition (both ATIII and APC action). In hemophilia (missing both fVIII and fIX), the amplification reaction did not occur, and thrombin was produced only by initiation (Fig. 8, center). Initiation was quickly inhibited by APC, and the thrombin level stabilized (eventually decaying at longer times because of ATIII activity). Lastly, when  $50\times$ trigger was used to induce thrombin formation in hemophilia (absence of fVIII/fIX), initiation mechanisms dominated for up to 6 min following initiation (Fig. 8, right). Similar to hemophilia alone, no amplification occurred in the  $50\times$ trigger+hemophilia case, and the rate of thrombin generation was extinguished by the combined action of ATIII and APC. Taken together, the hybrid modeling approach captured the transition between the modes of thrombin generation, as well as the role that inhibitors play in attenuating the thrombin generation rate. Thus, the transfer

function approach encoded the inhibitory logic of this cascade in the absence of specific mechanism.

**Global sensitivity analysis of the reduced order coagulation model.** We conducted a global sensitivity analysis to estimate which parameters controlled the performance of the reduced order model. We calculated the sensitivity of the time to maximum thrombin (peak time) and the thrombin exposure (area under the thrombin curve) for different levels of prothrombin, and protein C (Fig. 9). Globally, 41% of the parameters shifted in importance between the (fII,PC) = (50%, 0%) and (150%,100%) cases for the peak thrombin time (Fig. 9A). The majority of these shifts involved the interaction between increased prothrombin and the protein C pathway, while only 5% were directly associated with increased prothrombin alone. The rate constant for thrombin amplification was the most important parameter controlling the peak thrombin time. While this parameter was differentially important for different prothrombin levels, and in the presence or absence of the activated protein C pathway, it was consistently the most sensitive parameter in the model. The saturation constant governing thrombin amplification was the second most important parameter, followed by the initiation control gain parameter. Other important parameters influencing the thrombin peak time included the control gain for activated protein C formation, and the rate constant controlling ATIII inhibition of thrombin activity. On the other hand, only 27% of the model parameters were differentially sensitive between the (fII,PC) = (50%, 0%) and (150%,100%) cases for thrombin exposure (Fig. 9B). Of these parameters, all of the shifts were associated with the interplay between thrombin formation and the protein C pathway. The rate constant controlling ATIII inhibition was the most important parameter controlling the thrombin exposure. While this parameter was less important in the presence of protein C for 150% prothrombin levels, it was significantly above all other parameters. Similar to the peak time, for 150% prothrombin, the control gain for activated protein C formation was differentially important along with the

239 rate constant controlling amplification. However, the amplification parameter was much  
240 less important for thrombin exposure versus peak time.

## Discussion

In this study, we developed a reduced order model of the human coagulation cascade. We modeled coagulation because it is well studied, has a complex architecture, and has an abundance of experimental data available for model identification and validation. However, coagulation was just a proof of concept test of our approach. The proposed hybrid framework could also be used to dynamically model other biochemical networks, including intracellular metabolic networks, gene expression programs or potentially even cell free metabolic systems. The model consisted of five differential equations augmented with several logical rules describing regulatory connections between model components and unmodeled interactions in the network. We estimated model parameters from *in vitro* extrinsic coagulation data sets, in the presence of ATIII, with and without the protein C pathway. To estimate parameters, the residual between model simulations and experimental measurements was minimized using particle swarm optimization (PSO). However, not all of the model parameters were uniquely identifiable, given the training data. Instead, we estimated an ensemble of likely parameter sets ( $N = 20$ ) from eight *in vitro* time-series coagulation data sets with and without the protein C pathway. Ensemble approaches have been used previously for other signal transduction models [38–42], and for metabolic models [43] to estimate the impact of poorly constrained parameter values or poorly understood network structure on simulation performance. Thus, ensemble approaches are common in the dynamic modeling community. However, a unique feature of the current study is the direct connection between our particle swarm approach, and the parameter ensemble; each particle in our swarm uniquely corresponded to a parameter set in our ensemble. Thus, by constraining particles to operate in different parameter regions, giving each particle a different parameter combination to explore, or perhaps even supplying a different model formulation to each particle we can effectively traverse through complex parameter and model spaces. We validated the ensemble using thrombin data

sets taken from multiple laboratories for a variety of experimental conditions not used during training. The ensemble predicted thrombin trajectories for conditions not used for model training, including thrombin generation for normal and hemophilic coagulation in the presence of platelets (a significant unmodeled component). We then used flux analysis to understand how the network operated in a variety of conditions, and global sensitivity analysis to identify which parameters controlled the performance of the network. Flux analysis showed the logical rules formulation encoded the transitions between initiation, amplification and termination of thrombin generation. Sensitivity analysis suggested that the amplification rate constant was more important to the time to peak thrombin, while the ATIII inhibition constant controlled thrombin exposure. Taken together, the proposed hybrid framework produced a surprisingly predictive model, suggesting this approach could be used to effectively model other biochemical networks important to human health.

Malfunctions in coagulation can have potentially fatal consequences. Aggressive clotting involved with Coronary Artery Diseases (CADs), collectively accounts for 38% of all deaths in North America [44]. Coagulation management during surgery can also be challenging, particularly with the increase in clinical use of antithrombotic drugs [45]. Insufficient coagulation due to genetic disorders such as hemophilia can also result in recurrent bleeding. The coagulation factors VIII (fVIII) and IX (fIX) are deficient in Hemophilia A and B, respectively [46–48]. People with mild hemophilia have 5-40% of the normal clotting factor levels while severe hemophiliacs have  $<1\%$  [48]. Hemophilia can be controlled with regular infusions of the deficient clotting factors. However, clotting factor replacement sometimes leads to the formation of fVIII and fIX inhibitors *in vivo* [49]. Alternatively, recombinant factor VIIa (rFVIIa) has been used to treat bleeding disorders [50, 51] including hemophilia with and without factor VIII/IX inhibitors [52]. However, rFVIIa requires frequent administration (every 2-3 hr), and many questions remain about its mechanism of action, its effective dosage [49], and its overall utility for the treatment trauma-associated hem-

orrhage [53]. In this study, we did not model rFVIIa-induced coagulation directly. Rather, we modeled a general trigger which initiated the extrinsic coagulation cascade. Since we identified the model using TF/FVIIa, inherent to our rFVIIa simulations (and the rate constant governing initiation) was the presence of TF. However, even with this complication, the model generated potentially useful insight into the rFVIIa mechanism of action, and its possible shortcomings especially for the treatment of hemophilia. The addition of rFVIIa directly activated thrombin through the initiation pathway. However, no amplification of the thrombin signal occurred without fVIII or fIX. Thus, the peak thrombin signal was lower than normal coagulation, the peak thrombin time was longer, and thrombin generation was eventually inhibited by the combined action of ATIII and the protein C pathway. However, as the dose of rFVIIa increased, the peak thrombin time decreased (eventually saturating around  $200 \times$  nominal trigger), and the peak thrombin value increased such that the thrombin profile resembled normal coagulation. Butenas *et al.* performed an extensive *in vitro* study of rFVIIa-induced thrombin generation under normal and hemophilic conditions [54]. They found qualitatively similar trends, namely rFVIIa restored normal coagulation (even in the absence of TF) for large enough rFVIIa doses, although rFVIIa-induced coagulation in hemophilia (even for large rFVIIa doses) lagged the normal profile. These results suggest that rFVIIa administration alone might not be able to initiate normal coagulation in recurrent bleeding, unless the dosage is well above a critical threshold. However, defining this threshold, which is likely patient specific, is difficult as there is tremendous patient to patient variability even with a normal coagulation phenotype [55].

The performance of the proof of principle coagulation model was impressive given its limited size, however, there are several issues that could be further explored. First, the prediction of initiation time should be investigated. We were able to estimate initiation time within a data set, but unable to predict initiation time *across* independent data sets. This suggested that we should update the initiation module to distinguish between different



triggers, e.g., TF/FVIIa versus rFVIIa alone, and to include key biological milestones such as FXa activation (a prerequisite to thrombin formation). Next, there are several additional biological modules that could be added to the core model presented here. First, we could include thrombin-induced platelet activation and the role of activated platelets in amplification. We captured thrombin generation data in the presence of platelets, however, the initial shape of the activation curve and the time-scale of activation was not always consistent with the data. Platelets are activated by thrombin through the cleavage of the extracellular domain of protease-activated receptors (PARs) on the platelet surface. Once activated, platelets play an important role in amplification, and are key mediators of the positive feedback driving amplification. Thus, this biology is a potentially important component of an expanded model. We should also add the intrinsic pathway to the model. The intrinsic pathway is triggered by contact activation of the plasma protease factor XI (fXI) by negatively charged surfaces and by thrombin and upstream factors such as activated plasma protease factor XII (FXIIa) [56, 57]. Activated platelets may also release polyphosphate which directly activates fXII [58]. Arguably a minor player in acute bleeding, contact activation could also be important in other wound healing contexts. Finally, to make the model more clinically relevant, we should include the biochemical processes responsible for clot formation and clot dissolution (fibrinolysis). Clot formation is driven by thrombin activity, while fibrinolysis is driven by plasmin activity, a key enzyme that cleaves fibrin (one of the main materials in a clot). Similar to coagulation, fibrinolysis is managed by several activating and inhibitory factors which control the balance between clot formation and dissolution. Tissue plasminogen activator (t-PA) and urokinase activate plasmin, along with contact pathway factors such as fXIa. On the other hand, thrombin activatable fibrinolysis inhibitor (TAFI) inhibits the degradation of fibrin by plasmin. Also, similar to coagulation, there is considerable fibrinolysis and contact pathway data sets that can be used to train the model. Lastly, the choice of max/min integration rules or

the particular form of the transfer functions could be generalized to include other rule types and functions. Theoretically, an integration rule is a function whose domain is a set of transfer function inputs, and whose range is  $v \in [0, 1]$ . Thus, integration rules other than max/min could be used, such as the mean or the product, assuming the range of the transfer functions is always  $f \in [0, 1]$ . Alternative integration rules such as the mean might have different properties which could influence model identification or performance. For example, a mean integration rule would be differentiable, which allows derivative-based optimization approaches to be used. The particular form of the transfer function could also be explored. We choose a Hill-like function because of its prominence in the systems and synthetic biology community. However, the only mathematical requirement for a transfer function is that it map a non-negative continuous or categorical variable into the range  $f \in [0, 1]$ . Thus, many types of transfer functions are possible.

## Materials and Methods

**Formulation and solution of the model equations.** We used ordinary differential equations (ODEs) to model the time evolution of proteins ( $x_i$ ) in our reduced order coagulation model:

$$\frac{dx_i}{dt} = \sum_{j=1}^{\mathcal{R}} \sigma_{ij} r_j(\mathbf{x}, \epsilon, \mathbf{k}) \quad i = 1, 2, \dots, \mathcal{M} \quad (1)$$

where  $\mathcal{R}$  denotes the number of reactions,  $\mathcal{M}$  denotes the number of protein species in the model. The quantity  $r_j(\mathbf{x}, \epsilon, \mathbf{k})$  denotes the rate of reaction  $j$ . Typically, reaction  $j$  is a non-linear function of biochemical species abundance, as well as unknown kinetic parameters  $\mathbf{k}$  ( $\mathcal{K} \times 1$ ). The quantity  $\sigma_{ij}$  denotes the stoichiometric coefficient for species  $i$  in reaction  $j$ . If  $\sigma_{ij} > 0$ , species  $i$  is produced by reaction  $j$ . Conversely, if  $\sigma_{ij} < 0$ , species  $i$  is consumed by reaction  $j$ , while  $\sigma_{ij} = 0$  indicates species  $i$  is not connected with reaction  $j$ . The system material balances were subject to the initial conditions  $\mathbf{x}(t_o) = \mathbf{x}_o$ , which were specified by the experimental setup.

Each reaction rate was written as the product of two terms, a kinetic term ( $\bar{r}_j$ ) and a regulatory term ( $v_j$ ):

$$r_j(\mathbf{x}, \epsilon, \mathbf{k}) = \bar{r}_j v_j \quad (2)$$

We used multiple saturation kinetics to model the reaction term  $\bar{r}_j$ :

$$\bar{r}_j = k_j^{max} \epsilon_i \left( \prod_{s \in m_j^-} \frac{x_s}{K_{js} + x_s} \right) \quad (3)$$

where  $k_j^{max}$  denotes the maximum rate for reaction  $j$ ,  $\epsilon_i$  denotes the scaled enzyme activity which catalyzes reaction  $j$ , and  $K_{js}$  denotes the saturation constant for species  $s$  in reaction  $j$ . The product in Eqn. (3) was carried out over the set of *reactants* for reaction  $j$  (denoted as  $m_j^-$ ).

376 The control term  $v_j$  depended upon the combination of factors which influenced the  
 377 activity of enzyme  $i$ . For each enzyme, we used a rule-based approach to select from  
 378 competing control factors (Fig. 2). If an enzyme was activated by  $m$  metabolites, we  
 379 modeled this activation as:

$$v_j = \max(f_{1j}(\mathcal{Z}), \dots, f_{mj}(\mathcal{Z})) \quad (4)$$

380 where  $0 \leq f_{ij}(\mathcal{Z}) \leq 1$  was a regulatory transfer function that calculated the influence of  
 381 metabolite  $i$  on the activity of enzyme  $j$ . Conversely, if enzyme activity was inhibited by  $m$   
 382 metabolites, we modeled this inhibition as:

$$v_j = 1 - \max(f_{1j}(\mathcal{Z}), \dots, f_{mj}(\mathcal{Z})) \quad (5)$$

383 Lastly, if an enzyme had both  $m$  activating and  $n$  inhibitory factors, we modeled the regu-  
 384 latory term as:

$$v_j = \min(u_j, d_j) \quad (6)$$

385 where:

$$u_j = \max_{j^+}(f_{1j}(\mathcal{Z}), \dots, f_{mj}(\mathcal{Z})) \quad (7)$$

$$d_j = 1 - \max_{j^-}(f_{1j}(\mathcal{Z}), \dots, f_{nj}(\mathcal{Z})) \quad (8)$$

386 The quantities  $j^+$  and  $j^-$  denoted the sets of activating and inhibitory factors for enzyme  $j$ .  
 387 If a process has no modifying factors, we set  $v_j = 1$ . There are many possible functional  
 388 forms for  $0 \leq f_{ij}(\mathcal{Z}) \leq 1$ . However, in this study, each individual transfer function took the  
 389 form:

$$f_i(\mathbf{x}) = \frac{\kappa_{ij}^\eta \mathcal{Z}_j^\eta}{1 + \kappa_{ij}^\eta \mathcal{Z}_j^\eta} \quad (9)$$

where  $\mathcal{Z}_j$  denotes the abundance of the  $j$  factor (e.g., metabolite abundance), and  $\kappa_{ij}$  and  $\eta$  are control parameters.  $\kappa_{ij}$  was the species gain parameter, while  $\eta$  was a cooperativity parameter (similar to a Hill coefficient). Applying the general framework to the reduced coagulation network resulted in five ordinary differential equations:

$$\frac{dx_1}{dt} = -(r_{init}v_{init} + r_{amp}v_{amp}) \quad (10)$$

$$\frac{dx_2}{dt} = r_{amp}v_{amp} + r_{init}v_{init} - r_{inh,ATIII}v_{inh,ATIII} \quad (11)$$

$$\frac{dx_3}{dt} = -r_{apc}v_{apc} \quad (12)$$

$$\frac{dx_4}{dt} = r_{apc}v_{apc} \quad (13)$$

$$\frac{dx_5}{dt} = -r_{inh,ATIII}v_{inh,ATIII} \quad (14)$$

where  $\mathbf{x} = (fII, FIIa, PC, APC, ATIII)^T$ . The terms  $r_*v_*$  in the balance equations denote corrected kinetic expressions for initiation, amplification and inhibition processes. The rate of initiation  $\bar{r}_{init}$  was modeled as:

$$\bar{r}_{init} = k_{init} (trigger) \frac{x_1}{K_{init,fII} + x_1} \quad (15)$$

where  $k_{init}$ ,  $K_{init,fII}$  are the rate and saturation constants governing initiation, respectively. The rate of initiation was modified by  $v_{init}$ , the control parameter governing initiation. Initiation was sensitive to the level of trigger (activator) and TFPI (inhibitor):

$$v_{init} = \min (f_{init}^-(TFPI), f_{init}^+(trigger)) \quad (16)$$

where the transfer functions  $f$  took the form of Eqn (9). The rate of thrombin amplification was given by:

$$\bar{r}_{amp} = k_{amp} (x_2) \frac{x_1}{K_{amp,fII} + x_1} \quad (17)$$

where  $k_{amp}$ ,  $K_{amp,fII}$  denote the rate and saturation constants governing amplification, respectively. The amplification control term, which modified amplification rate, was modeled as a combination of multiple inhibition terms and one activation term:

$$v_{amp} = \min \left( f_{amp}^{-}(TFPI), f_{amp}^{-}(x_4), f_{amp}^{+}(\mathcal{Z}_{amp}) \right) \quad (18)$$

where  $\mathcal{Z}_{amp} = fV \times fX \times fVIII \times fIX$ . Although  $f_{amp}^{+}(\mathcal{Z}_{amp})$  is an activating term, we included it in the min integration rule; the factors in  $\mathcal{Z}_{amp}$  were essential for amplification (if any of these factors was missing the amplification reaction would not occur). Thus, the factors in  $\mathcal{Z}_{amp}$  were required components, a classification that we implemented by the min selection rule. The rate activated protein C formation was given by:

$$\bar{r}_{apc} = k_{APC,formation}(TM) \frac{x_3}{K_{formation,PC} + x_3} \quad (19)$$

where  $k_{APC,formation}$  and  $K_{formation,PC}$  denote the rate and saturation constants governing activated protein C formation, respectively and  $TM$  denotes the thrombomodulin abundance. We modeled the control term which governed APC formation as a single thrombin-dependent activation term:

$$v_{apc} = \max \left( f_{apc}^{+}(x_2) \right) \quad (20)$$

Lastly, we included direct irreversible inhibition of FIIa by ATIII:

$$\bar{r}_{inh,ATIII} = k_{ATIII,inhibition}(x_5 x_2^{\gamma}) \quad (21)$$

where  $\gamma$  was estimated to be  $\gamma = 1.26$ . For ATIII inhibition of FIIa, the control variables  $v_{inh,ATIII}$  was taken to be unity. The model equations were encoded using the Python programming language and solved using the ODEINT routine of the SciPy module [59].

The model files can be downloaded from <http://www.varnerlab.org>.

**Estimation of model parameters from experimental data.** Model parameters were estimated by minimizing the difference between simulations and experimental thrombin measurements (squared residual):

$$\min_{\mathbf{k}} \sum_{\tau=1}^{\mathcal{T}} \sum_{j=1}^{\mathcal{S}} \left( \frac{\hat{x}_j(\tau) - x_j(\tau, \mathbf{k})}{\omega_j(\tau)} \right)^2 \quad (22)$$

where  $\hat{x}_j(\tau)$  denotes the measured value of species  $j$  at time  $\tau$ ,  $x_j(\tau, \mathbf{k})$  denotes the simulated value for species  $j$  at time  $\tau$ , and  $\omega_j(\tau)$  denotes the experimental measurement variance for species  $j$  at time  $\tau$ . The outer summation is with respect to time, while the inner summation is with respect to state. We minimized the model residual using Particle swarm optimization (PSO) [60]. PSO uses a *swarming* metaheuristic to explore parameter spaces. A strength of PSO is its ability to find the global minimum, even in the presence of potentially many local minima, by communicating the local error landscape experienced by each particle collectively to the swarm. Thus, PSO acts both as a local and a global search algorithm. For each iteration, particles in the swarm compute their local error by evaluating the model equations using their specific parameter vector realization. From each of these local points, a globally best error is identified. Both the local and global error are then used to update the parameter estimates of each particle using the rules:

$$\Delta_i = \theta_1 \Delta_i + \theta_2 \mathbf{r}_1 (\mathcal{L}_i - \mathbf{k}_i) + \theta_3 \mathbf{r}_2 (\mathcal{G} - \mathbf{k}_i) \quad (23)$$

$$\mathbf{k}_i = \mathbf{k}_i + \Delta_i \quad (24)$$

where  $(\theta_1, \theta_2, \theta_3)$  are adjustable parameters,  $\mathcal{L}_i$  denotes the local best solution found by particle  $i$ , and  $\mathcal{G}$  denotes the best solution found over the entire population of particles. The quantities  $r_1$  and  $r_2$  denote uniform random vectors with the same dimen-

sion as the number of unknown model parameters ( $\mathcal{K} \times 1$ ). In this study, we used  $(\theta_1, \theta_2, \theta_3) = (1.0, 0.05564, 0.02886)$ . The quality of parameter estimates was measured using goodness of fit (model residual). The particle swarm optimization routine was implemented in the Python programming language. All plots were made using the Matplotlib module of Python [61].

**Global sensitivity analysis of model performance.** We conducted a global sensitivity analysis, using the variance-based method of Sobol, to estimate which parameters controlled the performance of the reduced order model [62]. We computed the total sensitivity index of each parameter relative to two performance objectives, the peak thrombin time and the area under the thrombin curve (thrombin exposure). We established the sampling bounds for each parameter from the minimum and maximum value of that parameter in the parameter set ensemble. We used the sampling method of Saltelli *et al.* [63] to compute a family of  $N(2d + 2)$  parameter sets which obeyed our parameter ranges, where  $N$  was the number of trials, and  $d$  was the number of parameters in the model. In our case,  $N = 10,000$  and  $d = 22$ , so the total sensitivity indices were computed from 460,000 model evaluations. The variance-based sensitivity analysis was conducted using the SALib module encoded in the Python programming language [64].

## Acknowledgements

This study was supported by an award from the Army Research Office (ARO #59155-LS).



## References

1. Kholodenko B, Yaffe MB, Kolch W (2012) Computational approaches for analyzing information flow in biological networks. *Sci Signal* 5: re1.
2. Gadkar KG, Varner J, Doyle FJ (2005) Model identification of signal transduction networks from data using a state regulator problem. *Syst Biol (Stevenage)* 2: 17–30.
3. Bailey JE (2001) Complex biology with no parameters. *Nat Biotechnol* 19: 503-4.
4. Machta BB, Chachra R, Transtrum MK, Sethna JP (2013) Parameter space compression underlies emergent theories and predictive models. *Science* 342: 604-7.
5. Tasseff R, Nayak S, Song SO, Yen A, Varner JD (2011) Modeling and analysis of retinoic acid induced differentiation of uncommitted precursor cells. *Integr Biol (Camb)* 3: 578-91.
6. Morris MK, Saez-Rodriguez J, Sorger PK, Lauffenburger DA (2010) Logic-based models for the analysis of cell signaling networks. *Biochemistry* 49: 3216-24.
7. Saez-Rodriguez J, Alexopoulos LG, Zhang M, Morris MK, Lauffenburger DA, et al. (2011) Comparing signaling networks between normal and transformed hepatocytes using discrete logical models. *Cancer Res* 71: 5400-11.
8. Morris MK, Saez-Rodriguez J, Clarke DC, Sorger PK, Lauffenburger DA (2011) Training signaling pathway maps to biochemical data with constrained fuzzy logic: quantitative analysis of liver cell responses to inflammatory stimuli. *PLoS Comput Biol* 7: e1001099.
9. Morris MK, Shriver Z, Sasisekharan R, Lauffenburger DA (2012) Querying quantitative logic models (q2lm) to study intracellular signaling networks and cell-cytokine interactions. *Biotechnol J* 7: 374-86.
10. Butenas S, Mann KG (2002) Blood coagulation. *Biochemistry (Mosc)* 67: 3-12.
11. Schenone M, Furie BC, Furie B (2004) The blood coagulation cascade. *Curr Opin Hematol* 11: 272-7.

12. Adams RLC, Bird RJ (2009) Review article: Coagulation cascade and therapeutics update: relevance to nephrology. part 1: Overview of coagulation, thrombophilias and history of anticoagulants. *Nephrology (Carlton)* 14: 462-70.
13. Goldhaber SZ, Colman RW, Clowes AW, editors (2006) *Hemostasis and Thrombosis: Basic Principles and Clinical Practice*. Lippincott Williams and Wilkins.
14. Brummel KE, Paradis SG, Butenas S, Mann KG (2002) Thrombin functions during tissue factor-induced blood coagulation. *Blood* 100: 148-52.
15. Mann K, Nesheim M, Church W, Haley P, Krishnaswamy S (1990) Surface-dependent reactions of vitamin k-dependent enzyme complexes. *Blood* 76: 1-16.
16. Roberts H, Monroe D, Oliver J, Chang J, Hoffman M (1998) Newer concepts of blood coagulation. *Haemophilia* 4: 331-334.
17. Mann K (1999) Biochemistry and physiology of blood coagulation. *Thromb Haemost* 82: 165-174.
18. Khanin MA, Semenov VV (1989) A mathematical model of the kinetics of blood coagulation. *J Theor Biol* 136: 127-34.
19. Willems GM, Lindhout T, Hermens WT, Hemker HC (1991) Simulation model for thrombin generation in plasma. *Haemostasis* 21: 197-207.
20. Baldwin SA, Basmadjian D (1994) A mathematical model of thrombin production in blood coagulation, part i: The sparsely covered membrane case. *Ann Biomed Eng* 22: 357-70.
21. Leipold RJ, Bozarth TA, Racanelli AL, Dicker IB (1995) Mathematical model of serine protease inhibition in the tissue factor pathway to thrombin. *J Biol Chem* 270: 25383-7.
22. Kuharsky AL, Fogelson AL (2001) Surface-mediated control of blood coagulation: the role of binding site densities and platelet deposition. *Biophys J* 80: 1050-74.
23. Luan D, Zai M, Varner JD (2007) Computationally derived points of fragility of a human

cascade are consistent with current therapeutic strategies. PLoS Comput Biol 3:  
e142.

24. Luan D, Szlam F, Tanaka KA, Barie PS, Varner JD (2010) Ensembles of uncertain  
mathematical models can identify network response to therapeutic interventions. Mol  
Biosyst 6: 2272-86.

25. Jones KC, Mann KG (1994) A model for the tissue factor pathway to thrombin. ii. a  
mathematical simulation. J Biol Chem 269: 23367-73.

26. Hockin MF, Jones KC, Everse SJ, Mann KG (2002) A model for the stoichiometric  
regulation of blood coagulation. J Biol Chem 277: 18322-33.

27. Butenas S, Orfeo T, Gissel MT, Brummel KE, Mann KG (2004) The significance of  
circulating factor ixa in blood. J Biol Chem 279: 22875-82.

28. Lo K, Denney WS, Diamond SL (2005) Stochastic modeling of blood coagulation  
initiation. Pathophysiol Haemost Thromb 34: 80-90.

29. Chatterjee MS, Denney WS, Jing H, Diamond SL (2010) Systems biology of coagu-  
lation initiation: kinetics of thrombin generation in resting and activated human blood.  
PLoS Comput Biol 6.

30. Stalker TJ, Traxler EA, Wu J, Wannemacher KM, Cermignano SL, et al. (2013) Hier-  
archical organization in the hemostatic response and its relationship to the platelet-  
signaling network. Blood 121: 1875-85.

31. Leiderman K, Fogelson A (2014) An overview of mathematical modeling of thrombus  
formation under flow. Thromb Res 133 Suppl 1: S12-4.

32. Bannish BE, Keener JP, Fogelson AL (2014) Modelling fibrinolysis: a 3d stochastic  
multiscale model. Math Med Biol 31: 17-44.


33. Voronov RS, Stalker TJ, Brass LF, Diamond SL (2013) Simulation of intrathrombus  
fluid and solute transport using in vivo clot structures with single platelet resolution.  
Ann Biomed Eng 41: 1297-307.

34. Runyon MK, Johnson-Kerner BL, Ismagilov RF (2004) Minimal functional model of hemostasis in a biomimetic microfluidic system. *Angew Chem Int Ed Engl* 43: 1531-6.
35. Kastrup CJ, Runyon MK, Shen F, Ismagilov RF (2006) Modular chemical mechanism predicts spatiotemporal dynamics of initiation in the complex network of hemostasis. *Proc Natl Acad Sci U S A* 103: 15747-52.
36. Runyon MK, Johnson-Kerner BL, Kastrup CJ, Van Ha TG, Ismagilov RF (2007) Propagation of blood clotting in the complex biochemical network of hemostasis is described by a simple mechanism. *J Am Chem Soc* 129: 7014-5.
37. Runyon MK, Kastrup CJ, Johnson-Kerner BL, Ha TGV, Ismagilov RF (2008) Effects of shear rate on propagation of blood clotting determined using microfluidics and numerical simulations. *J Am Chem Soc* 130: 3458-64.
38. Kuepfer L, Peter M, Sauer U, Stelling J (2007) Ensemble modeling for analysis of cell signaling dynamics. *Nat Biotechnol* 25: 1001-6.
39. Song SO, Varner J (2009) Modeling and analysis of the molecular basis of pain in sensory neurons. *PLoS One* 4: e6758.
40. Song SO, Chakrabarti A, Varner JD (2010) Ensembles of signal transduction models using pareto optimal ensemble techniques (poets). *Biotechnol J* 5: 768-80.
41. Tasseff R, Nayak S, Salim S, Kaushik P, Rizvi N, et al. (2010) Analysis of the molecular networks in androgen dependent and independent prostate cancer revealed fragile and robust subsystems. *PLoS One* 5: e8864.
42. Lequieu J, Chakrabarti A, Nayak S, Varner JD (2011) Computational modeling and analysis of insulin induced eukaryotic translation initiation. *PLoS Comput Biol* 7: e1002263.
43. Tran LM, Rizk ML, Liao JC (2008) Ensemble modeling of metabolic networks. *Biophys J* 95: 5606-17.

44. GKHansson (2005) Inflammation, Atherosclerosis and Coronary Artery Disease. N Engl J Med 352: 1685 - 1695.
45. Tanaka KA, Key NS, Levy JH (2009) Blood coagulation: hemostasis and thrombin regulation. Anesth Analg 108: 1433-46.
46. Tuddenham E, Cooper D (1994) The molecular genetics of haemostasis and its inherited disorders., volume 25 of *Oxford monographs in medical genetics*. Oxford University Press.
47. Mannucci MP, Tuddenham EGD (2001) The hemophilias - from royal genes to gene therapy. N Engl J Med 344: 1773 - 1780.
48. Mitchell J, Phillott A (2008) Haemophilia and inhibitors 1: diagnosis and treatment. Nursing Times 104: 26–27.
49. Tomokiyo K, Nakatomi Y, Araki T, Teshima K, Nakano H, et al. (2003) A novel therapeutic approach combining human plasma-derived factors viia and x for haemophiliacs with inhibitors: evidence of a higher thrombin generation rate in vitro and more sustained haemostatic activity in vivo than obtained with factor viia alone. Vox Sanguinis 85: 290-299.
50. Hedner U (2008) Factor viia and its potential therapeutic use in bleeding-associated pathologies. Thromb Haemost 100: 557–562.
51. Talbot M, Tien HC (2009) The use of recombinant factor viia in trauma patients. J Am Acad Orthop Surg 17: 477-81.
52. Shapiro AD (2008) Single-dose recombinant activated factor vii for the treatment of joint bleeds in hemophilia patients with inhibitors. Clin Adv Hematol Oncol 6: 579–586.
53. Duchesne JC, Mathew KA, Marr AB, Pinsky MR, Barbeau JM, et al. (2008) Current evidence based guidelines for factor viia use in trauma: the good, the bad, and the ugly. Am Surg 74: 1159-65.


- 586 54. Butenas S, Brummel KE, Branda RF, Paradis SG, Mann KG (2002) Mechanism of  
587 factor viia-dependent coagulation in hemophilia blood. *Blood* 99: 923-30.
- 588 55. Danforth CM, Orfeo T, Everse SJ, Mann KG, Brummel-Ziedins KE (2012) Defining  
589 the boundaries of normal thrombin generation: investigations into hemostasis. *PLoS*  
590 *One* 7: e30385.
- 591 56. Naito K, Fujikawa K (1991) Activation of human blood coagulation factor XI indepen-  
592 dent of factor XII. *J Biol Chem* 266: 7353-7358.
- 593 57. Gailani D, Broze GJ Jr (1991) Factor xi activation in a revised model of blood coagu-  
594 lation. *Science* 253: 909-12.
- 595 58. Smith SA, Mutch NJ, Baskar D, Rohloff P, Docampo R, et al. (2006) Polyphosphate  
596 modulates blood coagulation and fibrinolysis. *Proc Natl Acad Sci U S A* 103: 903-8.
- 597 59. Jones E, Oliphant T, Peterson P (2001–). *SciPy: Open source scientific tools for*  
598 *Python*. <http://www.scipy.org/>.
- 599 60. Kennedy J, Eberhart R (1995) Particle swarm optimization. In: *Proceedings of the*  
600 *International Conference on Neural Networks*. pp. 1942 - 1948.
- 601 61. Hunter JD (2007) Matplotlib: A 2d graphics environment. *Computing in Science and*  
602 *Engineering* 9: 90 - 95.
- 603 62. Sobol I (2001) Global sensitivity indices for nonlinear mathematical models and their  
604 monte carlo estimates. *Mathematics and Computers in Simulation* 55: 271 - 280.
- 605 63. Saltelli A, Annoni P, Azzini I, Campolongo F, Ratto M, et al. (2010) Variance based  
606 sensitivity analysis of model output. design and estimator for the total sensitivity index.  
607 *Computer Physics Communications* 181: 259 - 270.
- 608 64. Herman JD. <http://jdherman.github.io/salib/>.
- 609 65. Butenas S, van't Veer C, Mann KG (1999) "normal" thrombin generation. *Blood* 94:  
610 2169-78.
- 611 66. Allen GA, Hoffman M, Roberts HR, Monroe DM (2006) Manipulation of prothrombin

612 concentration improves response to high-dose factor VIIa in a cell-based model of  
613 haemophilia. Br J Haematology 134: 314 - 319.




**Fig. 1:** Schematic of the connectivity of the reduced order coagulation model. A trigger compound, e.g., TF/FVIIa initiates thrombin production (FIIa) from prothrombin (fII). Once activated, thrombin catalyzes its own activation (amplification step), as well as its own inhibition via the conversion of protein C to activated protein C (APC). APC and tissue factor pathway inhibitor (TFPI) inhibit initiation and amplification, while antithrombin III (ATIII) directly inhibits thrombin. All inhibition steps and trigger-induced initiation were modeled using a rule-based approach. Likewise, the dependence of amplification on other coagulation factors was also modeled using a rule-based approach. The abundance of the highlighted species (in the dashed boxes) was governed by an ordinary differential equation. All other species were assumed to be constant.







**Fig. 2:** Schematic of rule-based effective control laws. Traditional enzyme kinetic expressions, e.g., Michaelis–Menten or multiple saturation kinetics are multiplied by an enzyme activity control variable  $0 \leq v_j \leq 1$ . Control variables are functions of many possible regulatory factors encoded by arbitrary transfer functions of the form  $0 \leq f_j(\mathcal{Z}) \leq 1$ . At each simulation time step, the  $v_j$  variables are calculated by evaluating integration rules such as the max or min of the set of transfer functions  $f_1, \dots, f_n$  influencing the activity of enzyme  $E_j$ .




**Fig. 3:** Reduced order coagulation model training simulations. Reduced order coagulation model parameters were estimated using particle swarm optimization (PSO) with and without the protein C pathway as a function of prothrombin. Solid lines denote the simulated mean value of the thrombin profile for  $N = 20$  independent particles, points denote experimental data. The shaded region denotes the 99% confidence estimate of the mean simulated thrombin value (uncertainty in the model simulation). (A,B,C) training results for 150%, 100% and 50% of physiological prothrombin levels in the absence of the protein C pathway. Thrombin generation was initiated using 5 pmol/L FVIIa-TF in the presence of 200  $\mu$ mol/L of phospholipid vesicles (PCPS). All factors and control proteins were at their physiological concentration unless others denoted. The experimental training data was reproduced from the study of Butenas *et al.* [65].




**Fig. 4:** Reduced order coagulation model training simulations. Reduced order coagulation model parameters were estimated using particle swarm optimization (PSO) with and without the protein C pathway as a function of prothrombin. Solid lines denote the simulated mean value of the thrombin profile for  $N = 20$  independent particles, points denote experimental data. The shaded region denotes the 99% confidence estimate of the mean simulated thrombin value (uncertainty in the model simulation). (A,B,C) training results for 150%, 100% and 50% of physiological prothrombin levels in the presence of the protein C pathway. Only APC pathway parameters were allowed to vary in the simulations on the right. Thrombin generation was initiated using 5 pmol/L FVIIa-TF in the presence of 200  $\mu\text{mol/L}$  of phospholipid vesicles (PCPS). All factors and control proteins were at their physiological concentration unless others denoted. The experimental training data was reproduced from the study of Butenas *et al.* [65].




**Fig. 5:** Reduced order coagulation model predictions versus experimental data for normal coagulation. The reduced order coagulation model parameter estimates were tested against data not used during model training. Simulations of different levels of prothrombin and ATIII were compared with experimental data in the absence of the protein C pathway. Solid lines denote the simulated mean value of the thrombin profile for  $N = 20$  independent particles, points denote experimental data. The shaded region denotes the 99% confidence estimate of the mean simulated thrombin value (uncertainty in the model simulation). (A,B,C,D) prediction results for (FII,ATIII): (50%,150%), (100%, 100%), (125%, 75%) and (150%, 50%) of physiological prothrombin and ATIII levels in the absence of the protein C pathway. Thrombin generation was initiated using 5 pmol/L FVIIa-TF in the presence of 200  $\mu\text{mol/L}$  of phospholipid vesicles (PCPS). All factors and control proteins were at their physiological concentration unless others denoted. The experimental validation data was reproduced from the study of Butenas *et al.* [65]




**Fig. 6:** Reduced order coagulation model predictions versus experimental data with and without FVIII and FIX. The reduced order coagulation model parameter estimates were tested against data not used during model training. Simulations of normal thrombin formation with ATIII and the protein C pathway were compared with thrombin formation in the absence of fVIII and fIX. Solid lines denote the simulated mean value of the thrombin profile for  $N = 20$  independent particles, points denote experimental data. The shaded region denotes the 99% confidence estimate of the mean simulated thrombin value (uncertainty in the model simulation). (A,B) prediction results for normal thrombin generation and thrombin generation in hemophilia. All factors and control proteins were at their physiological concentration unless others noted. The experimental validation data was reproduced from the study of Allen *et al.* [66].



**Fig. 7:** Reduced order coagulation model predictions of rFVIIa administration. A: Simulations of thrombin formation in the presence of ATIII and the protein C pathway were conducted for a range of trigger values (1x - 200x nominal) in the absence of fVIII and fIX. B: Comparison of thrombin generation for normal versus hemophilia for 10x nominal trigger. Solid lines denote the simulated mean value of the thrombin profile for  $N = 20$  independent particles. The peak thrombin time for normal coagulation ( $t^*$ ) is less than rFVIIa induced coagulation in hemophilia ( $t^{**}$ ), while the peak thrombin value was greater in normal coagulation. The shaded region denotes the 99% confidence estimate of the mean thrombin value (uncertainty in the model simulation). All factors and control proteins were at their physiological concentration unless others noted.



**Fig. 8:** Reaction flux distribution as a function of time for thrombin generation under normal (left), hemophilia (center) and rFVIIa treated hemophilia (right). Reaction flux was calculated for each particle at  $T = 0, 4, 6, 8, 10, 12, 14$  min after the initiation of coagulation. Reaction fluxes were calculated for each particle in the parameter ensemble ( $N = 20$ ). Blue colors denote low flux values while red colors denote high flux values.



**Fig. 9:** Global sensitivity analysis of the reduced order coagulation model with respect to the model parameters. A: Sensitivity analysis of the thrombin peak time for different prothrombin levels (150%,100% and 50% of the physiological value) as a function of activated protein C. B: Sensitivity analysis of the thrombin exposure for different prothrombin levels (150%,100% and 50% of the physiological value) as a function of activated protein C. Points denote the mean total sensitivity value, while the area around each point denotes the uncertainty in the sensitivity value. The gray dashed line denotes the 45° degree diagonal, if sensitivity values are equal for different conditions they will lie on the diagonal. Sensitivity values significantly above or below the diagonal indicate differentially important model parameters. The radius of the shaded region around each total sensitivity value was the maximum uncertainty in that value estimated by the Sobol method.