

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human NF-L in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 985917
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human NF-L Met1-Asp338 Accession # P07196
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.


	Recommended Concentration	Sample
Immunocytochemistry	8-25 µg/mL	See Below
Confirmation of Specificity using RNAscope ISH		

DATA

Immunocytochemistry




ICC: Ms xhNF-L U251 MG




ICC: Ms xhNF-L K562 (-)

NF-L in U-251 MG Human Cell Line. NF-L was detected in immersion fixed U-251 MG human glioblastoma cell line (positive staining) and K562 human chronic myelogenous leukemia cell line (negative staining) using Mouse Anti-Human NF-L Monoclonal Antibody (Catalog # MAB22162) at 8 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). Specific staining was localized to cytoplasm. Staining was performed using our protocol for Fluorescent ICC Staining of Non-adherent Cells.

Immunocytochemistry



ISH: Hs-NEFL U251 MG



ISH: Hs-NEFL K562 (-)

NF-L in U-251 MG human glioblastoma cell line using RNAscope[®] ISH. NF-L mRNA was detected in formalin-fixed U-251 MG human glioblastoma cell line probed with ACD RNAscope[®] Probe-Hs-NEFL (Catalog # 468671) and stained using ACD RNAscope[®] 2.5 HD Detection Reagents-RED (Catalog # 32260). This is an In Situ Hybridization control for NF-L expression.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

NF-L is a 68 kDa light chain cytoskeletal intermediate filament protein that is expressed in neurons. It associates with the 125 kDa NF-M and the 200 kDa NF-H to form neurofilaments.