

NEFL Monoclonal Antibody (DA2)

Product Details

Size	200 µg
Species Reactivity	Human, Mouse, Rat
Published Species	Rat, Mouse, Human
Host/Isotype	Mouse / IgG1, kappa
Class	Monoclonal
Type	Antibody
Clone	DA2
Conjugate	Unconjugated
Immunogen	Specifically recognizes a phosphate-independent epitope on NF-L.
Form	Liquid
Concentration	0.5 mg/mL
Purification	purified
Storage buffer	PBS
Contains	0.1% sodium azide
Storage conditions	-20°C
RRID	AB_2532995

Applications	Tested Dilution	Publications
Western Blot (WB)	0.5-2 µg/mL	9 Publications
Immunohistochemistry (IHC)	-	3 Publications
Immunohistochemistry (Paraffin) (IHC (P))	1:20	-
Immunohistochemistry (Frozen) (IHC (F))	-	1 Publication
Immunocytochemistry (ICC/IF)	1:20	2 Publications
ELISA (ELISA)	0.1-0.5 µg/mL	-
Immunoprecipitation (IP)	2-5 µg	-
Miscellaneous PubMed (Misc)	-	3 Publications


Product Specific Information

13-0400 detects the neurofilament, light chain in human, rat, and mouse samples.

Product Images For NEFL Monoclonal Antibody (DA2)


NEFL Antibody (13-0400)

Antibody specificity was demonstrated by detection of differential basal expression of the target across tissue tested owing to their inherent genetic constitution. Relative expression of NEFL was observed in PC-12 differentiated to neurons in comparison to PC-12 undifferentiated cells. Relative expression was also observed in Mouse Cerebellum, Mouse Brain and Rat Brain in comparison to Mouse Heart and Mouse Spleen using Anti-NEFL Monoclonal Antibody (DA2) (Product # 13-0400) in Western Blot. {RE}




NEFL Antibody (13-0400) in IHC (P)

Immunofluorescent analysis of the neurofilament light chain in paraffin-embedded mouse brain tissue (right) compared to a negative control without primary antibody (left). Tissue sections were deparaffinized with xylene, and rehydrated with ethanol. To expose target proteins, antigen retrieval was performed using 10mM sodium citrate (pH 6.0) and microwaved for 8-15 min. Following antigen retrieval, tissues were washed with water and PBS, and then blocked in 0.3% BSA for 30 min at room temperature. Tissues were then probed with a neurofilament light chain monoclonal antibody (Product # 13-0400) in 0.3% BSA at a dilution of 1:20 for 1 hour at 37°C. Tissues were then incubated with a Goat anti-Mouse IgG (H+L) Secondary Antibody, DyLight 488 conjugate for 1 hour at 37°C (green). Nuclei (blue) were stained with DAPI. Images were taken at 40X magnification.



NEFL Antibody (13-0400) in IHC (P)

Immunofluorescent analysis of the neurofilament light chain in paraffin-embedded human brain tissue (right) compared to a negative control without primary antibody (left). Tissue sections were deparaffinized with xylene, and rehydrated with ethanol. To expose target proteins, antigen retrieval was performed using 10mM sodium citrate (pH 6.0) and microwaved for 8-15 min. Following antigen retrieval, tissues were washed with water and PBS, and then blocked in 0.3% BSA for 30 min at room temperature. Tissues were then probed with a neurofilament light chain monoclonal antibody (Product # 13-0400) in 0.3% BSA at a dilution of 1:20 for 1 hour at 37°C. Tissues were then incubated with a Goat anti-Mouse IgG (H+L) Secondary Antibody, DyLight 488 conjugate for 1 hour at 37°C (green). Nuclei (blue) were stained with DAPI. Images were taken at 40X magnification.



[View more figures on thermofisher.com](https://www.thermofisher.com)

Western Blot (9)

Scientific reports	Year 2020
Cep215 is essential for morphological differentiation of astrocytes.	Species Mouse
"13-0400 was used in Western Blotting to show that Cep215 is involved in morphological differentiation of astrocytes. Cep215 was specifically localized at the glial processes as well as centrosomes in developing astrocytes."	
Authors: Kang D,Shin W,Yoo H,Kim S,Lee S,Rhee K	
Frontiers in cellular neuroscience	Year 2020
Impaired Neurofilament Integrity and Neuronal Morphology in Different Models of Focal Cerebral Ischemia and Human Stroke Tissue.	Species Mouse
"13-0400 was used in Western Blotting to systematically address ischemia-induced alterations of the neurofilament light (NF-L), medium (NF-M), and heavy (NF-H) subunits as well as of -internexin (INA)."	Dilution 1:1000
Authors: Mages B,Aleithe S,Altmann S,Blietz A,Nitzsche B,Barthel H,Horn AKE,Hobusch C,Härtig W,Krueger M, Michalski D	

View more WB references on thermofisher.com

Immunohistochemistry (3)

Proceedings. Biological sciences	Year 2016
Substituting mouse transcription factor Pou4f2 with a sea urchin orthologue restores retinal ganglion cell development.	Dilution 1:1000
"13-0400 was used in immunohistochemistry to characterize restoration of retinal ganglion cell development by substitution of mouse transcription factor Pou4f2 with a sea urchin orthologue"	
Authors: Mao CA,Agca C,Mocko-Strand JA,Wang J,Ullrich-Lüter E,Pan P,Wang SW,Arnone MI,Frishman LJ,Klein WH	
Nature cell biology	Year 2004
A NUDEL-dependent mechanism of neurofilament assembly regulates the integrity of CNS neurons.	Species Human Mouse
Authors: Nguyen MD,Shu T,Sanada K,Larivière RC,Tseng HC,Park SK,Julien JP,Tsai LH	

View more IHC references on thermofisher.com

More applications with references on thermofisher.com

- IHC (F) (1)
- ICC/IF (2)
- Misc (3)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.