

Stochastic volatility  
Recent developments and future directions

Vladimir Piterbarg  
Barclays Capital

# Introduction

- ▶ SV models are an established part of a quant toolbox.
- ▶ Many tools have been developed, many more need to be developed
- ▶ Fundamental tools for calibration
  - ▶ Markovian projection (MP)
  - ▶ Parameter averaging (PA)
  - ▶ Direct calibration to market
- ▶ Applications
  - ▶ Short rate and forward Libor models of interest rates
  - ▶ Hybrids models
  - ▶ Basket and spread models
- ▶ Fundamental properties
  - ▶ Moment explosions
  - ▶ Smile asymptotics

# Calibration

- ▶ For calibration, need fast methods for European option pricing
- ▶ A number of SV models for interest rates and hybrids have been put forward recently, with various approaches to calibration
- ▶ Many of these approaches can be aggregated into what we call the Markovian Projection method:

a generic, powerful framework for deriving closed-form approximations to European option prices

**Step 1** Apply Markovian projection to  $S(\cdot)$ , a technique to replace a complicated process with a simple one, preserving European option prices

**Step 2** Approximate conditional expected values required

**Step 3** Apply parameter averaging techniques to obtain time-independent coefficients from time-dependent

**Step 4** Hopefully a simple model is obtained, use known results.

# The Markovian projection

Theorem (Dupire 97, Gyongy 86) Let  $X(t)$  be given by

$$dX(t) = \alpha(t) dt + \beta(t) dW(t), \quad (1)$$

where  $\alpha(\cdot)$ ,  $\beta(\cdot)$  are adapted bounded stochastic processes such that (1) admits a unique solution.

Define  $a(t, x)$ ,  $b(t, x)$  by

$$\begin{aligned} a(t, x) &= E(\alpha(t) | X(t) = x), \\ b^2(t, x) &= E(\beta^2(t) | X(t) = x), \end{aligned}$$

Then the SDE

$$\begin{aligned} dY(t) &= a(t, Y(t)) dt + b(t, Y(t)) dW(t), \\ Y(0) &= X(0), \end{aligned} \quad (2)$$

admits a weak solution  $Y(t)$  that has the same one-dimensional distributions as  $X(t)$ .

► See [Dup97], [Gyö86]

# The Markovian projection, cont

- Remark 1** Since  $X(\cdot)$  and  $Y(\cdot)$  have the same one-dimensional distributions, the prices of European options on  $X(\cdot)$  and  $Y(\cdot)$  for all strikes  $K$  and expiries  $T$  will be the same. Thus, for the purposes of European option valuation and/or calibration to European options, we can replace a potentially very complicated process  $X(\cdot)$  with a much simpler Markov process  $Y(\cdot)$ , which we call the Markovian projection of  $X(\cdot)$ .
- Remark 2** The process  $Y(\cdot)$  follows what is known as a “local volatility” process. The function  $b(t, x)$  is often called “Dupire’s local volatility”

## The Markovian projection, cont

- ▶ If  $X(\cdot)$  itself came from a local volatility model (perhaps complicated), then replacing it with a (simpler) local vol model is probably the right thing to do. But:
- ▶ For SV models, better to replace “like for like”. Replace a (complicated) SV model with a (simpler) SV model.
  - ▶ Approximations to conditional expected values (the hard part – see eg [Atl06]) may be simpler
  - ▶ Errors of approximations will tend to “cancel out”
- ▶ Dupire-Gyongy theorem still works

**Corollary** If two processes have the same Dupire’s local volatility, the European option prices on both are the same for all strikes and expiries

- ▶ Future directions: better methods for conditional expected value calculations

## Simple SV model

- ▶ After applying the MP method, often get the SDEs of the form

$$\begin{aligned}dz(t) &= \theta(1 - z(t)) dt + \gamma(t) \sqrt{z(t)} dV(t), \\dS(t) &= (\beta(t)S(t) + (1 - \beta(t))S(0))\sigma(t) \sqrt{z(t)} dW(t),\end{aligned}\tag{3}$$

- ▶ Or, rather, we apply the MP method with the goal of obtaining the SDEs in this form
  - ▶ Choose  $z$  to be the square root process
  - ▶ Linearize the volatility term of  $S$
- ▶ Why? When parameters are constant, this is the (shifted) Heston model, a model with very efficient numerical methods for European option valuation, see [AA02].
- ▶ How to replace time-dependent parameters with constant? Parameter averaging. Proofs and details in [Pit05b], [Pit05a]

## Example of a simple averaging formula

- For motivation, a log-normal model with time-dependent volatility,

$$dS(t) = \sigma(t) S(t) dW(t).$$

- It is known that, an option value with expiry  $T_n$  in this model is equal to the Black-Scholes option value with “effective” volatility

$$\sigma_n = \left( \frac{1}{T_n} \int_0^{T_n} \sigma^2(t) dt \right)^{1/2}.$$

- Direct link between “model” parameter  $\sigma(t)$  and “market” parameters ( $\sigma_n$ )
- Much faster than using Black-Scholes for valuation of options and implied calculations



## Non-trivial example: averaging skew

- ▶ Time-dependent skew

$$dS(t) = \sigma(t) (\beta(t) S(t) + (1 - \beta(t)) S(0)) dW(t),$$

- ▶ Constant skew

$$d\bar{S}(t) = \sigma(t) (b\bar{S}(t) + (1 - b)\bar{S}(0)) dW(t).$$

- ▶ Given  $\beta(\cdot)$ , find  $b$  such that option prices for different strikes (same expiry  $T$ ) are matched between two models
- ▶ The main result. In the “small skew” limit,

$$b = \int_0^T \beta(t) w(t) dt,$$

where

$$w(t) = \frac{v^2(t) \sigma^2(t)}{\int_0^T v^2(t) \sigma^2(t) dt}, \quad v^2(t) = \int_0^t \sigma^2(s) ds.$$

# Averaging skew, cont

- ▶ Comments:

- ▶ “Total skew”  $b$  is the average of “local skews”  $\beta(t)$  with weights  $w(t)$
- ▶ Weights proportional to total variance, i.e. local slope further away matters more

- ▶ Example: constant volatility  $\sigma(t) \equiv \sigma$ ,

$$b = (T^2/2)^{-1} \int_0^T t \beta(t) dt.$$

- ▶ SV extensions available, see [Pit05b], [Pit05a]

- ▶ Future directions:

- ▶ Averaging methods for other models/parameters
- ▶ More accurate/rigorous averaging methods

## Direct calibration to market

- In equity/FX: Let  $\sigma_{\text{mkt}}(T, K)$  be market volatilities for all expiries  $T$  and strikes  $K$  (assumed known). Given an exogenous SV process  $z(t)$ , find  $b(t, x)$  such that the model

$$dS(t) = b(t, S(t)) \sqrt{z(t)} dW(t), \quad S(0) = S_0,$$

matches the market

- Define Dupire's market local volatility  $b_{\text{mkt}}(t, x)$  by the requirement that the local volatility model with  $b_{\text{mkt}}(t, x)$  matches the whole market. Easy to compute

$$b_{\text{mkt}}(t, x) = \frac{2\partial C / \partial t}{\partial^2 C / \partial x^2}.$$

- Then, from Theorem and Corollary,

$$b^2(t, x) = \frac{b_{\text{mkt}}^2(t, x)}{E(z(t) | S(t) = x)}. \quad (4)$$

- In practice  $E(z(t) | S(t) = x)$  is often computed numerically in a forward PDE in  $(S, z)$ . Slow and noisy.

## Direct calibration to market, cont

- Define a “proxy” process  $X(t)$  by

$$dX(t) = \tilde{b}(t, X(t)) \sqrt{z(t)} dW(t), \quad X(0) = S_0, \quad (5)$$

where  $\tilde{b}(t, x)$  is such that European options on  $X$  are easy to compute

- Define the “proxy” Dupire’s local volatility  $b_{\text{proxy}}(t, x)$  as before but for European options on  $X$  (not on market). Then

$$E(z(t) | X(t) = x) = \frac{b_{\text{proxy}}^2(t, x)}{\tilde{b}^2(t, x)}, \quad (6)$$

thus having a stochastic volatility model which cheaply-computable European option prices allows us to compute the conditional expected values easily.

- Combining the two results we get

$$b(t, x) = \tilde{b}(t, x) \times \frac{b_{\text{mkt}}(t, x)}{b_{\text{proxy}}(t, x)} \times \left( \frac{E(z(t) | X(t) = x)}{E(z(t) | S(t) = x)} \right)^{1/2}.$$

## Direct calibration to market, cont

- ▶ Choice 1: Approximate

$$E(z(t)|X(t) = x) = E(z(t)|S(t) = x)$$

- ▶ Choice 2: Link  $S(t)$  and  $X(t)$ .

- ▶ Define  $H(t, s)$  by the requirement that  $H(t, S(t))$  has the same  $dW$  term as  $dX$  ( $H$  a function of  $b, \tilde{b}$ )

- ▶ Then approximate

$$\begin{aligned} X(t) &\approx H(t, S(t)), \\ E(z(t)|S(t) = x) &\approx E(z(t)|X(t) = H(t, x)). \end{aligned}$$

- ▶ Functional equation on  $b$ ,

$$b(t, x) = \tilde{b}(t, H(t, x)) \frac{b_{\text{mkt}}(t, x)}{b_{\text{proxy}}(t, H(t, x))}. \quad (7)$$

- ▶ Last derivation is an example of a clever way of computing conditional expectations. **Need more of these in the future!**
- ▶ Original result due to Forde ([For06]). More details in [Pit06a].

# Local volatility short rate model

- ▶ Simplest interest rate model: one-factor Gaussian (“Hull-White”)

$$r(t) = f(0, t) + x(t), \quad dx(t) = (\theta(t) - ax(t)) dt + \sigma(t) dW(t).$$

- ▶ Local-volatility extension: quasi-Gaussian (“Cheyette”)

$$\begin{aligned} dx(t) &= (y(t) - ax(t)) dt + \sigma(t, x(t), y(t)) dW(t), \\ dy(t) &= (\sigma^2(t, x(t), y(t)) - 2ay(t)) dt. \end{aligned}$$

- ▶ Swap rate (under swap measure),  $S(t) = S(t, x(t), y(t))$  for a known function  $S(t, x, y)$ ,

$$dS(t) = \left. \frac{\partial S(t, x, y)}{\partial x} \right|_{x=x(t), y=y(t)} \sigma(t, x(t), y(t)) dW^A(t).$$

## Local volatility short rate model, cont

- ▶ Markovian projection (preserves European swaptions)

$$dS(t) = \eta(t, S(t)) dW^A(t),$$

$$\eta^2(t, S) = E^A \left( \left( \frac{\partial S(t, x(t), y(t))}{\partial x} \right)^2 \sigma^2(t, x(t), y(t)) \middle| S(t) = S \right)$$

- ▶ Let  $y^*(t) = E^A(y(t))$ ,  $\xi(t, s)$  is the inverse of  $S(t, x, y^*(t))$  in  $x$ . Then

$$\eta^2(t, S) \approx \left( \frac{\partial S(t, x, y^*(t))}{\partial x} \middle|_{x=\xi(t, S)} \right) \sigma(t, \xi(t, S), y^*(t)).$$

- ▶ Local-volatility model for  $S$  with a known  $\eta$ . Apply parameter averaging (on skew and vol), then shifted-lognormal formula to get option prices.
- ▶ Similar approaches for SV extensions, forward Libor models with SV, and IR/Equity, IR/FX hybrids models. See details in [And05], [Pit05a], [Pit06b].

# Multi-stochastic volatility

- ▶ When modelling multiple underlyings, natural to have different SV processes for them: baskets, spreads
- ▶ Also in current models: in interest rates, turns out variance decorrelation is important for exotics linked to CMS spreads, see [Pit06c]

- ▶ Consider a “simple” multi-SV model for a basket

$$S(t) = \sum w_i S_i(t),$$

$$dS_i(t) = \varphi_i(S_i(t)) \sqrt{z_i(t)} dW_i(t),$$

$$dz_i(t) = \theta(1 - z_i(t)) dt + \eta_i \sqrt{z_i(t)} dW_{I+i}(t), \quad z_i(0) = 1,$$

$$i = 1, \dots, I.$$

- ▶ Options on index  $S(\cdot)$ . Apply MP to write SDE for  $S$ . Start

$$dS(t) = \sigma(t) dW(t),$$

$$\sigma^2(t) = \sum_{n,m=1}^N w_n w_m \varphi_n(S_n(t)) \varphi_m(S_m(t)) \sqrt{z_n(t) z_m(t)} \rho_{nm}.$$



## Multi-stochastic volatility, cont

- What to use for  $z(t)$ , the SV of the index? Inspired by constant case  $\varphi_i(S_i(t)) \equiv \varphi_i(S_i(0)) = p_i$ ,

$$\begin{aligned} z(t) &= p^{-2} \sum w_n w_m p_n p_m \sqrt{z_n(t) z_m(t)} \rho_{nm}, \\ p^2 &= \sum w_n w_m p_n p_m \rho_{nm}. \end{aligned}$$

Then

$$\begin{aligned} dS(t) &= \varphi(t, S(t)) \sqrt{z(t)} dW(t), \\ \varphi^2(t, x) &= E(\sigma^2(t) | S(t) = x) / E(z(t) | S(t) = x). \end{aligned}$$

- Use Gaussian approximation to  $(S_i, z_i)$  to compute  $\varphi^2(t, x)$
- Approximate the dynamics of  $z(\cdot)$  by the mean reverting square root process – also use MP to come up with the coefficients!
- See more in [Pit06a], and an alternative approach in [DK06]

# Moment explosions

- ▶ SV models widely used, but do we know their fundamental properties?

$$\begin{aligned}dX(t) &= \lambda X(t) \sqrt{z(t)} dW(t), \\dz(t) &= \kappa(\theta - z(t)) dt + \varepsilon z^p(t) dB(t), \quad \langle dW, dB \rangle = \rho.\end{aligned}$$

- ▶ Well-known failure of the martingale property: When  $p \leq \frac{1}{2}$  or  $p > \frac{3}{2}$ ,  $X$  is a martingale. When  $\frac{3}{2} > p > \frac{1}{2}$ ,  $X$  is a martingale for  $\rho \leq 0$  and a strict supermartingale for  $\rho > 0$ .
- ▶ Many tools not valid anymore (Girsanov's theorem, etc)
- ▶ Even worse. Moments of  $X(\cdot)$  may explode in finite time. See full details in [AP06]

## Moment explosions, cont

| p                | $\rho$          | $\omega$ | Result   |
|------------------|-----------------|----------|--|
| $0 < p < 1/2$    | $-1 < \rho < 1$ | $\geq 0$ | $EX^\omega(T) < \infty$ for $\forall T$                    |
| $p = 1/2$        | $-1 < \rho < 1$ | $> 1$    | $\exists T^*: EX^\omega(T) = \infty$ for $\forall T > T^*$ |
| $1/2 < p \leq 1$ | $-1 < \rho < 0$ | small    | $EX^\omega(T) < \infty$ for $\forall T > 0$ .              |
| $1/2 < p \leq 1$ | $-1 < \rho < 0$ | large    | $\exists T^*: EX^\omega(T) = \infty$ for $\forall T > T^*$ |
| $1/2 < p \leq 1$ | $\rho = 0$      | $> 1$    | $EX^\omega(T) = \infty$ for $\forall T$                    |
| $1/2 < p < 3/2$  | $0 < \rho < 1$  | large    | $EX^\omega(T) = \infty$ for $\forall T$                    |

- ▶ Second moment of X often important, in particular in interest rate applications
  - ▶ If X is a Libor rate ,then the price of Libor-in-arrears involves  $EX^2(T)$ .
  - ▶ If X is a swap rate, the price of CMS involves  $EX^2(T)$ .
  - ▶ Numerical tricks to make them finite? A lot of issues.

# Moment explosions, cont

- ▶ Consider SABR model

$$\begin{aligned}dX(t) &= \lambda X^c(t) \sqrt{z(t)} dW(t), \\dz(t) &= \frac{1}{4} \varepsilon^2 z(t) dt + \varepsilon z(t) dB(t),\end{aligned}$$

- ▶ Case  $c < 1$ : all moments finite,  $X$  is a non-negative martingale;
- ▶ Case  $c = 1$ : (same classification as before, despite the positive drift)
  - ▶ for small  $\omega$ ,  $EX^\omega(T) < \infty$ ,
  - ▶ for large  $\omega$ ,  $\exists T^*$ :  $EX^\omega(T) = \infty$  for  $\forall T > T^*$ .

# Smile asymptotics

- ▶ Roger Lee ([Lee04]) was the first to link the number of finite moments to the asymptotics of the smile.
- ▶ Define  $I(k)$  the implied BS volatility as a function of log strike,  $k = \log(K/S_0)$ . Define  $\beta = \limsup_{k \rightarrow \infty} \frac{I^2(k)}{k/T}$ . Then

$$\begin{aligned}\beta &= 2 - 4 \left[ \sqrt{\omega_{\max}^2 + \omega_{\max}} - \omega_{\max} \right], \\ \omega_{\max} &= \arg \sup \left\{ \omega : EX^{1+\omega}(T) < \infty \right\}.\end{aligned}$$

- ▶ Together with moment explosion results, gives smile asymptotics in many cases
- ▶ However, if all moments exist (eg SABR), not very informative:  $\limsup_{k \rightarrow \infty} \frac{I^2(k)}{k/T} = 0$ , so which one is it
  - ▶  $I(k)$  grows slower than  $\sqrt{k}$ ?
  - ▶  $I(k)$  approaches non-zero limit?
  - ▶  $I(k) \rightarrow 0$ ?

## Smile asymptotics, cont

- ▶ A refinement (stated by not proven in [Pit04], simpler case  $v(k) = 1$  proven in [For06])
- ▶ Let  $v(k)$  be such that  $\delta^2 = \lim_{k \rightarrow -\infty} v^2(k)/k$  exists. Then

$$\limsup_{k \rightarrow \infty} \frac{I(k)}{v(k)} = a^*,$$

where  $a^*$  is a solution to

$$a\sqrt{2q_{\max}T} = 1 - \frac{a^2\delta^2}{2}$$

and

$$q_{\max} = \arg \sup \left\{ q : \mathbb{E} \exp \left( q \left[ \frac{\log(1 + X(T))}{v(\log(1 + X(T)))} \right]^2 \right) < \infty \right\}.$$

- ▶ Lee's result:  $v(k) = \sqrt{k/T}$ .
- ▶  $v(k) = 1$ : the limit of  $I(k)$

## Smile asymptotics, cont

- ▶ Recently, Benaim and Friz (see [BF06]) completely characterized implied vol smile asymptotics. Define
  - ▶  $\psi(x) = 2 - 4 [\sqrt{x^2 + x} - x]$ ,
  - ▶  $f \sim g \Leftrightarrow f/g \rightarrow 1$  as  $x \rightarrow \infty$ ,
  - ▶  $R_\alpha$  to be regularly varying functions of index  $\alpha$  at  $+\infty$  (eg “similar” to  $x^\alpha$ )
  - ▶  $f(x)$  the density of  $\log X(T)$ .
- ▶ Main result:
  - ▶ If  $-\log f(k) \in R_\alpha$ , then  $I^2(k)/k \sim \psi(-1 - \log f(k)/k)$ ;
  - ▶ If  $-\log f(k)/k \rightarrow \infty$  (ie all moments exist) then  $I^2(k) \sim -1/(2 \log f(k))$
- ▶ Future challenge: **find smile asymptotics for SABR.**  
**Conjecture: for  $\text{cev} < 1$ , the limit is finite and non-zero.**





# Future challenges

Would like to see progress in the following directions

- ▶ New ways to compute/approximate  $E(z(t)|X(t))$  found: will lead to advances in SV calibration
- ▶ Parameter averaging methods extended to new models and improved
- ▶ More types of models handled by MP+PA
- ▶ Calibration and approximation tools developed for multi-stochastic volatility models (baskets, spreads)
- ▶ Volatility smile asymptotics for the SABR model found



# Bibliography I

-  Leif B.G. Andersen and Jesper Andreasen.  
Volatile volatilities.  
Risk, 15(12):163–168, December 2002.
-  Jesper Andreasen.  
Back to the future.  
Risk Magazine, 18(9):104–109, September 2005.
-  Leif B.G. Andersen and Vladimir V. Piterbarg.  
Moment explosions in stochastic volatility models.  
To appear in Finance and Stochastics, 2006.
-  Marc Atlan.  
Localizing volatilities.  
arXiv working paper, April 2006.

# Bibliography II



Shalom Benaim and Peter Friz.

Regular variation and smile asymptotics.  
arXiv working paper, March 2006.



Valdo Durrleman and Nicole El Karoui.

Coupling smiles.  
Working paper, May 2006.



Bruno Dupire.

A unified theory of volatility.  
Banque Paribas working paper, 1997.



Martin Forde.

Calibrating local-stochastic volatility models.  
University of Bristol working paper,  
<http://www.maths.bris.ac.uk/~mamsf/>, February 2006.

# Bibliography III



I. Gyöngy.

Mimicking the one-dimensional distributions of processes having an Ito differential.

Prob. Th. Rel. Fields, 71:501 – 516, 1986.



Roger W. Lee.

The moment formula for implied volatility at extreme strikes.

Mathematical Finance, 14:469 – 479, 2004.



Vladimir V. Piterbarg.

Implied volatility smile when all moments are finite.

Working paper, June 2004.



Vladimir V. Piterbarg.

Stochastic volatility model with time-dependent skew.

Applied Mathematical Finance, 12(2):147–185, June 2005.

# Bibliography IV



Vladimir V. Piterbarg.

Time to smile.

Risk Magazine, 18(5):71–75, May 2005.



Vladimir V. Piterbarg.

Markovian projection method for volatility calibration.

SSRN working paper, June 2006.



Vladimir V. Piterbarg.

Smiling hybrids.

Risk Magazine, 19(5):66–71, May 2006.

# Bibliography V



Vladimir V. Piterbarg.

Towards a multi-stochastic volatility model for CMS spread  
exotics.

Global Derivatives Trading and Risk Management Conference,  
Paris, May 2006.