




Probabilidad de que $t_{(10)}$ sea menor que -2

```
pt(q = -2, df = 10)
```

```
[1] 0.03669402
```

$$P(t_{(10)} < -2) = 0.0367$$

$$t_{10, 0.0367} = -2$$

$$t_{(0.49, 90)} = -0.0251 \approx Z_{0.49}$$

Media

$$\mu_X = E(X) = 0 \quad \text{para } \nu > 1$$

Varianza

$$\sigma_X^2 = V(X) = \frac{\nu}{\nu - 2} \quad \text{para } \nu > 2$$

```
> x1 = rt(n = 600, df = 140)
> mean(x1)
[1] 0.006157374 →  $\bar{x}$ 
> var(x1)
[1] 0.9923853 →  $s^2$ 
```

```
> x1 = rt(n = 600, df = 140)
> mean(x1)
[1] 0.02794208
> var(x1)
[1] 0.952294
```

$$V = 140$$

$$\sigma_x^2 = \frac{140}{138} \approx 1.01$$

Media

$$\mu_X = E(X) = \frac{d_2}{d_2 - 2}, \quad \text{si } d_2 > 2$$

Varianza

$$\sigma_X^2 = V(X) = \frac{2d_2^2(d_1 + d_2 - 2)}{d_1(d_2 - 2)^2(d_2 - 4)}, \quad \text{si } d_2 > 4$$

```
> x <- rf(n = 700, df1 = 25, df2 = 17)
> mean(x)
[1] 1.151256
> var(x)
[1] 0.318123
```

```
> x <- rf(n = 700, df1 = 25, df2 = 17)
> mean(x)
[1] 1.124213
> var(x)
[1] 0.3051615
```

$$1h \quad 1h \quad 1h \quad 1h \quad 1h \quad \dots$$

13	10	12	15	12	...
----	----	----	----	----	-----

$$1h \leftarrow 1h \leftarrow 1h \leftarrow \dots$$
