


Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/hen

Effect of liquid phase plasma on photocatalysis of water for hydrogen evolution

Sangmin Jeong^a, Kyong-Hwan Chung^a, Heon Lee^a, Hyunwoong Park^b,
Ki-Joon Jeon^c, Young-Kwon Park^d, Sang-Chul Jung^{a,*}

^a Department of Environmental Engineering, Sunchon National University, 255 Jungang-ro, Sunchon, Jeonnam 57922, Republic of Korea

^b School of Energy Engineering, Kyungpook National University, 80 Daehakro, Bukgu, Daegu 41566, Republic of Korea

^c Department of Environmental Engineering, Inha University, 100 Inharo, Nam-gu, Incheon 22212, Republic of Korea

^d School of Environmental Engineering, University of Seoul, 163 Seoulsiripdaero, Dongdaemun-gu, Seoul 02504, Republic of Korea

ARTICLE INFO

Article history:

Received 1 November 2016

Received in revised form

3 March 2017

Accepted 27 March 2017

Available online 15 April 2017

Keywords:

Liquid phase plasma

Photocatalytic decomposition

Hydrogen evolution

Metal-loaded TiO₂

Na-Y support

ABSTRACT

This study examined the effects of liquid phase plasma irradiation on the photocatalytic decomposition of water for hydrogen evolution. TiO₂ and metal-loaded TiO₂ nanocrystallites were introduced as photocatalysts. Na-Y zeolite was applied as a support for the TiO₂ nanocrystallites. The photocatalytic activities of the photocatalysts were estimated for hydrogen production from water. Hydrogen evolution appeared in the photodecomposition of water without photocatalysts. This was caused by the decomposition of water by plasma irradiation in water directly. The hydrogen evolution efficiency improved with increasing conductivity of water. The rate of hydrogen evolution was increased by the metal loading (Ni, Co, Fe) on the TiO₂ surface. Na-Y zeolite can be used as an efficient photocatalytic support for the fixation of TiO₂. The TiO₂ nanocrystallites were incorporated above 40 wt% on Na-Y support.

© 2017 Hydrogen Energy Publications LLC. Published by Elsevier Ltd. All rights reserved.

Introduction

Photocatalysis is a favorable method for hydrogen production because it can use solar energy sustainably and efficiently. This process is attractive economically more conventional methods compared to steam reforming process and water electrolysis [1,2]. Photocatalytic water splitting is an effective method for converting solar energy to hydrogen as clean and renewable hydrogen energy [3–5]. Many studies have been

attempted to develop a photosensitive catalysts under UV as well as visible light illumination [6–9]. Recently, the development of the visible light sensitive photocatalysts for the hydrogen generation from water has attracted considerable attention [9–15]. In addition, the light sources and photocatalysts are very important in a photochemical reaction. Although a range of light sources have been employed in photocatalysis, there are few reports on photocatalysis using liquid phase plasma (LPP) by irradiation into water directly [16].

* Corresponding author. Fax: +82 61 750 3810.

E-mail address: jsc@sunchon.ac.kr (S.-C. Jung).

<http://dx.doi.org/10.1016/j.ijhydene.2017.03.183>

0360-3199/© 2017 Hydrogen Energy Publications LLC. Published by Elsevier Ltd. All rights reserved.