


Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/hydro

The effect of solvent and ionomer on agglomeration in fuel cell catalyst inks: Simulation by the Discrete Element Method


Magnus So, Tomohiro Ohnishi, Kayoung Park, Masumi Ono, Yoshifumi Tsuge, Gen Inoue*

Department of Chemical Engineering, Kyushu University, 744 Motooka, Nishi-ku, Fukuoka 819-0395 Japan

HIGHLIGHTS

- A particle model to simulate agglomeration in fuel cell catalyst ink was developed.
- It comprises a new model of steric force due to ionomer between carbon aggregates.
- This force was modelled as a function of the dielectric constant of the solvent.
- The model can predict the agglomerate sizes observed in various ink experiments.
- It can be utilized to optimize the solvent for best catalyst porous structures.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 5 July 2019

Received in revised form

28 August 2019

Accepted 1 September 2019

Available online 8 October 2019

Keywords:

Dielectric constant

Discrete Element Method

ABSTRACT

We simulate agglomeration in different fuel cell catalyst ink solutions using Discrete Element Method. Carbon support is modelled as particles in various inks with ionomer and various solvents. The particles interact with particle-pair forces resulting in agglomerate build-up. The classical colloidal theory with van der Waals and electric double layer forces underestimates the ink stability, which motivates the development of a new model of polymer force between particles. The force is activated when there is a bridging of polymer between the carbon black particles, and the strength is dependent on the ionomer interaction with the solvent by the dielectric constant. A critical dielectric constant was defined for which ionomer form a web-like polymer network that increases the ink stability. This modification can explain the trend of the ink stability, and the model can simulate the

* Corresponding author.

E-mail address: ginoue@chem-eng.kyushu-u.ac.jp (G. Inoue).

<https://doi.org/10.1016/j.ijhydene.2019.09.012>

0360-3199/© 2019 Hydrogen Energy Publications LLC. Published by Elsevier Ltd. All rights reserved.