


Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/he

Effect of surface properties controlled by Ce addition on CO₂ methanation over Ni/Ce/Al₂O₃ catalyst

Min-Jae Kim^{a,c}, Jae-Rang Youn^{b,c}, Hye Jeong Kim^b, Myung Won Seo^c,
Doyeon Lee^c, Kang Seok Go^c, Ki Bong Lee^{a,**}, Sang Goo Jeon^{c,*}

^a Department of Chemical and Biological Engineering, Korea University, Anam-ro 145, Seongbuk-gu, Seoul, 02841, Republic of Korea


^b Graduate School of Energy Science and Technology, Chungnam National University, 99 Daehak-ro, Yuseong-gu, Daejeon, 305-764, Republic of Korea

^c Korea Institute of Energy Research, 152 Gajeong-ro, Yuseong-gu, Daejeon, 34129, Republic of Korea

HIGHLIGHTS

- Effect of CeO₂ addition to Ni/γ-Al₂O₃ catalysts on CO₂ methanation is investigated.
- Ni/15Ce/Al₂O₃ showed good selectivity toward CO₂ methanation at low temperatures.
- Physical properties of the catalyst have little influence on CO₂ methanation.
- Oxidation state and chemical properties significantly affect CO₂ methanation.
- CO₂ methanation efficiency is independent on the amount of Ce³⁺.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 8 April 2020

Received in revised form

9 June 2020

Accepted 15 June 2020

Available online 11 August 2020

Keywords:

CO₂ methanation

ABSTRACT

Ce-promoted Ni/Al₂O₃ catalysts with Ce contents of 0, 5, 10, 15, and 20 wt% were investigated for CO₂ methanation. Ni/15Ce/Al₂O₃ showed good selectivity and catalytic performance in CO₂ methanation and remained stable at 350 °C for 80 h with minor fluctuations. Interactions between Ni and the Ce/Al₂O₃ support was characterized using X-ray diffraction, temperature-programmed reduction of H₂, temperature-programmed desorption of CO₂, X-ray photoelectron spectroscopy, Raman spectroscopy, and thermogravimetric analysis. Addition of Ce did not increase the catalytic surface area, which can significantly enhance the heterogeneous catalytic activity. However, XPS analysis showed that the Ce on the Ni/Al₂O₃ catalyst changed the surface electron states of Ni, Ce, and O. Additionally,

* Corresponding author.

** Corresponding author.

E-mail addresses: kibonglee@korea.ac.kr (K.B. Lee), sgjeon@kier.re.kr (S.G. Jeon).

<https://doi.org/10.1016/j.ijhydene.2020.06.144>

0360-3199/© 2020 Hydrogen Energy Publications LLC. Published by Elsevier Ltd. All rights reserved.