




ELSEVIER

Contents lists available at ScienceDirect

Molecular Catalysis

journal homepage: www.elsevier.com/locate/mcat

Review

Catalytic decomposition of N₂O over cobalt based spinel oxides: The role of additives

Min-Jae Kim^{a,b}, Seung-Jae Lee^a, In-Soo Ryu^a, Min-Wook Jeon^{a,c}, Seung-Hyun Moon^a, Hyun-Seog Roh^b, Sang Goo Jeon^{a,*}

^a Climate Change Research Division, Korea Institute of Energy Research, 152 Gajeong-ro, Yuseong-gu, Daejeon 34129, Republic of Korea

^b Department of Environmental Engineering, Yonsei University, 1 Yonseidae-gil, Wonju, Gangwon 220-710, Republic of Korea

^c Department of Chemical Engineering, Chungnam National University, 99 Daehakro, Yuseong-gu, Daejeon, 305-764, Republic of Korea

ARTICLE INFO

Article history:

Received 13 October 2016

Received in revised form 24 May 2017

Accepted 28 May 2017

Available online 10 June 2017

Keywords:

N₂O decomposition

Cobalt spinel oxide

Surface area

Redox ability

ABSTRACT

Catalytic activity at temperature below 400 °C for N₂O decomposition was investigated over catalysts based on cobalt oxide synthesized by a co-precipitation and impregnation method. The characteristics of the catalysts were assessed by XRD, BET, H₂-TPR, O₂-TPD, and XPS techniques, and the catalytic activity for N₂O decomposition was examined in the presence of O₂ and H₂O. In general, it was observed that the catalytic activity was enhanced by the increase of surface area and redox ability of the catalyst. In particular, the N₂O conversion of K/Co₃O₄ was notably higher than that of pure Co₃O₄, while the Co-CeO₂ catalyst slightly improved the activity. The K/Co-CeO₂ catalyst decomposed 100% of N₂O at 375 °C reaction temperature without O₂ and H₂O in the reaction gas stream, and this value was lowered to 95.7% with the presence of O₂ and H₂O.

© 2017 Published by Elsevier B.V.

Contents

1. Introduction	202
2. Experimental	203
2.1. Catalyst preparation	203
2.2. Catalyst characterization	203
2.3. Catalyst activity test	203
3. Results & discussion	203
3.1. Catalytic decomposition of N ₂ O	203
3.2. H ₂ -TPR and O ₂ -TPD	204
3.3. XPS analysis	205
4. Conclusions	206
Acknowledgement	206
Appendix A. Supplementary data	207
References	207

1. Introduction

Nitrous oxide (N₂O) causes environmental problems such as global warming and the ozone layer destruction. Although annual emissions of N₂O are much smaller than those of CO₂, the Global Warming Potential (GWP) of N₂O is 310 times higher than that of CO₂ [1,2]. The concentration of N₂O in the earth's atmosphere

has been consistently increasing due to natural and anthropogenic activity (0.2% per year) [2,3]. One of the promising technologies for abatement of N₂O is catalytic decomposition, and various types of catalysts including noble metals [4–7], pure and mixed oxides [8–11], and ion-exchanged zeolites [12,13] accordingly have been studied. Recently, cobalt catalysts have been recognized as promising candidates for the decomposition of N₂O due to their comparatively high redox ability [14]. The adsorption of N₂O on the active site (Co²⁺) causes a weakening of the N–O bond, releasing N₂ [8]. The unsaturated Co²⁺ ions are oxidized to Co³⁺ by oxygen species adsorbed on the surface of the catalyst. Due to

* Corresponding author.

E-mail address: sgjeon@kier.re.kr (S.G. Jeon).