

linearAlgebra: Scaling and Distance

Ivan Jacob Agaloos Pesigan

Deviation - Mean Centering

The $n \times 1$ vector of deviations from the mean is given by

$$\mathbf{d} = \mathbf{x} - \bar{x} \quad (1)$$

where \mathbf{x} is an $n \times 1$ column vector and \bar{x} is the mean of \mathbf{x} .

```
d(x[, 1])  
  
##           [,1]  
## [1,]  0.24  
## [2,]  0.04  
## [3,] -0.16  
## [4,] -0.26  
## [5,]  0.14  
## [6,]  0.54  
## [7,] -0.26  
## [8,]  0.14  
## [9,] -0.46  
## [10,]  0.04
```

The $n \times k$ matrix of deviations from the mean is given by

$$\mathbf{D} = \mathbf{X} - \mathbf{1}_n \bar{\mathbf{x}} \quad (2)$$

where \mathbf{X} is an $n \times 1$ matrix, $\mathbf{1}_n$ is an $n \times 1$ column vector of ones, and $\bar{\mathbf{x}}$ is the $k \times 1$ mean vector of \mathbf{X} .

```
d(x)
```

##	Sepal.Length	Sepal.Width	Petal.Length	Petal.Width
## 1	0.24	0.19	-0.05	-0.02
## 2	0.04	-0.31	-0.05	-0.02
## 3	-0.16	-0.11	-0.15	-0.02
## 4	-0.26	-0.21	0.05	-0.02
## 5	0.14	0.29	-0.05	-0.02
## 6	0.54	0.59	0.25	0.18
## 7	-0.26	0.09	-0.05	0.08
## 8	0.14	0.09	0.05	-0.02
## 9	-0.46	-0.41	-0.05	-0.02
## 10	0.04	-0.21	0.05	-0.12

Standardized Score - Scaling

The standardized score of the vector \mathbf{x} is given by

$$z = \frac{\mathbf{x} - \bar{\mathbf{x}}}{s} \quad (3)$$

where $\bar{\mathbf{x}}$ is the mean and s is the standard deviation of \mathbf{x} .

```

z(x[, 1])

##           [,1]
## [1,]  0.8237318
## [2,]  0.1372886
## [3,] -0.5491545
## [4,] -0.8923761
## [5,]  0.4805102
## [6,]  1.8533965
## [7,] -0.8923761
## [8,]  0.4805102
## [9,] -1.5788192
## [10,]  0.1372886

```

The standardized score for the j^{th} column of a matrix \mathbf{X} is given by

$$z_j = \frac{\mathbf{x}_j - \bar{x}_j}{s_j} \quad (4)$$

where \mathbf{x}_j is the j^{th} column of a matrix \mathbf{X} , \bar{x}_j is the mean of \mathbf{x}_j , and s_j is the standard deviation of \mathbf{x}_j .

```

z(x)

##      Sepal.Length Sepal.Width Petal.Length Petal.Width
## 1      0.8237318    0.6186158      -0.46291  -0.2535463
## 2      0.1372886   -1.0093205      -0.46291  -0.2535463
## 3     -0.5491545   -0.3581460     -1.38873  -0.2535463
## 4     -0.8923761   -0.6837333       0.46291  -0.2535463
## 5      0.4805102    0.9442031     -0.46291  -0.2535463
## 6      1.8533965    1.9209649       2.31455   2.2819165

```

```
## 7    -0.8923761    0.2930285    -0.46291    1.0141851
## 8      0.4805102    0.2930285     0.46291   -0.2535463
## 9     -1.5788192   -1.3349078    -0.46291   -0.2535463
## 10     0.1372886   -0.6837333     0.46291   -1.5212777
```

Δ^2 - Squared Mahalanobis Distance

The squared Mahalanobis distance of the vector \mathbf{x} is given by

$$\Delta^2 = (\mathbf{x} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \quad (5)$$

where $\boldsymbol{\mu}$ is the mean vector and $\boldsymbol{\Sigma}$ is the covariance matrix.

```
deltacapsq(
  x,
  mu  = colMeans(x),
  sigmacap = stats::cov(x)
)

##           1           2           3           4           5           6           7           8
## 2.1836824  6.0959842  2.3508914  2.6450391  3.3674652  7.2574119  3.7700132  0.7755954
##           9          10
## 3.0556619  4.4982552
```