

Reticular Action Model (RAM) Notation Notes

Ivan Jacob Agaloos Pesigan

2021-01-23

Contents

1	Description	5
2	Reticular Action Model (RAM) Matrix Notation	7
2.1	Full Model	7
2.2	Observed/Manifest/Given Variables vs. Unobserved/Latent/Hidden Variables	8
3	Reticular Action Model (RAM) Path Diagram	11
4	Student's t-test	17
4.1	Symbolic	17
4.2	Numerical Example	21
4.3	Equations to RAM	28
4.4	Equations to Expectations	30

Chapter 1

Description

This is a collection of my personal notes on the Reticular Action Model (RAM) notation that accompanies the **ramR** package (Pesigan, 2021) in the **R** statistical environment (R Core Team, 2020). You can install the released version of **ramR** from GitHub with:

```
remotes::install_github("jeksterslab/ramR")
```

These notes are based on the following resources:

- Boker and McArdle (2005)
- McArdle and McDonald (1984)
- McArdle (2005)

See GitHub Pages for the html deployment.

Chapter 2

Reticular Action Model (RAM) Matrix Notation

2.1 Full Model

Definition 2.1.

$$\mathbf{v} = \mathbf{A}\mathbf{v} + \mathbf{u} \quad (2.1)$$

where

- \mathbf{v} and \mathbf{u} are $t \times 1$ vectors of random variables
- \mathbf{A} is a $t \times t$ matrix of *directed* or *asymmetric* relationship from column variable v_j to row variable v_i
 - \mathbf{A} represent the regression of each of the t variables \mathbf{v} on the other $t - 1$ variables
 - diagonal $a_{i,i}$ is zero
 - u_i represent the residual of v_i
 - if all regression coefficients on other variables are zero, then the variable v_i is considered the same as its own residual u_i

Definition 2.2.

$$\mathbf{S} = \mathbb{E} \{ \mathbf{u}\mathbf{u}' \}, \quad (2.2)$$

where

- \mathbf{S} is a $t \times t$ matrix of *undirected* or *symmetric* relationship
 - the notation Ω is used in other sources for \mathbf{S}
- \mathbb{E} is the expectation operator

Definition 2.3.

$$\mathbf{C} = \mathbb{E} \{ \mathbf{v}\mathbf{v}' \}, \quad (2.3)$$

where

- \mathbf{C} is a $t \times t$ variance-covariance matrix
 - the notation Σ is used in other sources for \mathbf{C}

Definition 2.4.

$$\mathbf{v} = \mathbf{A}\mathbf{v} + \mathbf{u}$$

can be rewritten as

$$\begin{aligned}\mathbf{v} - \mathbf{A}\mathbf{v} &= \mathbf{u} \\ \mathbf{u} &= \mathbf{v} - \mathbf{A}\mathbf{v} \\ \mathbf{u} &= (\mathbf{I} - \mathbf{A})\mathbf{v}\end{aligned}\tag{2.4}$$

assuming that $(\mathbf{I} - \mathbf{A})$ is non-singular,

$$\mathbf{E} = (\mathbf{I} - \mathbf{A})^{-1}\tag{2.5}$$

then

$$\begin{aligned}\mathbf{v} &= (\mathbf{I} - \mathbf{A})^{-1}\mathbf{u} \\ &= \mathbf{E}\mathbf{u}.\end{aligned}\tag{2.6}$$

Using the definitions above, \mathbf{S} and \mathbf{C} are given by

$$\begin{aligned}\mathbf{S} &= (\mathbf{I} - \mathbf{A})\mathbf{C}(\mathbf{I} - \mathbf{A})^{-1} \\ &= \mathbf{E}^{-1}\mathbf{C}(\mathbf{E}^{-1})^{\top}\end{aligned}\tag{2.7}$$

$$\begin{aligned}\mathbf{C} &= (\mathbf{I} - \mathbf{A})^{-1}\mathbf{S}[(\mathbf{I} - \mathbf{A})^{-1}]^{\top} \\ &= \mathbf{E}\mathbf{S}\mathbf{E}^{\top}\end{aligned}\tag{2.8}$$

2.2 Observed/Manifest/Given Variables vs. Unobserved/Latent/Hidden Variables

Definition 2.5.

$$\mathbf{v} = \begin{bmatrix} \mathbf{g}_{p \times 1} \\ \mathbf{h}_{q \times 1} \end{bmatrix}\tag{2.9}$$

$$t = p + q\tag{2.10}$$

- \mathbf{g} may be considered observed, manifest or *given* variables
- \mathbf{h} may be considered unobserved, latent, or *hidden* variables

Definition 2.6.

$$\mathbf{F} = [\mathbf{I}_{p \times p} : \mathbf{0}_{p \times q}]\tag{2.11}$$

- the \mathbf{F} matrix acts as a *filter* to select the manifest variables out of the full set of manifest and latent variables

$$\mathbf{g} = \mathbf{F}\mathbf{v} \quad (2.12)$$

$$\begin{aligned} \mathbf{g} &= \mathbf{F}(\mathbf{I} - \mathbf{A})^{-1} \mathbf{u} \\ &= \mathbf{F}\mathbf{E}\mathbf{u} \end{aligned} \quad (2.13)$$

Definition 2.7.

$$\mathbf{M} = \mathbb{E} \{ \mathbf{g}\mathbf{g}^T \} \quad (2.14)$$

$$\begin{aligned} \mathbf{M} &= \mathbf{F}(\mathbf{I} - \mathbf{A})^{-1} \mathbf{S} [(\mathbf{I} - \mathbf{A})^{-1}]^T \mathbf{F}^T \\ &= \mathbf{F}\mathbf{E}\mathbf{S}\mathbf{E}^T \mathbf{F}^T \\ &= \mathbf{F}\mathbf{C}\mathbf{F}^T \end{aligned} \quad (2.15)$$

- when components of \mathbf{v} are permuted, the columns of \mathbf{F} can be correspondingly permuted
- the rows and columns of \mathbf{C} that are filtered out by \mathbf{F} contain useful information about the latent variable structure.

The equations above completely define RAM.

Chapter 3

Reticular Action Model (RAM) Path Diagram






Figure 3.1: Path Diagram Elements



$$y = \alpha + \beta x + \varepsilon$$

Figure 3.2: Two-Variable Regression Model

Figure 3.3: k -Variable Regression Model




Figure 3.4: Two-Factor Confirmatory Factor Analysis Model




Figure 3.5: Two-Factor Confirmatory Factor Analysis Model with Mean Structure




Figure 3.6: Path Model with Latent Variables

Chapter 4

Student's t -test

In this section, the Student's t -test is presented as a structural equation model using the RAM notation. Let y be a continuous dependent variable, x be a dichotomous independent variable ($x = \{0, 1\}$), and ε be the stochastic error term with mean 0 and constant variance of σ_ε^2 across the values of x . The associations of the variables are given by

$$y = \alpha + \beta x + \varepsilon$$

where

- α is the expected value of y when $x = 0$
- β is the unit change in y for unit change in x
- $\alpha + \beta$ is the expected value of y when $x = 1$




Figure 4.1: Student's t -test

4.1 Symbolic

Let $\{y, x, \varepsilon\}$ be the variables of interest.

$$\mathbf{A} = \begin{pmatrix} 0 & \beta & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\mathbf{S} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \sigma_x^2 & 0 \\ 0 & 0 & \sigma_\varepsilon^2 \end{pmatrix}$$

$$\mathbf{C} = (\mathbf{I} - \mathbf{A})^{-1} \mathbf{S} [(\mathbf{I} - \mathbf{A})^{-1}]^\top$$

$$= \mathbf{E} \mathbf{S} \mathbf{E}^\top$$

$$\begin{aligned} &= \begin{pmatrix} 1 & \beta & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \sigma_x^2 & 0 \\ 0 & 0 & \sigma_\varepsilon^2 \end{pmatrix} \begin{pmatrix} 1 & \beta & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}^\top \\ &= \begin{pmatrix} \sigma_x^2 \beta^2 + \sigma_\varepsilon^2 & \beta \sigma_x^2 & \sigma_\varepsilon^2 \\ \sigma_x^2 \beta & \sigma_x^2 & 0 \\ \sigma_\varepsilon^2 & 0 & \sigma_\varepsilon^2 \end{pmatrix} \end{aligned}$$

$$\mathbf{F} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

$$\mathbf{M} = \mathbf{F} (\mathbf{I} - \mathbf{A})^{-1} \mathbf{S} [(\mathbf{I} - \mathbf{A})^{-1}]^\top \mathbf{F}^\top$$

$$= \mathbf{F} \mathbf{E} \mathbf{S} \mathbf{E}^\top \mathbf{F}^\top$$

$$= \mathbf{F} \mathbf{C} \mathbf{F}^\top$$

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \sigma_x^2 \beta^2 + \sigma_\varepsilon^2 & \beta \sigma_x^2 & \sigma_\varepsilon^2 \\ \sigma_x^2 \beta & \sigma_x^2 & 0 \\ \sigma_\varepsilon^2 & 0 & \sigma_\varepsilon^2 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^\top$$

$$= \begin{pmatrix} \sigma_x^2 \beta^2 + \sigma_\varepsilon^2 & \beta \sigma_x^2 \\ \sigma_x^2 \beta & \sigma_x^2 \end{pmatrix}$$

$$\mathbf{v} = (\mathbf{I} - \mathbf{A})^{-1} \mathbf{u}$$

$$= \left[\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 0 & \beta & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right]^{-1} \begin{pmatrix} \alpha \\ \mu_x \\ 0 \end{pmatrix}$$

$$= \begin{pmatrix} \alpha + \beta \mu_x \\ \mu_x \\ 0 \end{pmatrix}$$

$$\begin{aligned}
\mathbf{u} &= (\mathbf{I} - \mathbf{A}) \mathbf{v} \\
&= \left[\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 0 & \beta & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right] \begin{pmatrix} \alpha + \beta\mu_x \\ \mu_x \\ 0 \end{pmatrix} \\
&= \begin{pmatrix} \alpha \\ \mu_x \\ 0 \end{pmatrix}
\end{aligned}$$

$$\begin{aligned}
\mathbf{g} &= \mathbf{F} (\mathbf{I} - \mathbf{A})^{-1} \mathbf{u} \\
&= \left[\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 0 & \beta & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right]^{-1} \begin{pmatrix} \alpha \\ \mu_x \\ 0 \end{pmatrix} \\
&= \begin{pmatrix} \alpha + \beta\mu_x \\ \mu_x \\ 0 \end{pmatrix}
\end{aligned}$$

4.1.1 Using the ramR Package

A

```
##   y   x       e
## y "0" "beta" "1"
## x "0" "0"    "0"
## e "0" "0"    "0"
```

S

```
##   y   x       e
## y "0" "0"      "0"
## x "0" "sigma[x]^2" "0"
## e "0" "0"      "sigma[varepsilon]^2"
```

u

```
##   u
## y "alpha"
## x "mu[x]"
## e "0"
```

filter

```
##   y x e
## y 1 0 0
## x 0 1 0
```

The covariance expectations can be symbolically derived using the `ramR::C_sym()` function.

```
ramR::C_sym(A, S)
```

```
## {{sigma[x]^2*beta^2+sigma[varepsilon]^2,          beta*sigma[x]^2,
## {          sigma[x]^2*beta,          sigma[x]^2,
## {          sigma[varepsilon]^2,          0,
```

$$\mathbf{C} = \begin{pmatrix} \sigma_x^2 \beta^2 + \sigma_\varepsilon^2 & \beta \sigma_x^2 & \sigma_\varepsilon^2 \\ \sigma_x^2 \beta & \sigma_x^2 & 0 \\ \sigma_\varepsilon^2 & 0 & \sigma_\varepsilon^2 \end{pmatrix}$$

The covariance expectations for the observed variables can be symbolically derived using the `ramR::M_sym()` function.

```
ramR::M_sym(A, S, filter)
```

```
## {{sigma[x]^2*beta^2+sigma[varepsilon]^2,          beta*sigma[x]^2},
## {          sigma[x]^2*beta,          sigma[x]^2}}
```

$$\mathbf{M} = \begin{pmatrix} \sigma_x^2 \beta^2 + \sigma_\varepsilon^2 & \beta \sigma_x^2 \\ \sigma_x^2 \beta & \sigma_x^2 \end{pmatrix}$$

The mean expectations can be symbolically derived using the `ramR::v_sym()` function.

```
ramR::v_sym(A, u)
```

```
## {{alpha+beta*mu[x]},
## {          mu[x]},
## {          0}}
```

$$\mathbf{v} = \begin{pmatrix} \alpha + \beta \mu_x \\ \mu_x \\ 0 \end{pmatrix}$$

The mean expectations for the observed variables can be symbolically derived using the `ramR::g_sym()` function.

```
ramR::g_sym(A, u, filter)
```

```
## {{alpha+beta*mu[x]},
## {          mu[x]}}
```

$$\mathbf{g} = \begin{pmatrix} \alpha + \beta \mu_x \\ \mu_x \end{pmatrix}$$

4.2 Numerical Example

Let \mathbf{df} be a random sample with the following parameters

Parameter	$x = 0$	$x = 1$
Sample Size	500	500
μ	0	1
σ^2	1	1


Parameter	Description	Estimate
α	Mean of $x = 0$.	0
β	Mean of $x = 1$ minus $x = 0$.	1


```
head(df)
```

```
##           y x
## 1  1.3709584 0
## 2 -0.5646982 0
## 3  0.3631284 0
## 4  0.6328626 0
## 5  0.4042683 0
## 6 -0.1061245 0
```

```
summary(df)
```

```
##           y           x
## Min.   :-2.9931  Min.   :0.0
## 1st Qu.: -0.2770  1st Qu.:0.0
## Median :  0.4503  Median :0.5
## Mean    :  0.4742  Mean    :0.5
## 3rd Qu.:  1.2492  3rd Qu.:1.0
## Max.    :  4.4953  Max.    :1.0
```





4.2.1 t-test

```
t.test(y ~ x, data = df)
```

```
##
## Welch Two Sample t-test
##
## data: y by x
## t = -15.897, df = 994.36, p-value < 2.2e-16
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
## -1.1329278 -0.8839594
## sample estimates:
## mean in group 0 mean in group 1
## -0.03004622 0.97839737
```

4.2.2 Linear Regression

```
summary(lm(y ~ x, data = df))
```

```
##
## Call:
## lm(formula = y ~ x, data = df)
##
```

```
## Residuals:
##      Min       1Q   Median       3Q      Max
## -3.3501 -0.6517  0.0086  0.6858  3.5169
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -0.03005    0.04486   -0.67   0.503
## x            1.00844    0.06344   15.90 <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.003 on 998 degrees of freedom
## Multiple R-squared:  0.2021, Adjusted R-squared:  0.2013
## F-statistic: 252.7 on 1 and 998 DF,  p-value: < 2.2e-16
```

4.2.3 Structural Equation Modeling

4.2.3.1 lavaan (Rosseel, 2012)

```
model <- "
  y ~ x
"
fit <- lavaan::sem(
  model,
  data = df,
  meanstructure = TRUE,
  fixed.x = FALSE
)
lavaan::summary(fit)

## lavaan 0.6-7 ended normally after 12 iterations
##
##      Estimator                      ML
##      Optimization method          NLMINB
##      Number of free parameters      5
##
##      Number of observations          1000
##
## Model Test User Model:
##
##      Test statistic                  0.000
##      Degrees of freedom              0
##
## Parameter Estimates:
##
##      Standard errors                Standard
##      Information                    Expected
##      Information saturated (h1) model Structured
##
## Regressions:
##              Estimate Std.Err  z-value  P(>|z|)
##  y ~
```

```
##      x              1.008    0.063   15.913    0.000
##
## Intercepts:
##              Estimate Std.Err  z-value  P(>|z|)
##      .y             -0.030    0.045   -0.671    0.503
##      x              0.500    0.016   31.623    0.000
##
## Variances:
##              Estimate Std.Err  z-value  P(>|z|)
##      .y              1.004    0.045   22.361    0.000
##      x              0.250    0.011   22.361    0.000
```

4.2.3.2 OpenMx (Boker et al., 2020)

RAM matrices can be used to specify models in `OpenMx`. Note, however, that the `u` vector in the RAM notation is `M` in the `OpenMx` notation.

```
mxData <- OpenMx::mxData(
  observed = df,
  type = "raw"
)
mxA <- OpenMx::mxMatrix(
  type = "Full",
  nrow = 3,
  ncol = 3,
  free = c(
    F, T, F,
    F, F, F,
    F, F, F
  ),
  values = c(
    0, 0.20, 1,
    0, 0, 0,
    0, 0, 0
  ),
  labels = c(
    NA, "beta", NA,
    NA, NA, NA,
    NA, NA, NA
  ),
  byrow = TRUE,
  name = "mxA"
)
mxS <- OpenMx::mxMatrix(
  type = "Symm",
  nrow = 3,
  ncol = 3,
  free = c(
    F, F, F,
    F, T, F,
    F, F, T
  ),
  values = c(
```



```

    0, 0, 0,
    0, 0.20, 0,
    0, 0, 0.20
  ),
  labels = c(
    NA, NA, NA,
    NA, "sigma2x", NA,
    NA, NA, "sigma2e"
  ),
  byrow = TRUE,
  name = "mxS"
)
mxM <- OpenMx::mxMatrix(
  type = "Full",
  nrow = 1,
  ncol = 3,
  free = c(
    T, T, F
  ),
  values = c(
    0.20,
    0.20,
    0
  ),
  labels = c(
    "alpha",
    "mux",
    NA
  ),
  byrow = TRUE,
  name = "mxM"
)
mxF <- OpenMx::mxMatrix(
  type = "Full",
  nrow = 2,
  ncol = 3,
  free = FALSE,
  values = c(
    1, 0, 0,
    0, 1, 0
  ),
  byrow = TRUE,
  name = "mxF"
)
expRAM <- OpenMx::mxExpectationRAM(
  A = "mxA",
  S = "mxS",
  F = "mxF",
  M = "mxM",
  dimnames = c(
    "y",
    "x",
    "e"
  )

```

```

)
)
objML <- OpenMx::mxFitFunctionML()
mxMod <- OpenMx::mxModel(
  name = "Model",
  data = mxData,
  matrices = list(
    mxA,
    mxS,
    mxF,
    mxM
  ),
  expectation = expRAM,
  fitfunction = objML
)
fit <- OpenMx::mxRun(mxMod)

```

```
## Running Model with 5 parameters
```

```
summary(fit)
```

```
## Summary of Model
##
## free parameters:
##      name matrix row col   Estimate Std.Error A
## 1   beta    mxA   1   2  1.00844356 0.06337369
## 2 sigma2x    mxS   2   2  0.25000000 0.01118034
## 3 sigma2e    mxS   3   3  1.00402596 0.04490152
## 4   alpha    mxM   1   y -0.03004621 0.04481202
## 5     mux    mxM   1   x  0.49999999 0.01581140
##
## Model Statistics:
##           | Parameters | Degrees of Freedom | Fit (-2lnL units)
##      Model:              5              1995          4293.478
##   Saturated:              5              1995              NA
## Independence:            4              1996              NA
## Number of observations/statistics: 1000/2000
##
## Information Criteria:
##           | df Penalty | Parameters Penalty | Sample-Size Adjusted
## AIC:      303.4776          4303.478          4303.538
## BIC:      -9487.4941          4328.016          4312.136
## CFI: NA
## TLI: 1 (also known as NNFI)
## RMSEA: 0 [95% CI (NA, NA)]
## Prob(RMSEA <= 0.05): NA
## To get additional fit indices, see help(mxRefModels)
## timestamp: 2021-01-23 18:43:07
## Wall clock time: 0.04290342 secs
## optimizer: SLSQP
## OpenMx version number: 2.18.1
## Need help? See help(mxSummary)
```

4.2.4 Using the ramR Package

A

```
##      y      x e
## y 0 1.008444 1
## x 0 0.000000 0
## e 0 0.000000 0
```

S

```
##      y      x      e
## y 0 0.0000000 0.000000
## x 0 0.2502503 0.000000
## e 0 0.0000000 1.006038
```

u

```
##      u
## y -0.03004622
## x 0.50000000
## e 0.00000000
```

filter

```
##      y x e
## y 1 0 0
## x 0 1 0
```

The covariance expectations can be numerically derived using the `ramR::C_num()` function.

```
ramR::C_num(A, S)
```

```
##      y      x      e
## y 1.2605321 0.2523633 1.006038
## x 0.2523633 0.2502503 0.000000
## e 1.0060380 0.0000000 1.006038
```

The covariance expectations for the observed variables can be numerically derived using the `ramR::M_num()` function.

```
ramR::M_num(A, S, filter)
```

```
##      y      x
## y 1.2605321 0.2523633
## x 0.2523633 0.2502503
```

The mean expectations can be numerically derived using the `ramR::v_num()` function.

```
ramR::v_num(A, u)
```

```
##           v
## y 0.4741756
## x 0.5000000
## e 0.0000000
```

The mean expectations for the observed variables can be numerically derived using the `ramR::v_num()` function.

```
ramR::g_num(A, u, filter)
```

```
##           g
## y 0.4741756
## x 0.5000000
```

4.3 Equations to RAM

The `ramR` package has a utility function to convert structural equations to RAM notation. The Student's *t*-test can be expressed in the following equations


```
eq <- "
# VARIABLE1 OPERATION VARIABLE2 LABEL
e          by          y          1
y          on          x          beta
e          with        e          sigma[varepsilon]^2
x          with        x          sigma[x]^2
y          on          1          alpha
x          on          1          mu[x]
"
```

The error term is treated as a latent variable and defined with the operation `by`. It's value is constrained to 1. The regression of y on x is defined by operation `on`. It is labeled as `beta`. The variance of x and the error variance are defined using the operation `with`. These are labeled `sigma[x]^2` and `sigma[varepsilon]^2` respectively. The intercept and the mean of x are defined using the operation `on` 1. These are labeled `alpha` and `mu[x]` respectively.

The `ramR::eq2ram` converts the equations to RAM notation.

```
ramR::eq2ram(eq)
```

```
## $eq
##   var1  op var2          label
## 1    e  by   y           1
## 2    y  on   x          beta
## 3    e with  e sigma[varepsilon]^2
## 4    x with  x    sigma[x]^2
## 5    y  on   1          alpha
## 6    x  on   1          mu[x]
##
```

Figure 4.2: Student's t -test's Structural Equations

```
## $variables
## [1] "y" "x" "e"
##
## $A
##   y   x   e
## y "0" "beta" "1"
## x "0" "0"    "0"
## e "0" "0"    "0"
##
## $S
##   y   x           e
## y "0" "0"         "0"
## x "0" "sigma[x]^2" "0"
## e "0" "0"         "sigma[varepsilon]^2"
##
## $filter
##   y x e
## y 1 0 0
## x 0 1 0
##
## $u
##   u
## y "alpha"
## x "mu[x]"
## e "0"
```

4.4 Equations to Expectations

The `ramR` package has a utility function to convert structural equations to expectations both symbolically and numerically.

```
eq <- "
# VARIABLE1 OPERATION VARIABLE2 LABEL
e      by      y      1
y      on      x      beta
e      with    e      sigma[varepsilon]^2
x      with    x      sigma[x]^2
y      on      1      alpha
x      on      1      mu[x]
"
```

```
ramR::eq2exp_sym(eq)
```

```
## $variables
## [1] "y" "x" "e"
##
## $A
## {{ 0, beta, 1},
## { 0, 0, 0},
## { 0, 0, 0}}
##
## $S
## {{ 0, 0, 0},
## { 0, sigma[x]^2, 0},
## { 0, 0, sigma[varepsilon]^2}}
##
## $u
## {{alpha},
## {mu[x]},
## { 0}}
##
## $filter
## {{1, 0, 0},
## {0, 1, 0}}
##
## $v
## {{alpha+beta*mu[x]},
## { mu[x]},
## { 0}}
##
## $g
## {{alpha+beta*mu[x]},
## { mu[x]}}
##
## $C
## {{sigma[x]^2*beta^2+sigma[varepsilon]^2, beta*sigma[x]^2,
## { sigma[x]^2*beta, sigma[x]^2,
## { sigma[varepsilon]^2, 0,
##
```

```
## $M
## {{sigma[x]^2*beta^2+sigma[varepsilon]^2,          beta*sigma[x]^2},
## {          sigma[x]^2*beta,                      sigma[x]^2}}
```

```
eq <- "
# VARIABLE1 OPERATION VARIABLE2 VALUE
e          by          y          1.00
y          on          x          1.00
e          with        e          1.00
x          with        x          0.25
y          on          1          0.00
x          on          1          0.50
"
```

```
ramR::eq2exp_num(eq)
```

```
## $variables
## [1] "y" "x" "e"
##
## $A
##   y x e
## y 0 1 1
## x 0 0 0
## e 0 0 0
##
## $S
##   y   x e
## y 0 0.00 0
## x 0 0.25 0
## e 0 0.00 1
##
## $u
##   u
## y 0.0
## x 0.5
## e 0.0
##
## $filter
##   y x e
## y 1 0 0
## x 0 1 0
##
## $v
##   v
## y 0.5
## x 0.5
## e 0.0
##
## $g
##   g
## y 0.5
## x 0.5
##
```

```
## $C
##      y      x e
## y 1.25 0.25 1
## x 0.25 0.25 0
## e 1.00 0.00 1
##
## $M
##      y      x
## y 1.25 0.25
## x 0.25 0.25
```


Bibliography

- Boker, S. M. and McArdle, J. J. (2005). Path analysis and path diagrams. In Everitt, B. S. and Howell, D. C., editors, *Encyclopedia of Statistics in Behavioral Science*, pages 1529–1531. John Wiley & Sons, Ltd, Chichester, UK.
- Boker, S. M., Neale, M. C., Maes, H. H., Wilde, M. J., Spiegel, M., Brick, T. R., Estabrook, R., Bates, T. C., Mehta, P., von Oertzen, T., Gore, R. J., Hunter, M. D., Hackett, D. C., Karch, J., Brandmaier, A. M., Pritikin, J. N., Zahery, M., Kirkpatrick, R. M., Wang, Y., Goodrich, B., Driver, C., Massachusetts Institute of Technology, Johnson, S. G., Association for Computing Machinery, Kraft, D., Wilhelm, S., Medland, S., Falk, C. F., Keller, M., Manjunath B G, The Regents of the University of California, Ingber, L., Shao Voon, W., Palacios, J., Yang, J., Guennebaud, G., and Niesen, J. (2020). *OpenMx 2.18.1 User Guide*.
- McArdle, J. J. (2005). The development of the RAM rules for latent variable structural equation modeling. In Maydeu-Olivares, A. and McArdle, J. J., editors, *Contemporary psychometrics: A festschrift for Rodrick P. McDonald*, Multivariate applications book series, pages 225–273. Lawrence Erlbaum Associates, Mahwah, NJ.
- McArdle, J. J. and McDonald, R. P. (1984). Some algebraic properties of the reticular action model for moment structures. *British Journal of Mathematical and Statistical Psychology*, 37(2):234–251.
- Pesigan, I. J. A. (2021). *ramR: Reticular Action Model (RAM) Notation*. R package version 0.9.0.
- R Core Team (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48(2):1–36.