

**REFER NOW
SUSPECTED SEVERE MALARIA****PATIENT**

Age: 2 years
State / LGA: Kano

Sex: —

PRESENTING FINDINGS

- Fever for 3 day(s)
- Malaria RDT: positive
- **Danger sign: Convulsions**
- **Danger sign: Unable to drink or breastfeed**

TRIAGE DECISION**REFER NOW**

The patient exhibits convulsions and inability to drink or breastfeed, which are IMCI general danger signs indicating very severe disease requiring urgent referral.

SUSPECTED CONDITION

1. Severe malaria (*high*)
2. Lassa fever (*low*)
3. Severe bacterial infection (sepsis) (*moderate*)

ADVISED ACTIONS

1. Refer now: danger signs present (Convulsions, Unable to drink or breastfeed).
2. Put patient in a holding area and institute infection prevention measures
3. Alert the relevant authorities: infectious disease team or responsible physician
4. Start rehydration. Commence Fluid therapy (see guidance on fluid replacement in Chapter 4)
5. Monitor vital signs every 4 hours (Pulse Rate, Blood Pressure, Respiratory Rate, Temperature)
6. Give first dose of artesunate or quinine for severe malaria Give first dose of an appropriate antibiotic
7. Treat the child to prevent low blood sugar

OUTBREAK CONTEXT

- Lassa Fever — Ondo (live report, 2026-09-06)
- Lassa Fever — Bauchi (live report, 2026-09-06)
- Lassa Fever — Taraba (live report, 2026-09-06)
- Lassa Fever — Edo (live report, 2026-09-06)
- Lassa Fever — Benue (live report, 2026-09-06)
- Diphtheria — Kano (live report, 2026-07-12 (older report))
- Cholera — Nigeria (live report, date unknown)
- Cerebrospinal meningitis — Kano (NCDC Cerebrospinal Meningitis Situation Report, Week 14, 2025 (cumulative Epi week 40 2024 - 14 2025), p.1 (older surveillance))

TO BE COMPLETED BY HEALTH WORKER

Referred to (facility): _____

Referred by (name and cadre): _____

Date: _____ Time: _____ Phone: _____

Notes: _____

Decision support only, not a diagnosis. A qualified health worker must confirm every decision. Doses come from the guideline table only.

Generated by Ibà — decision support only. Not a diagnosis.