

REFER NOW
SUSPECTED SEVERE MALARIA

PATIENT	
Age: 2 years State / LGA: Kano	Sex: —

PRESENTING FINDINGS
- Fever for 3 day(s) - Malaria RDT: positive Danger sign: Convulsions Danger sign: Unable to drink or breastfeed

TRIAGE DECISION
REFER NOW Patient exhibits convulsions and unable to drink or breastfeed, which are general danger signs requiring urgent referral per IMCI guidelines.

SUSPECTED CONDITION
- Severe malaria (<i>high</i>) - Cerebrospinal meningitis (<i>moderate</i>) - Sepsis/bacterial infection (<i>low</i>)

ADVISED ACTIONS
- Refer now: danger signs present (Convulsions, Unable to drink or breastfeed). - Put patient in a holding area and institute infection prevention measures - Give first dose of artesunate or quinine for severe malaria - Give first dose of an appropriate antibiotic - Treat to prevent low blood sugar

OUTBREAK CONTEXT
- Lassa Fever — Ondo (live report, 2026-09-06) - Lassa Fever — Bauchi (live report, 2026-09-06) - Lassa Fever — Taraba (live report, 2026-09-06) - Lassa Fever — Edo (live report, 2026-09-06) - Lassa Fever — Benue (live report, 2026-09-06) - Diphtheria — Kano (live report, 2026-07-12 (older report)) - Cholera — Nigeria (live report, date unknown) - Cerebrospinal meningitis — Kano (NCDC Cerebrospinal Meningitis Situation Report, Week 14, 2025 (cumulative Epi week 40 2024 - 14 2025), p.1 (older surveillance))

TO BE COMPLETED BY HEALTH WORKER

Referred to (facility): _____

Referred by (name and cadre): _____

Date: _____ Time: _____ Phone: _____

Notes: _____

Decision support only, not a diagnosis. A qualified health worker must confirm every decision. Doses come from the guideline table only.

Generated by Ibà — decision support only. Not a diagnosis.