


Whitening & Anti-Wrinkle & Anti-Oxidation

The Most Stabilized Vitamin C derivative

* Trade Name : V2G (INCI Name : Ascorbyl Glucoside)


(CAS No. : 129499-78-1)

Introduction



V2G is a conjugate consisting of one molecule of glucose linked at the C-2 hydroxyl group of vitamin C. The C-2 hydroxyl group is the primary site of natural vitamin C degradation. V2G is manufactured by transferring a glucose molecule from starch to vitamin C, using an enzyme, glycosyltransferase, obtained from a microorganism.


Characteristic 1 :
Release of Vitamin C by α-Glucosidase




Specification

Appearance	Fine powder
Colour	White to cream
Odour	Odourless
Assay	≥ 98.0%
Melting Point	158.0 ~ 163.0℃
Specific optical rotation [α]D 20	+186.0 ~ + 188.0˚
Loss on Drying	≤ 1.0%
Residue on Ignition	≤ 0.2%
pH	2.3 ~ 2.4
Free Ascorbic Acid	≤ 0.1%
Free Glucose	≤ 0.1%

Characteristic 2 : High Stability



Stability in solution



Stability with Cu^{2+}

MacroCare Tech Co.Ltd

Tel +82-43-214-3655~6
Fax +82-43-214-3658
E-mail Sales Team isales@macrocare.net msupport@macrocare.net
Address 32, Gangni-1gil, Ochang-eup, Cheongwon-gu, Cheongju-si, Chungcheongbuk-do, 28126, S.KOREA