

Interpretable Machine Learning

Conditional Feature Importance (CFI)




Figure: Bike Sharing Dataset

Figure: Bike Sharing Dataset

Learning goals

- Extrapolation and Conditional Sampling
- Conditional Feature Importance (CFI)
- Interpretation of CFI and difference to PFI

Learning goals

- Extrapolation and Conditional Sampling
- Conditional Feature Importance (CFI)
- Interpretation of CFI and difference to PFI

CONDITIONAL FEATURE IMPORTANCE IDEA

- **Permutation Feature Importance Idea:** Replace the feat. of interest x_j with an indep. sample from the marginal dist. $P(x_j)$, e.g. by randomly perm. obs. in x_j .




CONDITIONAL FEATURE IMPORTANCE IDEA

- **Permutation Feature Importance Idea:** Replace the feat. of interest x_j with an indep. sample from the marginal dist. $P(x_j)$, e.g. by randomly perm. obs. in x_j
- **Problem:** Under dependent features, permutation leads to extrapolation
- **Problem:** Under dependent features, permutation leads to extrapolation




CONDITIONAL FEATURE IMPORTANCE IDEA

- **Permutation Feature Importance Idea:** Replace the feat. of interest x_j with an indep. sample from the marginal dist. $P(x_j)$, e.g. by randomly perm. obs. in x_j
- **Problem:** Under dependent features, permutation leads to extrapolation
- **Conditional Feature Importance Idea:** Resample x_j from the cond. dist. $P(x_j|x_{-j})$, s.t. the joint dist. is preserved, i.e., $P(x_j|x_{-j})P(x_{-j}) = P(x_j, x_{-j})$
- **Conditional Feature Importance Idea:** Resample x_j from the conditional distribution $P(x_j|x_{-j})$, such that the joint distribution is preserved, i.e., $P(x_j|x_{-j})P(x_{-j}) = P(x_j, x_{-j})$



CONDITIONAL FEATURE IMPORTANCE IDEA




- **Permutation Feature Importance Idea:** Replace the feat. of interest x_j with an indep. sample from the marginal dist. $P(x_j)$, e.g. by randomly perm. obs. in x_j
- **Problem:** Under dependent features, permutation leads to extrapolation
- **Conditional Feature Importance Idea:** Resample x_j from the cond. dist. $P(x_j|x_{-j})$, s.t. the joint dist. is preserved, i.e., $P(x_j|x_{-j})P(x_{-j}) = P(x_j, x_{-j})$

Example: Conditional permutation scheme

Example: Conditional permutation scheme

► Molnar et al (2020)



- $X_2 \sim U(0, 1)$ and $X_1 \sim N(0, 1)$ if $X_2 < 0.5$, else $X_1 \sim N(4, 4)$ (black dots)

- **Left:** For $X_2 < 0.5$, permuting X_1 (crosses) preserves marginal (but not joint) distribution


~ Bottom: Marginal density of X_1

- **Right:** Permuting X_1 within subgroups $X_2 < 0.5$ & $X_2 \geq 0.5$ reduces extrapolation

~ Bottom: Density of X_1 conditional on groups

RECALL: EXTRAPOLATION IN PFI

Example: Let $y = x_3 + \epsilon_3$ with $\epsilon_3 \sim N(0, 0.1)$ where $x_1 = \epsilon_1$, $x_2 = x_1 + \epsilon_2$ are highly correlated ($\epsilon_1 \sim N(0, 1)$, $\epsilon_2 \sim N(0, 0.01)$) and $x_3 = \epsilon_3$ with $\epsilon_3 \sim N(0, 1)$. All noise terms are independent. Fitting a LM yields $\hat{f}(x) \approx 0.3x_1 - 0.3x_2 + x_3$.



Hexbin plot of x_1, x_2 before permuting x_1 (left), after permuting x_1 (center), and PFI scores (right) $\Rightarrow x_1$ and x_2 should be irrelevant for the prediction $\hat{f}(x)$ for

$\{x : P(x) > 0\}$ as $0.3x_1 - 0.3x_2 \approx 0$
 \Rightarrow PFI evaluates model on unrealistic obs. outside $P(x) \sim x_1$ and x_2 are considered relevant

Hooker et al. (2021)

Conditional feature importance (CFI) for features x_S using test data \mathcal{D} :

Conditional feature importance (CFI) for features x_S using test data \mathcal{D} :


- Measure the error with unperturbed features.
- Measure the error with perturbed feature values $\tilde{x}^{S|-S}$, where $\tilde{x}^{S|-S} \sim \mathbb{P}(x_S | x_{-S})$.

- Repeat permuting the feature (e.g., m times) and average the difference of both errors:

- Repeat permuting the feature (e.g., m times) and average the difference of both errors:

Here, $\tilde{\mathcal{D}}^{S|-S}$ denotes the dataset where features x_S where sampled conditional on the remaining features x_{-S} .


Here, $\tilde{\mathcal{D}}^{S|-S}$ denotes the dataset where features x_S where sampled conditional on the remaining features x_{-S} .




IMPLICATIONS OF CFI

König et al. (2020)

Interpretation: Due to the conditional sampling w.r.t. all other features, CFI quantifies a feature's unique contribution to the model performance.



Interpretation: Due to the conditional sampling w.r.t. all other features, CFI quantifies a feature's unique contribution to the model performance.



Entanglement with data:

- If feature x_S does not contribute unique information about y , i.e. $x_S \perp\!\!\!\perp y | x_{-S} \Rightarrow \text{CFI} = 0$
- $\text{CFI} = 0$ Why? Under the conditional independence $P(\tilde{x}^{S|-S}, y) = P(x, y)$
- Why? Under the conditional independence $P(\tilde{x}^{S|-S}, y) = P(x, y)$
 - \leadsto no prediction-relevant information is destroyed by permutation of x_S conditional on x_{-S}

Interpretation: Due to the conditional sampling w.r.t. all other features, CFI quantifies a feature's unique contribution to the model performance.



Entanglement with data:

- If feature x_S does not contribute unique information about y , i.e. $x_S \perp\!\!\!\perp y | x_{-S} \Rightarrow \text{CFI} = 0$
- $\text{CFI} = 0$ Why? Under the conditional independence $P(\tilde{x}^{SI-S}, y) = P(x, y)$
- Why? Under the conditional independence $P(\tilde{x}^{SI-S}, y) = P(x, y)$
no prediction-relevant information is destroyed by permutation of x_S conditional on x_{-S}

Entanglement with model:

- If the model does not use a feature $\Rightarrow \text{CFI} = 0$

Entanglement with model:

- Why? Then the prediction is not affected by any perturbation of the feature
- If the model does not use a feature $\Rightarrow \text{CFI} = 0$
model performance does not change after conditional permutation
- Why? Then the prediction is not affected by any perturbation of the feature
model performance does not change after conditional permutation

IMPLICATIONS OF CFI

Can we gain insight into whether ...

1 the feature x_j is causal for the prediction?

- $CFI_j \neq 0 \Rightarrow$ model relies on x_j (converse does not hold, see next slide)



IMPLICATIONS OF CFI

Can we gain insight into whether ...

1 the feature x_j is causal for the prediction?

- $CFI_j \neq 0 \Rightarrow$ model relies on x_j (converse does not hold, see next slide)

2 the variable x_j contains prediction-relevant information?

- If $x_j \perp\!\!\!\perp y$ but $x_j \perp\!\!\!\perp y | x_{-j}$ (e.g. x_j and x_{-j} share information) $\Rightarrow CFI_j \neq 0$
- x_j is not exploited by model (regardless of whether it is useful for y or not) $\Rightarrow CFI_j = 0$



IMPLICATIONS OF CFI

Can we gain insight into whether ...



❶ the feature x_j is causal for the prediction?

- $CFI_j \neq 0 \Rightarrow$ model relies on x_j (converse does not hold, see next slide)

❷ the variable x_j contains prediction-relevant information?

- If $x_j \perp\!\!\!\perp y$ but $x_j \not\perp\!\!\!\perp y | x_{-j}$ (e.g. x_j and x_{-j} share information) $\Rightarrow CFI_j \neq 0$
- x_j is not exploited by model (regardless of whether it is useful for y or not) $\Rightarrow CFI_j = 0$

❸ Does the model require access to x_j to achieve it's prediction performance?

❹ Does the model require access to x_j to achieve its prediction performance?

- $CFI_j \neq 0 \Rightarrow x_j$ contributes unique information (meaning $x_j \neq y(x_{-j})$)
- Only uncovers the relationships that were exploited by the model

COMPARISON: PFI AND CFI

Example: Let $y = x_3 + \epsilon_3$ with $\epsilon_3 \sim N(0, 0.1)$ where $x_1 = \epsilon_1$, $x_2 = x_1 + \epsilon_2$ are highly correlated ($\epsilon_1 \sim N(0, 1)$, $\epsilon_2 \sim N(0, 0.01)$) and $x_3 = \epsilon_3$ with $\epsilon_3 \sim N(0, 1)$. All noise terms are independent. Fitting a LM yields $\hat{f}(x) \approx 0.3x_1 - 0.3x_2 + x_3$.




Figure: Density plot for x_1, x_2 before permuting x_1 (left). PFI and CFI (right).

Figure: Density plot for x_1, x_2 before permuting x_1 (left). PFI and CFI (right).

- $\Rightarrow x_1$ and x_2 are irrelevant for the prediction $\hat{f}(x)$ for $\{x : P(x) > 0\}$ as $0.3x_1 - 0.3x_2 \approx 0$
- \Rightarrow PFI evaluates model on unrealistic obs. outside $P(x) \leadsto x_1, x_2$ are considered relevant (PFI > 0)
- \Rightarrow Since x_1 can be reconstructed from x_2 and vice versa, CFI considers x_1 and x_2 to be irrelevant
- \Rightarrow Since x_1 can be reconstructed from x_2 and vice versa, CFI considers x_1 and x_2 to be irrelevant