


Optimization in Machine Learning

Mathematical Concepts: Conditions for optimality





Learning goals

- Local and global
- First & second order conditions

DEFINITION LOCAL AND GLOBAL MINIMUM

Given $\mathcal{S} \subseteq \mathbb{R}^d$, $f : \mathcal{S} \rightarrow \mathbb{R}$:

- f has **global minimum** in $\mathbf{x}^* \in \mathcal{S}$, if $f(\mathbf{x}^*) \leq f(\mathbf{x})$ for all $\mathbf{x} \in \mathcal{S}$
- f has a **local minimum** in \mathbf{x}^* , if $f(\mathbf{x}^*) \leq f(\mathbf{x})$ for all $\mathbf{x} \in B_\epsilon(\mathbf{x}^*)$, with $B_\epsilon(\mathbf{x}^*) := \{\mathbf{x} \in \mathcal{S} \mid \|\mathbf{x} - \mathbf{x}^*\| < \epsilon\}$ (“ ϵ ”-ball round \mathbf{x}^*).



Source (left): https://en.wikipedia.org/wiki/Maxima_and_minima.

Source (right): <https://wngaw.github.io/linear-regression/>.


EXISTENCE OF OPTIMA

$$f : \mathcal{S} \rightarrow \mathbb{R}$$

- f continuous:
 - A real-valued function f defined on a **compact set** must attain a minimum and a maximum (extreme value theorem).
- f not continuous:
 - In general no statement possible about existence of maximum/minimum.

FIRST ORDER CONDITION FOR OPTIMALITY

Let $f \in \mathcal{C}^1$. **Observation:** At a local minimum (for an interior point) 1st order Taylor series approx is perfectly flat; 1st order derivs are 0.



(Strictly) convex functions (left: univariate; right: multivariate) with unique local minimum, which is the global one. Tangent (hyperplane) is perfectly flat at the optimum.

Source: Watt, 2020, Machine Learning Refined.

FIRST ORDER CONDITION FOR OPTIMALITY

At every (interior) local minimum \mathbf{x}^* the first derivative is necessarily always zero; it is therefore called **first-order** or **necessary** condition.

- **First-order condition (univariate):** Let $\mathbf{x}^* \in \mathbb{R}$ be a local minimum of f . Then:

$$f'(\mathbf{x}^*) = 0$$


- **First-order condition (multivariate):** Let $\mathbf{x}^* \in \mathbb{R}^d$ be a local minimum of f . Then:

$$\nabla f(\mathbf{x}^*) = (0, 0, \dots, 0)^\top$$

The points at which the first order derivative is zero are called **stationary points**.

FIRST ORDER CONDITION FOR OPTIMALITY

The condition is **not sufficient**: Not every stationary point ($\nabla f(\mathbf{x}) = 0$) is a local minimum.



Left: Four points fulfill the necessary conditions; but two of the points are local maxima (not minima). Middle: One point fulfills the necessary condition, but is not a local optimum. Right: Multiple local minima and maxima.

Source: Watt, 2020, Machine Learning Refined.

SECOND ORDER CONDITION FOR OPTIMALITY

Let $f \in \mathcal{C}^2$. If the function is locally convex, so:

- **Second-order condition (univariate):** A **stationary** point $x^* \in \mathcal{S} \subseteq \mathbb{R}$ fulfills

$$f''(x^*) > 0.$$

- **Second-order condition (multivariate):** A **stationary** point $\mathbf{x}^* \in \mathcal{S} \subseteq \mathbb{R}^d$ fulfills

$$\nabla^2 f(\mathbf{x}^*) \text{ is positive definite}$$

(all EVs positive), hence curvature is positive in all directions.

Then the second-order condition is **sufficient** to prove a local minimum.

CONDITIONS FOR OPTIMALITY AND CONVEXITY

Let $f : \mathcal{S} \rightarrow \mathbb{R}$ be convex on convex set \mathcal{S} . Then the following holds:

- Any local minimum is also global minimum
- If f strictly convex, f has exactly one local minimum which is also unique global minimum on \mathcal{S}

CONDITIONS FOR OPTIMALITY AND CONVEXITY

Def.: **Saddle point**

- all gradients of x are zero but still, x is no optimum
- Hessian is indefinite

