

# Linear algebra for AI & ML

September - 15

- [ Fundamentals of Matrix  
Computations  
D. Watkins

- [ Matrix computations  
Golub, Van Loan

$$Ax = b \quad ; \quad A \in \mathbb{R}^{n \times n}, \quad b \in \mathbb{R}^n$$

find  $x$  s.t.  $Ax = b$



- sensitivity analysis:

$$A(x + \delta x) = b + \delta b$$

Allowing perturbation  
in  $b$  only.

$$\frac{\|\delta x\|_2}{\|x\|_2} \leq \underbrace{\|A\|_2 \|A^{-1}\|_2}_{\|k_2(A)\|} \frac{\|\delta b\|_2}{\|b\|_2}$$

Geometrically,  $k_2(A)$ .



$$\text{maxmag}(A) = \max_{\|x\|_2=1} \|Ax\|_2$$

$$\text{minmag}(A) = \min_{\|x\|_2=1} \|Ax\|_2$$

orthogonal matrices :  $\text{maxmag}(Q) = 1$  ;  $\text{minmag}(Q) = 1$   
 unit circle  $\rightarrow$  unit circle

---

$$k_2(A) = \frac{\text{maxmag}(A)}{\text{minmag}(A)} \quad \leftarrow \text{want to prove!!}$$

$$k_2(A) = \|A\|_2 \|A^{-1}\|_2 \quad \leftarrow \text{we know!!}$$

Lemma:  $A$  is non-singular matrix.

$$\text{maxmag}(A) = \frac{1}{\text{minmag}(A^T)}$$

and  $\boxed{\text{maxmag}(A^{-1}) = \frac{1}{\text{minmag}(A)}}$

Pf:  $A$  is nonsingular (invertible)

$$\Rightarrow Ax = 0 \Leftrightarrow x = 0 \quad \checkmark$$

Also,  $A$  is invertible  $\Rightarrow A$  is one-one.  $\left. \vphantom{\begin{matrix} A \\ A \end{matrix}} \right\}$

$$Ax = y \Rightarrow x = A^{-1}y$$

$$\text{maxmag}(A) = \max_{x \neq 0} \frac{\|Ax\|_2}{\|x\|_2}$$

$$= \max_{y \neq 0} \frac{\|y\|_2}{\|A^{-1}y\|_2}$$

$$= \frac{1}{\min_{y \neq 0} \frac{\|A^{-1}y\|_2}{\|y\|_2}}$$

$$= \frac{1}{\text{minmag}(A^{-1})}$$

$$k_2(A) = \|A\|_2 \|A^{-1}\|_2$$

$$= \text{maxmag}(A) \cdot \text{maxmag}(A^{-1})$$

$$k_2(A) = \frac{\text{maxmag}(A)}{\text{minmag}(A)}$$

Q:

$$Ax = b$$

$$A(x + \delta x) = b + \delta b$$

If  $\text{cond}(A)$  is very large. (ill-conditioned matrix)

$\frac{\|\delta b\|_2}{\|b\|_2}$  is small.



$$\frac{\|\delta x\|_2}{\|x\|_2}$$

may be small.  
may be large.

$$\cancel{Ax} + A\delta x = \cancel{b} + \delta b$$

$$A\delta x = \delta b$$

$$\Rightarrow \boxed{\delta x = A^{-1} \delta b} \quad \checkmark$$



$$\textcircled{Ax} = b$$

$$\textcircled{X} \beta = Y$$

$$\textcircled{\beta} = \underbrace{(X^T X)^{-1}}_{\text{wavy line}} X^T Y$$

Another interpretation of ill-conditioned matrices:

$$A \in \mathbb{R}^{n \times n}$$

Let  $A$  be NOT invertible.

$\Rightarrow$  columns of  $A$  are linearly dependent.

$\Rightarrow \exists x \in \mathbb{R}^n ; x \neq 0$  such that

$$Ax = 0$$

Without loss of generality, assume  $\|x\|_2 = 1$

$$\Rightarrow \underline{\min \text{mag}(A) = 0}$$

$$\min \text{mag}(A) = \min_{\|x\|_2=1} \|Ax\|_2$$

Let  $A$  be matrix which is ill-conditioned.

$$k_2(A) \gg 1$$

$$\frac{\max_{\text{mag}}(A)}{\min_{\text{mag}}(A)} \gg 1$$

WLOG

assume

that

$$\max_{\text{mag}}(A) = \|A\|_2 = 1$$

$$k_2(A) \gg 1$$

$$\Rightarrow \frac{1}{\min_{\text{mag}}(A)} \gg 1$$

$$\Rightarrow \min_{\text{mag}}(A) \ll 1$$

$$\frac{1}{0.001}$$



there exists a vector  $x \in \mathbb{R}^n$ ;  $\|x\|_2 = 1$

s.t.  $\|Ax\|_2$  is "very very small".

$$\frac{\|Ax\|_2}{\|x\|_2} = \epsilon$$

$\|Ax\|_2$  is "almost zero".

$\Rightarrow Ax$  is "almost zero".

$\Rightarrow$  columns of  $A$  are "almost linearly dependent".

$\Rightarrow$  matrix  $A$  is "almost non-invertible"  
"almost singular".

$$\|Qx\|_2 = \|x\|_2 \quad \forall x \in \mathbb{R}^n$$

$$\max_{x \neq 0} \frac{\|Qx\|_2}{\|x\|_2} = 1$$

$$A = \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix}$$

$$\alpha > 0, \beta > 0$$

$$\alpha, \beta \in \underline{(0, 1)}$$

