

On the properties $\alpha - z$ Rényi divergences on general von Neumann algebras

Fumio Hiai and Anna Jenčová

February 22, 2024

1 Introduction

2 Preliminaries

2.1 Basic definitions

Let \mathcal{M} be a von Neumann algebra and let \mathcal{M}^+ be the cone of positive elements in \mathcal{M} . We denote the predual by \mathcal{M}_* , its positive part by \mathcal{M}_*^+ and the set of normal states by $\mathfrak{S}_*(\mathcal{M})$. For $\psi \in \mathcal{M}_*^+$, we will denote by $s(\psi)$ the support projection of ψ .

For $0 < p \leq \infty$, let $L_p(\mathcal{M})$ be the Haagerup L_p -space over \mathcal{M} and let $L_p(\mathcal{M})^+$ its positive cone, [?]. We will use the identifications $\mathcal{M} \simeq L_\infty(\mathcal{M})$, $\mathcal{M}_* \ni \psi \leftrightarrow h_\psi \in L_1(\mathcal{M})$ and the notation $\text{Tr } h_\psi = \psi(1)$ for the trace in $L_1(\mathcal{M})$. In this way, \mathcal{M}_*^+ is identified with the positive cone $L_1(\mathcal{M})^+$ and $\mathfrak{S}_*(\mathcal{M})$ with subset of elements in $L_1(\mathcal{M})^+$ with unit trace. Precise definitions and further details on the spaces $L_p(\mathcal{M})$ can be found in the notes [?].

2.2 The $\alpha - z$ -Rényi divergences

In [2?], the $\alpha - z$ -Rényi divergence for $\psi, \varphi \in \mathcal{M}_*^+$ was defined as follows:

Definition 1. Let $\psi, \varphi \in \mathcal{M}_*^+$, $\psi \neq 0$ and let $\alpha, z > 0$, $\alpha \neq 1$. The $\alpha - z$ -Rényi divergence is defined as

$$D_{\alpha,z}(\psi \parallel \varphi) := \frac{1}{\alpha - 1} \log \frac{Q_{\alpha,z}(\psi \parallel \varphi)}{\psi(1)},$$

where

$$Q_{\alpha,z} = \begin{cases} \text{Tr} \left(h_\varphi^{(1-\alpha)/2z} h_\psi^{\alpha/z} h_\varphi^{(1-\alpha)/2z} \right)^z, & \text{if } 0 < \alpha < 1 \\ \|x\|_z^z, & \text{if } \alpha > 1 \text{ and} \\ h_\psi^{\alpha/z} = h_\varphi^{(\alpha-1)/2z} x h_\varphi^{(\alpha-1)/2z}, \text{ with } x \in s(\varphi)L_z(\mathcal{M})s(\varphi) \\ \infty & \text{otherwise.} \end{cases}$$

In the case $\alpha > 1$, the following alternative form will be useful.

Lemma 1. [2] Let $\alpha > 1$ and $\psi, \varphi \in \mathcal{M}_*^+$. Then $Q_{\alpha,z}(\psi\|\varphi) < \infty$ if and only if there is some $y \in L_{2z}(\mathcal{M})s(\varphi)$ such that

$$h_\psi^{\alpha/2z} = y h_\varphi^{(\alpha-1)/2z}.$$

Moreover, in this case, such y is unique and we have $Q_{\alpha,z}(\psi\|\varphi) = \|y\|_{2z}^{2z}$.

The standard Rényi divergence [?] is contained in this range as $D_\alpha(\psi\|\varphi) = D_{\alpha,1}(\psi\|\varphi)$. The sandwiched Rényi divergence is obtained as $\tilde{D}_\alpha(\psi\|\varphi) = D_{\alpha,\alpha}(\psi\|\varphi)$, see [1?] for some alternative definitions and properties of \tilde{D}_α . The definition in [?] and [1] is based on the Kosaki interpolation spaces $L_p(\mathcal{M}, \varphi)$ with respect to a state [?]. These spaces and complex interpolation method will be used frequently also in the present work.

Many of the properties of $D_{\alpha,z}(\psi\|\varphi)$ were extended from the finite dimensional case in [2]. In particular, the following variational expressions will be an important tool for our work.

Theorem 1. Let $\psi, \varphi \in \mathcal{M}_*^+$, $\psi \neq 0$.

(i) Let $0 < \alpha < 1$ and $\max\{\alpha, 1 - \alpha\} \leq z$. Then

$$Q_{\alpha,z}(\psi\|\varphi) = \inf_{a \in \mathcal{M}^{++}} \left\{ \alpha \operatorname{Tr} \left((a^{1/2} h_\psi^{\alpha/z} a^{1/2})^{z/\alpha} \right) + (1 - \alpha) \operatorname{Tr} \left((a^{-1/2} h_\varphi^{(1-\alpha)/2z} a^{-1/2})^{z/(1-\alpha)} \right) \right\}.$$

Moreover, if $\lambda^{-1}\varphi \leq \psi \leq \lambda\varphi$ for some $\lambda > 0$, then the infimum is attained at some $a_0 \in \mathcal{M}^{++}$ satisfying

$$h_\varphi^{(1-\alpha)/2z} a_0^{-1} h_\varphi^{(1-\alpha)/2z} = \left(h_\varphi^{(1-\alpha)/2z} h_\psi^{\alpha/z} h_\varphi^{(1-\alpha)/2z} \right)^{1-\alpha}$$

and

$$\operatorname{Tr} \left((h_\psi^{\alpha/2z} a_0 h_\psi^{\alpha/2z})^{z/\alpha} \right) = \operatorname{Tr} \left((h_\varphi^{(1-\alpha)/2z} h_\psi^{\alpha/z} h_\varphi^{(1-\alpha)/2z})^z \right).$$

(ii) Let $1 < \alpha \leq 2z$, then

$$Q_{\alpha,z}(\psi\|\varphi) = \sup_{a \in \mathcal{M}_+} \left\{ \alpha \operatorname{Tr} \left((a^{1/2} h_\psi^{\alpha/z} a^{1/2})^{z/\alpha} \right) - (\alpha - 1) \operatorname{Tr} \left((a^{1/2} h_\varphi^{(\alpha-1)/2z} a^{1/2})^{z/(\alpha-1)} \right) \right\}.$$

Proof. For part (i) see [2, Theorem 1 (vi)] and its proof. The inequality \geq in part (ii) holds for all α and z and was proved in [2, Theorem 2 (vi)]. We now prove the opposite inequality.

Assume first that $Q_{\alpha,z}(\psi\|\varphi) < \infty$, so that there is some $x \in s(\varphi)L_z(\mathcal{M})^+s(\varphi)$ such that $h_\psi^{\alpha/z} = h_\varphi^{(\alpha-1)/2z} x h_\varphi^{(\alpha-1)/2z}$. Plugging this into the right hand side, we obtain

$$\begin{aligned} & \sup_{a \in \mathcal{M}_+} \left\{ \alpha \operatorname{Tr} \left((a^{1/2} h_\psi^{\alpha/z} a^{1/2})^{z/\alpha} \right) - (\alpha - 1) \operatorname{Tr} \left((a^{1/2} h_\varphi^{(\alpha-1)/2z} a^{1/2})^{z/(\alpha-1)} \right) \right\} \\ &= \sup_{a \in \mathcal{M}_+} \left\{ \alpha \operatorname{Tr} \left((a^{1/2} h_\varphi^{(\alpha-1)/2z} x h_\varphi^{(\alpha-1)/2z} a^{1/2})^{z/\alpha} \right) - (\alpha - 1) \operatorname{Tr} \left((a^{1/2} h_\varphi^{(\alpha-1)/2z} a^{1/2})^{z/(\alpha-1)} \right) \right\} \\ &= \sup_{a \in \mathcal{M}_+} \left\{ \alpha \operatorname{Tr} \left((x^{1/2} h_\varphi^{(\alpha-1)/2z} a h_\varphi^{(\alpha-1)/2z} x^{1/2})^{z/\alpha} \right) - (\alpha - 1) \operatorname{Tr} \left((h_\varphi^{(\alpha-1)/2z} a h_\varphi^{(\alpha-1)/2z})^{z/(\alpha-1)} \right) \right\} \\ &= \sup_{w \in L_{z/(\alpha-1)}(\mathcal{M})^+} \left\{ \alpha \operatorname{Tr} \left((x^{1/2} w x^{1/2})^{z/\alpha} \right) - (\alpha - 1) \operatorname{Tr} \left(w^{z/(\alpha-1)} \right) \right\}, \end{aligned}$$

where we used the fact that $\operatorname{Tr}((a^*a)^p) = \operatorname{Tr}((aa^*)^p)$ for $p > 0$ and $a \in L_{p/2}(\mathcal{M})$ and the fact that the set of elements of the form $h_\varphi^{(\alpha-1)/2z} a h_\varphi^{(\alpha-1)/2z}$ with $a \in \mathcal{M}^+$ is dense in the positive cone $L_{z/(\alpha-1)}(\mathcal{M})^+$. Putting $w = x^{\alpha-1}$ we get

$$\sup_{w \in L_{z/(\alpha-1)}(\mathcal{M})^+} \left\{ \alpha \operatorname{Tr} \left((x^{1/2} w x^{1/2})^{z/\alpha} \right) - (\alpha - 1) \operatorname{Tr} \left(w^{z/(\alpha-1)} \right) \right\} \geq \operatorname{Tr}(x^z) = \|x\|_z^z = Q_{\alpha,z}(\psi\|\varphi).$$

This finishes the proof in the case that $Q_{\alpha,z}(\psi\|\varphi) < \infty$. Note that this holds if $\psi \leq \lambda\varphi$ for some $\lambda > 0$. Indeed, since $\alpha/2z \in (0, 1]$ by the assumption, we then have

$$h_\psi^{\alpha/2z} \leq \lambda^{\alpha/2z} h_\varphi^{\alpha/2z},$$

hence by [?, Lemma A.58] there is some $b \in \mathcal{M}$ such that

$$h_\psi^{\alpha/2z} = b h_\varphi^{\alpha/2z} = y h_\varphi^{(\alpha-1)/2z},$$

where $y = b h_\varphi^{1/2z} \in L_{2z}(\mathcal{M})$. By Lemma 1 we get $Q_{\alpha,z}(\psi\|\varphi) = \|y\|_{2z}^{2z} < \infty$.

In the general case, note that lower semicontinuity [2], we have

$$Q_{\alpha,z}(\psi\|\varphi) \leq \liminf_{\epsilon \searrow 0} Q_{\alpha,z}(\psi\|\varphi + \epsilon\psi)$$

and by the previous paragraph, the variational expression holds for $Q_{\alpha,z}(\psi\|\varphi)$ for all $\epsilon > 0$. The proof is finished by using norm continuity of the map $L_1(\mathcal{M})^+ \ni h \mapsto h^{1/p} \in L_p(\mathcal{M})^+$ for $p > 1$.

I am not entirely sure about this proof, since the convergence of the expressions

$$\text{Tr} \left((a^{1/2} (h_\varphi + \epsilon h_\psi)^{(\alpha-1)/2z} a^{1/2})^{z/(\alpha-1)} \right) \rightarrow \text{Tr} \left((a^{1/2} h_\varphi^{(\alpha-1)/2z} a^{1/2})^{z/(\alpha-1)} \right)$$

also depends on $\|a\|$, which is not bounded. But probably I misinterpreted something, or I am just being stupid.

□

3 Data processing inequality and reversibility of quantum channels

Let $\gamma : \mathcal{N} \rightarrow \mathcal{M}$ be a normal positive unital map. Then the predual of γ defines a positive linear map $\gamma_* : L_1(\mathcal{M}) \rightarrow L_1(\mathcal{N})$ that preserves the trace. The support of γ will be denoted by $s(\gamma)$, recall that this is the largest projection $p \in \mathcal{N}$ such that $\gamma(p) = 1$. For any $\rho \in \mathcal{M}_*^+$ we clearly have $s(\rho \circ \gamma) \leq s(\gamma)$, with equality if ρ is faithful. It follows that γ_* maps $L_1(\mathcal{M})$ to $s(\gamma)L_1(\mathcal{N})s(\gamma) \equiv L_1(s(\gamma)\mathcal{N}s(\gamma))$. For any $\rho \in \mathcal{M}_*^+$, $\rho \neq 0$, the map

$$s(\gamma)\mathcal{N}s(\gamma) \rightarrow s(\rho)\mathcal{M}s(\rho), \quad a \mapsto s(\rho)\gamma(a)s(\rho)$$

is a faithful normal positive unital map, so using such restrictions we may always assume that both ρ and $\rho \circ \gamma$ are faithful.

The Petz dual of γ with respect to a faithful $\rho \in \mathcal{M}_*^+$ is a map $\gamma_\rho^* : \mathcal{M} \rightarrow \mathcal{N}$, introduced in [? ?]. It was proved that it is again normal, positive and unital, in addition, it is n -positive whenever γ is. As explained in [?] γ_ρ^* is determined by the equality

$$(\gamma_\rho^*)_*(h_{\rho \circ \gamma}^{1/2} b h_{\rho \circ \gamma}^{1/2}) = h_\rho^{1/2} \gamma(b) h_\rho^{1/2}, \tag{1}$$

for all $b \in \mathcal{N}^+$, here $(\gamma_\rho^*)_*$ is the predual map of γ_ρ^* . We also have

$$(\gamma_\rho^*)_*(h_{\rho \circ \gamma}) = (\gamma_\rho^*)_* \circ \gamma_*(h_\rho) = h_\rho.$$

3.1 Data processing inequality

In this paragraph we prove the data processing inequality (DPI) for $D_{\alpha,z}$ with respect to normal positive unital maps. In the case of the sandwiched divergences \tilde{D}_α with $1/2 \leq \alpha \neq 1$, DPI was proved in [1?], see also [?] for an alternative proof in the case when the maps are also completely positive.

Lemma 2. *Let $\gamma : \mathcal{N} \rightarrow \mathcal{M}$ be a normal positive unital map and let $\rho \in \mathcal{M}_*^+$, $b \in \mathcal{N}^+$.*

(i) *If $p \in [1/2, 1)$, then*

$$\|h_{\rho \circ \gamma}^{\frac{1}{2p}} b h_{\rho \circ \gamma}^{\frac{1}{2p}}\|_p \leq \|h_\rho^{\frac{1}{2p}} \gamma(b) h_\rho^{\frac{1}{2p}}\|_p.$$

(ii) *If $p \in [1, \infty]$, the inequality reverses.*

Proof. Let us denote $\beta := \gamma_\rho^*$ and let $\omega \in \mathcal{M}_*^+$ be such that $h_\omega := h_{\rho \circ \gamma}^{1/2} b h_{\rho \circ \gamma}^{1/2} \in L_1(\mathcal{N})^+$. Then β is a normal positive unital map and we have

$$\beta_*(h_\omega) = h_\rho^{1/2} \gamma(b) h_\rho^{1/2}, \quad \beta_*(h_{\rho \circ \gamma}) = h_\rho.$$

Let $p \in [1/2, 1)$, then

$$\begin{aligned} \|h_\rho^{\frac{1}{2p}} \gamma(b) h_\rho^{\frac{1}{2p}}\|_p^p &= \|h_\rho^{\frac{1-p}{2p}} \beta_*(h_\omega) h_\rho^{\frac{1-p}{2p}}\|_p^p = Q_{p,p}(\beta_*(h_\omega) \|h_\rho\|) = Q_{p,p}(\beta_*(h_\omega) \|\beta_*(h_{\rho \circ \gamma})\|) \\ &\geq Q_{p,p}(h_\omega \|h_{\rho \circ \gamma}\|) = \|h_{\rho \circ \gamma}^{\frac{1-p}{2p}} h_\omega h_{\rho \circ \gamma}^{\frac{1-p}{2p}}\|_p^p = \|h_{\rho \circ \gamma}^{\frac{1}{2p}} b h_{\rho \circ \gamma}^{\frac{1}{2p}}\|_p^p. \end{aligned}$$

Here we have used the DPI for the sandwiched Rényi divergence $D_{\alpha,\alpha}$ for $\alpha \in [1/2, 1)$, [1, Theorem 4.1]. This proves (i). The case (ii) was proved in [2] (see Eq. (22) therein), using the relation of the sandwiched Rényi divergence to the Kosaki L_p norms as

$$\begin{aligned} \|h_\rho^{1/2p} \gamma(b) h_\rho^{1/2p}\|_p^p &= Q_{p,p}(h_\rho^{1/2} \gamma(b) h_\rho^{1/2} \|h_\rho\|) = Q_{p,p}(\beta_*(h_\omega) \|\beta_*(h_{\rho \circ \gamma})\|) \\ &\leq Q_{p,p}(h_\omega \|h_{\rho \circ \gamma}\|) = \|h_{\rho \circ \gamma}^{\frac{1}{2p}} b h_{\rho \circ \gamma}^{\frac{1}{2p}}\|_p^p, \end{aligned}$$

here the inequality follows from the DPI for sandwiched Rényi divergence $D_{p,p}$ with $p > 1$, [?]. \square

Theorem 2 (DPI). *Let $\psi, \varphi \in \mathcal{M}_*^+$, $\psi \neq 0$ and let $\gamma : \mathcal{N} \rightarrow \mathcal{M}$ be a normal positive unital map. Assume either of the following conditions:*

(i) $0 < \alpha < 1$, $\max\{\alpha, 1 - \alpha\} \leq z$

(ii) $\alpha > 1$, $\max\{\alpha/2, \alpha - 1\} \leq z \leq \alpha$.

Then we have

$$D_{\alpha,z}(\psi \circ \gamma \| \varphi \circ \gamma) \leq D_{\alpha,z}(\psi \| \varphi).$$

Proof. Under the conditions (i), the DPI was proved in [2, Theorem 1 (viii)]. There an additional assumption was used, namely that \mathcal{N} is σ -finite, to construct a faithful map out of γ . This can be done using the restriction described above, so the additional condition is not needed.

We next prove that DPI holds under the condition (ii). \square

Let ψ be a faithful normal state on a von Neumann algebra \mathcal{M} . We will prove the following inequality:

$$\|h_{\psi \circ \gamma}^{\frac{1}{2p}} b h_{\psi \circ \gamma}^{\frac{1}{2p}}\|_p \leq \|h_{\psi}^{\frac{1}{2p}} \gamma(b) h_{\psi}^{\frac{1}{2p}}\|_p \quad (2)$$

for all $p \in [1/2, 1]$, all $b \in \mathcal{N}^+$ and any unital positive map $\gamma : \mathcal{N} \rightarrow \mathcal{M}$ (Eq. (19) in [2]). This then implies DPI for the $\alpha - z$ -Rényi divergence for $\alpha/2, \alpha - 1 \leq z \leq \alpha$.

Let γ_{ψ}^* be the Petz dual of γ with respect to ψ , then its predual satisfies

$$(\gamma_{\psi}^*)_*(h_{\psi \circ \gamma}^{1/2} b h_{\psi \circ \gamma}^{1/2}) = h_{\psi}^{1/2} \gamma(b) h_{\psi}^{1/2}$$

(this is eq. (21) in [2]). Put $h_{\omega} := h_{\psi \circ \gamma}^{1/2} b h_{\psi \circ \gamma}^{1/2} \in L_1(\mathcal{N})^+$. We then have, using Thm. 4.1 in [1]

$$\begin{aligned} \|h_{\psi}^{\frac{1}{2p}} \gamma(b) h_{\psi}^{\frac{1}{2p}}\|_p^p &= \|h_{\psi}^{\frac{1-p}{2p}} (\gamma_{\psi}^*)_*(h_{\omega}) h_{\psi}^{\frac{1-p}{2p}}\|_p^p = \tilde{Q}_p((\gamma_{\psi}^*)_*(h_{\omega}) \|(\gamma_{\psi}^*)_*(h_{\psi \circ \gamma})\|) \\ &\geq \tilde{Q}_p(h_{\omega} \|h_{\psi \circ \gamma}\|) = \|h_{\psi \circ \gamma}^{\frac{1-p}{2p}} h_{\omega} h_{\psi \circ \gamma}^{\frac{1-p}{2p}}\|_p^p = \|h_{\psi \circ \gamma}^{\frac{1}{2p}} b h_{\psi \circ \gamma}^{\frac{1}{2p}}\|_p^p. \end{aligned}$$

References

- [1] A. Jenčová, Rényi relative entropies and noncommutative L_p -spaces II, Ann. Henri Poincaré 22, 3235–3254 (2021)
- [2] Shinya Kato, On $\alpha - z$ -Rényi divergence in the von Neumann algebra setting, arXiv:2311.01748.