

# The testing range of quantum statistical models and measurements

Michele Dall'Arno\* and Francesco Buscemi†

## Abstract

A positive operator-valued measure (i.e., a quantum measurement) can be regarded as a linear map from the space of states to the space of probability distributions. The image of the set of states through such a map is naturally defined in parametric form, that is, as a body in the space of probability distributions parameterized by the set of states itself. In such a form, the image of a quantum measurement is impractical to treat analytically. Our first result is to provide an implicit outer approximation of the image of any given quantum measurement in any finite dimension: namely, a region in probability space that contains the desired image, but is defined implicitly, using a formula that depends only on the given quantum measurement. This generalizes the bounding recently provided in Ref. [22] by Xu, Schwonnek, and Winter: first, the extension is from Pauli strings to arbitrary measurements; second, the optimization is not restricted to the radius of fixed-axis ellipsoids, but it is a *global* optimization over all the parameters of the ellipsoid. The outer approximation that we construct is *minimal* among all such outer approximations, and *close*, in the sense that it becomes the *maximal inner* approximation up to a constant scaling factor. We also obtain a similar result for the dual problem of implicitizing the image of the set of effects through a family of quantum states (i.e., a quantum statistical model). Finally, we apply our approximation formulas to characterize, in a semi-device independent way, the ability to transform one quantum measurement into another, or one quantum statistical model into another.

---

\*Department of Computer Science and Engineering, Toyohashi University of Technology, Japan, and Yukawa Institute for Theoretical Physics, Kyoto University, Japan

†Graduate School of Informatics, Nagoya University, Chikusa-ku, 464-8601 Nagoya, Japan

In statistics, information theory, and mathematical economics one is often faced with the problem of comparing two setups in terms of their expected performances on a particular task of interest. For example, one might compare two statistical models by comparing their informativeness in a given parameter estimation problem, or two noisy channels with respect to a given communication figure of merit, or again two portfolios with respect to their expected utility in a given betting scenario. The comparison could also be extended, so to ask when a given setup is *always* better than another one, i.e., independent of any particular task at hand. Such “global” comparisons, generally described by a preorder relation, play a crucial role in the formulation of mathematical statistics.

The simplest example of one such preorder in statistics is given by the *majorization preorder* of probability distributions [1, 2, 3, 4]. Generalizing this, we find the comparison of families comprising two or more probability distributions. The case of pairs of probability distributions (i.e., *dichotomies*) is also known as *relative majorization* [5, 6, 7, 8], whereas the case of multiple elements is usually referred to as comparison of statistical *experiments* or *models* [5, 6, 7, 9].

The relevance of such preorder relations is epitomized by Blackwell’s theorem [5, 6], which establishes the equivalence between the above mentioned statistical comparisons, and the existence of a suitable stochastic map that transforms one setup (the “always better” one) into the other (the “always worse” one). For this reason, Blackwell’s theorem and its variants provide a powerful framework for general resource theories [10], and indeed recent quantum extensions of Blackwell’s theorem [11, 12, 13] have found fruitful application in the study of quantum entanglement [14], quantum thermodynamics [15, 16], and quantum measurement theory [17, 18, 19], for example.

Unfortunately, due to the non-commutativity of the underlying algebra, the quantum version of Blackwell’s theorem [11] turns out to be necessarily more convoluted than its original classical variant. This is particularly evident in the case of relative majorization: while classical relative majorization can be summarized in a finite collection of easily computable inequalities [5, 8], in the quantum case (with the notable exceptions of qubits [20, 21]) an infinite number of scalar inequalities must be evaluated [13]. The situation becomes even more cumbersome in the case of quantum statistical models [11].

The image  $\pi(\mathbb{S}_d)$  of the set  $\mathbb{S}_d$  of states through a measurement  $\pi$  is by definition given in parametric form, that is, it is a body in the probability space parameterized by states in the state space. Ideally, one would aim at implicitizing it, that is, writing it in the form  $f(p) \leq 1$ , for probability distributions  $p$ . However, due to intractability of the structure of the state

space, we provide here inclusion conditions in terms of implicit bodies.

**Definition 1.** For any  $d$ -dimensional,  $n$ -outcome measurement  $\boldsymbol{\pi} = \{\pi_i\}_{i=1}^n$ , let  $\{\mathcal{E}_r(\boldsymbol{\pi})\}_{r \in \mathbb{R}}$  be the following family of hyper-ellipsoids parameterized by  $r$ :

$$\mathcal{E}_r(\boldsymbol{\pi}) := \left\{ \mathbf{p} \in \boldsymbol{\pi}(\mathbb{C}^d) \text{ s.t. } \left| \sqrt{Q^+}(\mathbf{p} - \mathbf{t}) \right|_2^2 \leq \frac{1}{r^2} \right\},$$

where  $Q \in \mathbb{R}^{n \times n}$  is the positive semi-definite covariance matrix given by

$$Q_{ij} = \frac{d-1}{d} \left( \text{Tr}[\pi_i \pi_j] - \frac{\text{Tr}[\pi_i] \text{Tr}[\pi_j]}{d} \right), \quad 0 \leq i, j \leq n,$$

and  $\mathbf{t} \in \mathbb{R}^n$  is the vector with entries  $t_i = \text{Tr}[\pi_i]/d$ .

**Theorem 1.** For any  $d$ -dimensional,  $n$ -outcome informationally complete measurement  $\boldsymbol{\pi}$ , one has that  $\mathcal{E}_{\sqrt{d^2-1}}(\boldsymbol{\pi})$  is the maximum volume ellipsoid enclosed in  $\boldsymbol{\pi}(\mathbb{S}_d)$  and  $\mathcal{E}_1(\boldsymbol{\pi})$  is the minimum volume ellipsoid enclosing  $\boldsymbol{\pi}(\mathbb{S}_d)$ .

This generalizes the bounding recently provided in Ref. [22] by Xu, Schwonek, and Winter: first, the extension is from Pauli strings to arbitrary measurements; second, the optimization is not restricted to the radius of fixed-axis ellipsoids, but it is a *global* optimization over all the parameters of the ellipsoid.

Now that we have a close approximation of the image of the set of states through any given measurement, we turn our attention to applying it to semi-device independent tests of simulability. We say that a  $d_1$ -dimensional,  $n$ -outcome measurement  $\boldsymbol{\pi}_1$  simulates a  $d_0$ -dimensional,  $n$ -outcome measurement  $\boldsymbol{\pi}_0$  if and only if there exists a completely positive map  $\mathcal{C} : \mathcal{L}(\mathbb{C}^{d_0}) \rightarrow \mathcal{L}(\mathbb{C}^{d_1})$  such that

$$\boldsymbol{\pi}_1 \circ \mathcal{C} = \boldsymbol{\pi}_0. \quad (1)$$

The following corollary generalizes Corollary 2 of Ref. [21] to the arbitrary dimensional case, providing a semi-device independent test of Eq. (1).

**Corollary 1** (Semi-device independent simulability test). *Given a set  $\mathcal{P}$  of  $n$ -element probability distributions generated by a  $d_1$ -dimensional (otherwise unspecified) measurement  $\boldsymbol{\pi}_1$ , for any  $d_0$  and for any  $d_0$ -dimensional  $n$ -outcome measurement  $\boldsymbol{\pi}_0$  such that*

$$\mathcal{E}_1(\boldsymbol{\pi}_0) \subseteq \mathcal{P},$$

*there exists a trace preserving map  $\mathcal{C}$  that is positive on the support of  $\boldsymbol{\pi}_0$  such that Eq. (1) holds. Moreover, if  $D = 2$ ,  $n \leq 3$ , and  $d \leq 3$ , map  $\mathcal{C}$  in Eq. (1) is completely positive, that is, measurement  $\boldsymbol{\pi}_1$  simulates measurement  $\boldsymbol{\pi}_0$ .*

We also obtain a similar result for the dual problem of implicitizing the image of the set of effects through a family of quantum states (i.e., a quantum statistical model).

## References

- [1] Max O. Lorenz. Methods of measuring the concentration of wealth. *Publ. Am. Stat. Assoc.*, 9(70):209–219, 1905.
- [2] Godfrey Harold Hardy, John Edensor Littlewood, and George Polya. *Inequalities*. Cambridge university press, 1952.
- [3] Barry C. Arnold. *Majorization and the Lorenz Order: A Brief Introduction*. Springer New York, 1987. doi:10.1007/978-1-4615-7379-1.
- [4] Albert W. Marshall, Ingram Olkin, and Barry C. Arnold. *Inequalities: theory of majorization and its applications*. Springer, 2010.
- [5] David Blackwell. Equivalent Comparisons of Experiments. *The Annals of Mathematical Statistics*, 24(2):265–272, 1953. URL: <http://www.jstor.org/stable/2236332>, doi:10.1214/aoms/1177729032.
- [6] D.A. Blackwell and M.A. Girshick. *Theory of Games and Statistical Decisions*. Dover Books on Mathematics. Dover Publications, 1979. URL: <https://books.google.co.jp/books?id=fwTLAgAAQBAJ>.
- [7] Erik Torgersen. *Comparison of Statistical Experiments*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1991. doi:10.1017/CB09780511666353.
- [8] Joseph M. Renes. Relative submajorization and its use in quantum resource theories. *Journal of Mathematical Physics*, 57(12):122202, 12 2016. arXiv:[https://pubs.aip.org/aip/jmp/article-pdf/doi/10.1063/1.4972295/16083406/122202\\_1\\_online.pdf](https://pubs.aip.org/aip/jmp/article-pdf/doi/10.1063/1.4972295/16083406/122202_1_online.pdf), doi:10.1063/1.4972295.
- [9] F. Liese and K.-J. Miescke. *Statistical Decision Theory*. Springer, 2008.
- [10] Eric Chitambar and Gilad Gour. Quantum resource theories. *Rev. Mod. Phys.*, 91:025001, Apr 2019. URL: <https://link.aps.org/doi/10.1103/RevModPhys.91.025001>, doi:10.1103/RevModPhys.91.025001.

- [11] Francesco Buscemi. Comparison of quantum statistical models: Equivalent conditions for sufficiency. *Communications in Mathematical Physics*, 310(3):625–647, 2012. doi:10.1007/s00220-012-1421-3.
- [12] Anna Jencova. Comparison of quantum channels and statistical experiments, 2015. URL: <https://arxiv.org/abs/1512.07016>, doi:10.48550/ARXIV.1512.07016.
- [13] Francesco Buscemi and Gilad Gour. Quantum relative lorenz curves. *Phys. Rev. A*, 95:012110, Jan 2017. URL: <https://link.aps.org/doi/10.1103/PhysRevA.95.012110>, doi:10.1103/PhysRevA.95.012110.
- [14] Francesco Buscemi. All entangled quantum states are nonlocal. *Phys. Rev. Lett.*, 108:200401, May 2012. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.108.200401>, doi:10.1103/PhysRevLett.108.200401.
- [15] Francesco Buscemi. Fully quantum second-law-like statements from the theory of statistical comparisons, 2015. URL: <https://arxiv.org/abs/1505.00535>, doi:10.48550/ARXIV.1505.00535.
- [16] Gilad Gour, David Jennings, Francesco Buscemi, Runyao Duan, and Iman Marvian. Quantum majorization and a complete set of entropic conditions for quantum thermodynamics. *Nature Communications*, 9(1):5352, 2018. doi:10.1038/s41467-018-06261-7.
- [17] Francesco Buscemi, Eric Chitambar, and Wenbin Zhou. Complete resource theory of quantum incompatibility as quantum programmability. *Phys. Rev. Lett.*, 124:120401, Mar 2020. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.124.120401>, doi:10.1103/PhysRevLett.124.120401.
- [18] Francesco Buscemi, Kodai Kobayashi, Shintaro Minagawa, Paolo Perinotti, and Alessandro Tosini. Unifying different notions of quantum incompatibility into a strict hierarchy of resource theories of communication, 2022. URL: <https://arxiv.org/abs/2211.09226>, doi:10.48550/ARXIV.2211.09226.
- [19] Francesco Buscemi, Kodai Kobayashi, and Shintaro Minagawa. A complete and operational resource theory of measurement sharpness, 2023. arXiv:2303.07737.

- [20] Peter M. Alberti and Armin Uhlmann. *Stochasticity and Partial Order Doubly Stochastic Maps and Unitary Mixing*. Springer Netherlands, 1982.
- [21] Michele Dall’Arno, Francesco Buscemi, and Valerio Scarani. Extension of the Alberti-Uhlmann criterion beyond qubit dichotomies. *Quantum*, 4:233, February 2020. doi:10.22331/q-2020-02-20-233.
- [22] Zhen-Peng Xu, René Schwonnek, and Andreas Winter. Bounding the joint numerical range of pauli strings by graph parameters, 2023. arXiv:2308.00753.