

On the properties $\alpha - z$ Rényi divergences on general von Neumann algebras

Fumio Hiai and Anna Jenčová

February 20, 2024

1 Introduction

2 Preliminaries

2.1 Basic definitions

Let \mathcal{M} be a von Neumann algebra and let \mathcal{M}^+ be the cone of positive elements in \mathcal{M} . We denote the predual by \mathcal{M}_* , its positive part by \mathcal{M}_*^+ and the set of normal states by $\mathfrak{S}_*(\mathcal{M})$. For $\psi \in \mathcal{M}_*^+$, we will denote by $s(\psi)$ the support projection of ψ .

For $0 < p \leq \infty$, let $L_p(\mathcal{M})$ be the Haagerup L_p -space over \mathcal{M} and let $L_p(\mathcal{M})$ its positive cone, [?]. We will use the identifications $\mathcal{M} \simeq L_\infty(\mathcal{M})$, $\mathcal{M}_* \ni \psi \leftrightarrow h_\psi \in L_1(\mathcal{M})$ and the notation $\text{Tr } h_\psi = \psi(1)$ for the trace in $L_1(\mathcal{M})$. In this way, \mathcal{M}_*^+ is identified with the positive cone $L_1(\mathcal{M})^+$ and $\mathfrak{S}_*(\mathcal{M})$ with subset of elements in $L_1(\mathcal{M})^+$ with unit trace. Precise definitions and further details on the spaces $L_p(\mathcal{M})$ can be found in the notes [?].

2.2 The $\alpha - z$ -Rényi divergences

In [? ?], the $\alpha - z$ -Rényi divergence for $\psi, \varphi \in \mathcal{M}_*^+$ was defined as follows:

Definition 1. Let $\psi, \varphi \in \mathcal{M}_*^+$, $\psi \neq 0$ and let $\alpha, z > 0$, $\alpha \neq 1$. The $\alpha - z$ -Rényi divergence is defined as

$$D_{\alpha,z}(\psi \parallel \varphi) := \frac{1}{\alpha - 1} \log \frac{Q_{\alpha,z}(\psi \parallel \varphi)}{\psi(1)},$$

where

$$Q_{\alpha,z} = \begin{cases} \text{Tr} \left(h_\varphi^{(1-\alpha)/2z} h_\psi^{\alpha/z} h_\varphi^{(1-\alpha)/2z} \right)^z, & \text{if } 0 < \alpha < 1 \\ \|x\|_z^z, & \text{if } \alpha > 1 \text{ and} \\ h_\psi^{\alpha/z} = h_\varphi^{(\alpha-1)/2z} x h_\varphi^{(\alpha-1)/2z}, \text{ with } x \in s(\varphi)L_z(\mathcal{M})s(\varphi) \\ \infty & \text{otherwise.} \end{cases}$$

In the case $\alpha > 1$, the following alternative form will be useful.

Lemma 1. [?] Let $\alpha > 1$ and $\psi, \varphi \in \mathcal{M}_*^+$. Then $Q_{\alpha,z}(\psi\|\varphi) < \infty$ if and only if there is some $y \in L_{2z}(\mathcal{M})s(\varphi)$ such that

$$h_{\psi}^{\alpha/2z} = y h_{\varphi}^{(\alpha-1)/2z}.$$

Moreover, in this case, such y is unique and we have $Q_{\alpha,z}(\psi\|\varphi) = \|y\|_{2z}^{2z}$.

The standard Rényi divergence [?] is contained in this range as $D_{\alpha}(\psi\|\varphi) = D_{\alpha,1}(\psi\|\varphi)$. The sandwiched Rényi divergence is obtained as $\tilde{D}_{\alpha}(\psi\|\varphi) = D_{\alpha,\alpha}(\psi\|\varphi)$, see [?] for some alternative definitions and properties of \tilde{D}_{α} .

Many of the properties of $D_{\alpha,z}(\psi\|\varphi)$ were extended from the finite dimensional case in [?]. In particular, the following variational formulas will be an important tool for our work.

Theorem 1. (i) $\alpha < 1$, + attained

(ii) $\alpha > 1$

Proof. [?] for (i).

□

3 Data processing inequality and reversibility of quantum channels