

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/268799105>

Stochasticity and partial order. Doubly stochastic maps and unitary mixing

Article · January 1982

CITATIONS

105

READS

103

2 authors:



[Peter M. Alberti](#)

University of Leipzig

50 PUBLICATIONS 431 CITATIONS

[SEE PROFILE](#)



[Armin Uhlmann](#)

University of Leipzig

86 PUBLICATIONS 3,453 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



FRAP - Framework for biodiversity Reconciliation Action Plans [View project](#)



Geometry of state spaces [View project](#)

Mathematics and Its Applications

Managing Editor:

M. HAZEWINKEL

Department of Mathematics, Erasmus University, Rotterdam, The Netherlands

Editorial Board:

R. W. BROCKETT, *Harvard University, Cambridge, Mass., U.S.A.*

J. CORONES, *Iowa State University, Ames, Iowa, U.S.A. and Ames Laboratory,
U.S. Dept. of Energy, Iowa, U.S.A.*

Yu. I. MANIN, *Steklov Institute of Mathematics, Moscow, U.S.S.R.*

G.-C. ROTA, *M.I.T., Cambridge, Mass., U.S.A.*

Volume 9

Peter M. Alberti and Armin Uhlmann

Department of Physics, Karl-Marx-University, Leipzig

Stochasticity and Partial Order

Doubly Stochastic Maps and Unitary Mixing



VEB Deutscher Verlag der Wissenschaften, Berlin

D. Reidel Publishing Company

Dordrecht : Holland / Boston : U.S.A. / London : England

Contents

Introduction	7
1. Some classical results	9
1.1. Elementary notations	9
1.2. The linear space \mathbb{R}^n in n dimensions	9
1.3. Stochastic and doubly stochastic maps	11
1.4. The relation \succ	16
1.5. An example: Comparison of Gibbsian states	19
1.6. An example: Localization of spectra for Hermitian matrices	20
1.7. Convexity and the relation \succ	23
1.8. Examples: Some maps and processes	28
1.9. A partial order of m -tuples a_1, \dots, a_m	31
2. Order structures of matrices	37
2.1. The relation \succ in the space B^1	37
2.2. Some doubly stochastic maps	40
2.3. Convexity and the relation \succ	45
2.4. Another partial order	47
2.5. Further examples: $A^k \succ A ^k$ and related inequalities	52
2.6. Miscellaneous	55
3. The order structure in the state space of C^*- and W^*-algebras	58
3.1. Preliminaries	58
3.2. The relation \succ ; definitions and elementary results	60
3.3. Some essential results for W^* -algebras	63
4. The c-ideal	65
4.1. Definition and the existence of the c -ideal	65
4.2. Ky Fan functionals and von Neumann's relation	68
4.3. A centre-valued convex trace	71
4.4. Maximally mixed states (chaotic states)	76
4.5. Technical Lemmata	78
5. The Σ-property	81
5.1. The definition, preliminary remarks	81
5.2. The case of properly infinite projections — type III	82
5.3. The finite case	83
5.4. The universality of the Σ -property for W^* -algebras	85
5.5. A duality theorem, remarks	87
5.6. Positive linear forms on C^* -algebras and the Σ -property	87

6. The dual structure in W^*-algebras	91
6.1. Introduction	91
6.2. Technical preliminaries	92
6.3. The purely infinite case	94
6.4. Approximation results	97
6.5. A characterization of (\mathcal{M}_+, \succ)	105
6.6. The main theorem	107
6.7. A special case: Finite W^* -algebras	111
6.8. Examples for \succ on \mathcal{M}_+	113
References	116
Symbols	121
Key words	122

Introduction

It is our intention to explain and to prove certain relations between stochasticity and partial order within this book.

An elementary but typical example is the following theorem due to RADO. Let us consider two n -tuples of real numbers, $\alpha = \{a_1, a_2, \dots\}$ and $\mathbf{b} = \{b_1, b_2, \dots\}$, which are decreasingly ordered, i.e. $a_1 \geq a_2 \geq \dots$ and $b_1 \geq b_2 \geq \dots$ respectively, and which fulfil $a_1 + a_2 + \dots + a_n = b_1 + b_2 + \dots + b_n$. Then α is a convex sum of n -tuples $\mathbf{b}^{(k)}$, where every $\mathbf{b}^{(k)}$ denotes a permutation of the n -tuple \mathbf{b} , if and only if

$$\forall m: \sum_{j=1}^m a_j \leq \sum_{j=1}^m b_j. \quad (1)$$

Using Birkhoff's result concerning the structure of doubly stochastic matrices, one at once infers the equivalence of (1) under the assumptions above with the existence of a doubly stochastic matrix T mapping \mathbf{b} onto α , i.e. $T\mathbf{b} = \alpha$.

For reasons of physical interpretation we call α "more chaotic" than \mathbf{b} iff $\alpha = T\mathbf{b}$ with a doubly stochastic T . In the first chapter, there are several examples explaining this point. In particular we refer to the increase of all "entropy-like quantities" if probability vectors become more chaotic.

After mainly dealing with now-classical results and some of their recent applications in Chapter 1, we extend these considerations to matrices in Chapter 2. Though there is little known of the structure of doubly stochastic transformations in matrix spaces (see Definition 2-2), one can establish similar connections between such doubly stochastic and related maps and appropriate partial (pre-)orderings. In the first two chapters we select, hopefully, a representative part of our present knowledge in that domain, although several branches cannot even be touched.

In Chapter 3 we indicate how to relate all that to the geometry of state spaces of finite-dimensional \ast -algebras and intend to convince the reader of the inescapability of extending it to general W^\ast -algebras.

The main tools and proofs for doing so are given in Chapters 4 and 5. We believe part of our techniques to be of general interest in the theory of W^\ast -algebras; for example, the Ky Fan functionals, the c -ideal, the central-valued convex trace, and the Σ -property.

In the last chapter, we introduce the dual structure to the partial (pre-)ordering \succsim of the state space. Let a, b be two Hermitian elements of a W^\ast -algebra. We write $a \preceq b$ iff for all positive linear functionals ω of the algebra, we have

$$\sup \omega(a^u) \geq \sup \omega(b^u). \quad (2)$$

Here a^* , b^* denote unitary transforms of the elements a , b , and the supremum has to run through all unitaries of the algebra in question. In this way one gets an extension of the famous von Neumann-Murray classification of pairs of projectors to pairs of Hermitian elements.

We do not present analogues in W^* -algebras for all the theorems of the first two chapters. However, using the technicalities developed, the skilled reader can find many of them by himself. Furthermore, one may obviously extend this theory completely to C^* -algebras with "sufficiently many projections", i.e. to AW^* -algebras. For general C^* -algebras, however, the results are essentially incomplete.

As already mentioned, there are parts of our work that are related to physics, including the description of some irreversible processes, the classification of mixed (in the sense of GIBBS and VON NEUMANN) states, and the problem of general diffusions. There are connections to non-commutative probability theory and non-commutative ergodicity.

Last not least, we point to the following statement: With the exception of the first chapter, the theory presented relies on the use of the unitary transformations of a W^* -algebra. Therefore, without some care, one gets trivialities for commutative algebras. In 1.9. a relevant idea for the commutative case is explained. Its implications for general commutative C^* -algebras will be considered elsewhere. *)

*) P. M. ALBERTI and A. UHLMANN, Dissipative motion in state spaces, Teubner-Texte zur Mathematik Bd. 33, Leipzig 1981.