

Rényi relative entropies and sufficiency of quantum channels

Anna Jenčová

Mathematical Institute, Slovak Academy of Sciences

Beyond IID, Cambridge, July 2018

Dedicated to the memory of Dénes Petz

Classical Rényi relative entropies

For p, q probability measures over a finite set X , $0 < \alpha \neq 1$:

$$D_{\alpha}(p\|q) = \frac{1}{\alpha - 1} \log \sum_x p(x)^{\alpha} q(x)^{1-\alpha}$$

In the limit $\alpha \rightarrow 1$: relative entropy

$$S(p\|q) = \sum_x p(x) \log(p(x)/q(x))$$

- ▶ introduced as the unique family of divergences satisfying a set of postulates
- ▶ fundamental quantities appearing in many information - theoretic tasks

Standard quantum Rényi relative entropies

For density matrices ρ, σ , $0 < \alpha \neq 1$,

$$D_\alpha(\rho\|\sigma) = \frac{1}{\alpha - 1} \log \operatorname{Tr} \rho^\alpha \sigma^{1-\alpha}$$

In the limit $\alpha \rightarrow 1$: quantum (Umegaki) relative entropy

$$S(\rho\|\sigma) = \operatorname{Tr} \rho(\log(\rho) - \log(\sigma))$$

- ▶ obtained from [Petz quasi-entropies](#)^{1,2}
- ▶ defined for normal states of a von Neumann algebra, using the [relative modular operator](#)

¹D. Petz, *Rep. Math. Phys.*, 1986

²D. Petz, Publ. RIMS, Kyoto Univ., 1985

Standard quantum Rényi relative entropies

It follows from the properties of quasi-entropies that: if $\alpha \in (0, 2]$

- ▶ strict **positivity**: $D_\alpha(\rho\|\sigma) \geq 0$ with equality iff $\rho = \sigma$;
- ▶ **data processing inequality**:

$$D_\alpha(\rho\|\sigma) \geq D_\alpha(\Phi(\rho)\|\Phi(\sigma))$$

for any quantum channel Φ

- ▶ **joint lower semicontinuity**
- ▶ **joint (quasi)-convexity**: the map

$$(\rho, \sigma) \mapsto \exp\{(\alpha - 1)D_\alpha(\rho\|\sigma)\}$$

is jointly convex.

Equality in DPI: sufficient quantum channels

Let the quantum states ρ, σ and a channel Φ be such that

$$S(\Phi(\rho) \parallel \Phi(\sigma)) = S(\rho \parallel \sigma) < \infty$$

This condition was introduced as a quantum extension of classical **sufficient statistics**:

A statistic T is sufficient with respect to a pair of probability distributions $\{p, q\}$ if

- ▶ the **conditional expectation** satisfies $E_p[\cdot | T] = E_q[\cdot | T]$
- ▶ an equivalent **Kullback-Leibler characterization** by the classical relative entropy:

$$S(p^T \parallel q^T) = S(p \parallel q) \text{ (if } < \infty \text{)}$$

Sufficient quantum channels

Let ρ, σ be quantum states (normal states of a von Neumann algebra), σ faithful. Assume that $S(\rho\|\sigma) < \infty$

Theorem

The following are equivalent.

- ▶ $S(\rho\|\sigma) = S(\Phi(\rho)\|\Phi(\sigma))$;
- ▶ There is a quantum channel Ψ such that

$$\Psi \circ \Phi(\rho) = \rho, \quad \Psi \circ \Phi(\sigma) = \sigma$$

We say in this case that Φ is **sufficient (reversible)** with respect to $\{\rho, \sigma\}$.

Sufficient quantum channels: divergences

A divergence D characterizes sufficiency if

$$D(\Phi(\rho)\|\Phi(\sigma)) = D(\rho\|\sigma) < \infty$$

implies that Φ is sufficient with respect to $\{\rho, \sigma\}$.

The following divergences characterize sufficiency:

- ▶ relative entropy
- ▶ D_α , with $\alpha \in (0, 2)^{3,4}$
- ▶ a class of f -divergences (in finite dimension)^{5,6}

³D. Petz, *Quart. J. Math. Oxford*, 1988

⁴AJ and D. Petz, 2006

⁵Hiai, Mosonyi, Petz, Bény, 2011

⁶F. Hiai and M. Mosonyi, 2017

Sufficient quantum channels: universal recovery channel

The **Petz recovery channel** is defined as

$$\Phi_{\sigma}(Y) = \sigma^{1/2} \Phi^{*}(\Phi(\sigma)^{-1/2} Y \Phi(\sigma)^{-1/2}) \sigma^{1/2}$$

Note that we always have $\Phi_{\sigma} \circ \Phi(\sigma) = \sigma$.

Theorem

Φ is sufficient with respect to $\{\rho, \sigma\}$ if and only if

$$\Phi_{\sigma} \circ \Phi(\rho) = \rho.$$

Sufficient quantum channels: a conditional expectation

Note that by the last condition, ρ (and σ) must be invariant states of the channel $\Phi_\sigma \circ \Phi$.


There is a conditional expectation E , $\sigma \circ E = \sigma$, such that Φ is sufficient with respect to $\{\rho, \sigma\}$ if and only if $\rho \circ E = \rho$.

Structure of the states⁷: in finite dimensions (or on $B(\mathcal{H})$), there is a decomposition

$$U\sigma U^* = \bigoplus_n \lambda_n \sigma_n^L \otimes \sigma_n^R, \quad \sigma_n^L, \sigma_n^R \text{ states, } \lambda_n \text{ probabilities}$$

such that Φ is sufficient with respect to $\{\rho, \sigma\}$ iff

$$U\rho U^* = \bigoplus_n \mu_n \rho_n^L \otimes \sigma_n^R, \quad \rho_n^L \text{ states, } \mu_n \text{ probabilities}$$

⁷M. Mosonyi and D. Petz, 2004; AJ and D. Petz, 2006 

Sufficient quantum channels: applications

Characterization of equality in various entropic inequalities:

- ▶ strong subadditivity: characterization of quantum Markov states⁸
- ▶ monotonicity of quantum Fisher information, Holevo quantity, etc.

Approximate version - **recoverability**⁹:

$$S(\rho\|\sigma) - S(\Phi(\rho)\|\Phi(\sigma)) \geq d(\rho\|\tilde{\Phi}_\sigma \circ \Phi(\rho))$$

d some divergence measure, $\tilde{\Phi}_\sigma$ a modification of Petz recovery channel

⁸Hayden, Josza, Petz, Winter, 2004

⁹O. Fawzi and R. Renner, 2014; M. M. Wilde, 2015; Junge et. al, 2015; ..

Sandwiched Rényi relative entropy

Another version:

for density matrices ρ, σ :

$$\tilde{D}_\alpha(\rho\|\sigma) = \frac{1}{\alpha-1} \log \text{Tr} \left[\left(\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}} \right)^\alpha \right]$$

M. Müller-Lennert et al., *J. Math. Phys.*, 2013

M. M. Wilde et al., *Commun. Math. Phys.*, 2014

- ▶ satisfies DPI (+ other properties) if $\alpha \in [1/2, 1) \cup (1, \infty)$
- ▶ $\lim_{\alpha \rightarrow 1} \tilde{D}_\alpha(\rho\|\sigma) = S(\rho\|\sigma)$
- ▶ operational interpretation for $\alpha > 1$: strong converse exponents in quantum hypothesis testing¹⁰

¹⁰M. Mosonyi, and T. Ogawa, *Commun. Math. Phys.*, 2017

The purpose of the rest of this talk

Extend the sandwiched Rényi relative entropies to normal states of von Neumann algebras and show some properties

- ▶ the standard version D_α (quasi-entropies) is defined in this setting and has an operational interpretation in hypothesis testing as in finite dimensions¹¹

In this general setting, prove that \tilde{D}_α characterize sufficiency of channels, for $\alpha \in (1/2, 1) \cup (1, \infty)$.

¹¹V. Jaksic et al., *Rev. Math. Phys.*, 2012

Extensions of \tilde{D}_α to von Neumann algebras

Let ρ, σ be normal states on a von Neumann algebra \mathcal{M} .

Two constructions, using noncommutative L_p -spaces:

- ▶ Araki-Masuda divergences, defined for $\alpha \in [1/2, 1) \cup (1, \infty]$, uses Araki-Masuda L_p -spaces¹²
- ▶ sandwiched Rényi relative entropies, defined for $\alpha > 1$, uses Kosaki L_p -spaces¹³

¹²M. Berta, V. B. Scholz, and M. Tomamichel, Ann. H. Poincaré, 2018

¹³AJ, Ann. H. Poincaré, 2018

Haagerup L_p -spaces

For $1 \leq p \leq \infty$, $L_p(\mathcal{M})$ - Haagerup L_p -space, with norm $\|\cdot\|_p$

- ▶ $\mathcal{M} \simeq L_\infty(\mathcal{M})$;
- ▶ the predual $\mathcal{M}_* \simeq L_1(\mathcal{M})$: $\rho \mapsto h_\rho$, $\text{Tr } h_\rho = \rho(1)$;
- ▶ $L_2(\mathcal{M})$ a Hilbert space: $\langle h, k \rangle = \text{Tr } k^* h$

If $\mathcal{M} = B(\mathcal{H})$, we can use the Schatten classes:

$$L_p(\mathcal{M}) = \{X \in B(\mathcal{H}), \text{Tr } |X|^p < \infty\}, \quad \|X\|_p = (\text{Tr } |X|^p)^{1/p}$$

The standard form

Standard form: $(\lambda(\mathcal{M}), L_2(\mathcal{M}), J, L_2(\mathcal{M})^+)$

a **representation** of \mathcal{M} on $L_2(\mathcal{M})$ by left multiplication:

$$\lambda(x)h = xh, \quad x \in \mathcal{M}, \quad h \in L_2(\mathcal{M})$$

Any $\rho \in \mathcal{M}_*^+$ has a unique **vector representative** $h_\rho^{1/2}$ in $L_2(\mathcal{M})^+$:

$$\rho(a) = \langle ah_\rho^{1/2}, h_\rho^{1/2} \rangle$$

Kosaki L_p -spaces with respect to a faithful normal state

Let σ be a faithful normal state:

- ▶ continuous embedding

$$\mathcal{M} \rightarrow L_1(\mathcal{M}), \quad x \mapsto h_\sigma^{1/2} x h_\sigma^{1/2}$$

- ▶ Put

$$L_\infty(\mathcal{M}, \sigma) := h_\sigma^{1/2} \mathcal{M} h_\sigma^{1/2}, \quad \|h_\sigma^{1/2} x h_\sigma^{1/2}\|_{\infty, \sigma} = \|x\|_\infty$$

- ▶ let $\rho \in \mathcal{M}_*^+$, then $h_\rho \in L_\infty(\mathcal{M}, \sigma)$ iff $\rho \leq \lambda \sigma$ and

$$\|h_\rho\|_{\infty, \sigma} = \inf\{\lambda > 0, \rho \leq \lambda \sigma\}.$$

Kosaki L_p -spaces with respect to a faithful normal state

Let $1 < p < \infty$:

- ▶ $(L_\infty(\mathcal{M}, \sigma), L_1(\mathcal{M}))$ compatible pair of Banach spaces
- ▶ interpolation space

$$L_p(\mathcal{M}, \sigma) := C_{1/p}(L_\infty(\mathcal{M}, \sigma), L_1(\mathcal{M})), \text{ with norm } \|\cdot\|_{p,\sigma}$$

- ▶ Let $1/p + 1/q = 1$, then

$$L_p(\mathcal{M}, \sigma) = \{h_\sigma^{1/2q} k h_\sigma^{1/2q}, k \in L_p(\mathcal{M})\},$$

$$\|h_\sigma^{1/2q} k h_\sigma^{1/2q}\|_{p,\sigma} = \|k\|_p$$

A definition of \tilde{D}_α , $\alpha > 1$

Extension to non-faithful σ : by restriction to support $s(\sigma) = e$

$$L_p(\mathcal{M}, \sigma) = \{h \in L_1(\mathcal{M}), h = ehe \in L_p(e\mathcal{M}e, \sigma|_{e\mathcal{M}e})\}.$$

For normal states ρ, σ and $1 < \alpha < \infty$:

$$\tilde{D}_\alpha(\rho||\sigma) = \begin{cases} \frac{\alpha}{\alpha-1} \log(\|h_\rho\|_{\alpha,\sigma}) & \text{if } h_\rho \in L_\alpha(\mathcal{M}, \sigma) \\ \infty & \text{otherwise.} \end{cases}$$

Some properties of \tilde{D}_α

Using complex interpolation, we can prove

- ▶ **strict positivity**: $\tilde{D}_\alpha(\rho\|\sigma) \geq 0$, with equality iff $\rho = \sigma$.
- ▶ **joint lower semicontinuity** (on $L_1(\mathcal{M})^+ \times L_1(\mathcal{M})^+$)
- ▶ if $\rho \neq \sigma$ and $\tilde{D}_\alpha(\rho\|\sigma) < \infty$, then

$\alpha' \mapsto \tilde{D}_{\alpha'}(\rho\|\sigma)$ is **strictly increasing** for $\alpha' \in (1, \alpha]$.

- ▶ **quasi-convexity**

Relation to the standard version D_α , limit values

For normal states ρ, σ , $\alpha > 1$:

$$D_{2-1/\alpha}(\rho\|\sigma) \leq \tilde{D}_\alpha(\rho\|\sigma) \leq D_\alpha(\rho\|\sigma)$$



Limit values:

$$\lim_{\alpha \rightarrow 1} \tilde{D}_\alpha(\rho\|\sigma) = S(\rho\|\sigma)$$

$$\lim_{\alpha \rightarrow \infty} \tilde{D}_\alpha(\rho\|\sigma) = \tilde{D}_\infty(\rho\|\sigma) \text{ relative max entropy}$$

Data processing inequality

$\Phi : L_1(\mathcal{M}) \rightarrow L_1(\mathcal{N})$ positive, trace-preserving. Let $\sigma_0 = \Phi(\sigma)$.

- ▶ Φ is a contraction $L_1(\mathcal{M}) \rightarrow L_1(\mathcal{N})$.
- ▶ if $0 \leq \rho \leq \lambda\sigma$, then also $0 \leq \Phi(\rho) \leq \lambda\sigma_0$, hence

$$\Phi(L_\infty(\mathcal{M}, \sigma)^+) \rightarrow L_\infty(\mathcal{N}, \sigma_0)^+$$

- ▶ this extends to a map $x \mapsto y$, $\Phi(h_\sigma^{1/2} x h_\sigma^{1/2}) = h_{\sigma_0}^{1/2} y h_{\sigma_0}^{1/2}$
(adjoint of) the [Petz recovery channel](#)¹⁴ Φ_σ
- ▶ Φ defines a contraction $L_\infty(\mathcal{M}, \sigma) \rightarrow L_\infty(\mathcal{N}, \sigma_0)$.

Φ defines a contraction $L_p(\mathcal{M}, \sigma) \rightarrow L_p(\mathcal{N}, \sigma_0)$ for all $1 \leq p \leq \infty$.

¹⁴D. Petz, Quart. J. Math. Oxford, 1984

Data processing inequality

For $\alpha > 1$, normal states ρ, σ , **positive**, trace-preserving Φ :

$$\tilde{D}_\alpha(\rho\|\sigma) \geq \tilde{D}_\alpha(\Phi(\rho)\|\Phi(\sigma))$$

Consequently, by the limit $\alpha \rightarrow 1$:

For normal states ρ, σ ,

$$S(\rho\|\sigma) \geq S(\Phi(\rho)\|\Phi(\sigma))$$

holds for any **positive** trace-preserving map Φ .

A. Müller-Hermes, D. Reeb, Ann. H. Poincaré, 2017 (for $\mathcal{M} = B(\mathcal{H})$)

The Araki-Masuda divergences

The Araki-Masuda L_p -norm: defined on $L_2(\mathcal{M})$

- ▶ for $2 \leq p \leq \infty$, $\xi \in L_2(\mathcal{M})^+$,

$$\|\xi\|_{p,\sigma}^{AM} = \sup_{\omega \in \mathcal{M}_*^+, \omega(1)=1} \|\Delta_{\omega,\sigma}^{1/2-1/p} \xi\|_2$$

if $s(\omega_\xi) \leq s(\sigma)$ and is infinite otherwise

- ▶ for $1 \leq p < 2$,

$$\|\xi\|_{p,\sigma}^{AM} = \inf_{\omega \in \mathcal{M}_*^+, \omega(1)=1, s(\omega) \geq s(\omega_\xi)} \|\Delta_{\omega,\sigma}^{1/2-1/p} \xi\|_2$$

The Araki-Masuda divergences

For normal states ρ, σ and $\alpha \in [1/2, 1) \cup (1, \infty)$:

$$\tilde{D}_{\alpha}^{AM}(\rho\|\sigma) = \frac{2\alpha}{\alpha-1} \log(\|h_{\rho}^{1/2}\|_{2\alpha,\sigma}^{AM})$$

- ▶ can be defined using any $*$ -representation of \mathcal{M} on a Hilbert space \mathcal{H} and any vector $\xi \in \mathcal{H}$ representing ρ
- ▶ duality relation: for $1/p + 1/q = 1$

$$|\langle \eta, \xi \rangle| \leq \|\eta\|_{p,\sigma}^{AM} \|\xi\|_{q,\sigma}^{AM}, \quad \xi, \eta \in \mathcal{H}$$

- ▶ if $1 < p \leq 2$, there is a (unique) unit vector $\eta_0 \in \mathcal{H}$ such that

$$\langle \eta, \eta_0 \rangle = \|\eta\|_{p,\sigma}^{AM} \|\eta_0\|_{q,\sigma}^{AM}$$

$$\tilde{D}_\alpha^{AM} \text{ and } \tilde{D}_\alpha$$

Araki-Masuda L_p -norms can be introduced by interpolation:

- ▶ For $2 \leq p \leq \infty$: a continuous embedding $\mathcal{M} \rightarrow L_2(\mathcal{M})$

$$x \mapsto h_\sigma^{1/2} x, \quad x \in \mathcal{M}$$

the interpolation norm $\|\cdot\|_{p,\sigma}^{AM}$ in $C_{1/p}(\mathcal{M}, L_2(\mathcal{M}))$.

- ▶ For $1 \leq p \leq 2$: a continuous embedding $L_2(\mathcal{M}) \rightarrow L_1(\mathcal{M})$

$$k \mapsto kh_\sigma^{1/2}, \quad k \in L_2(\mathcal{M})$$

the interpolation norm $\|\cdot\|_{p,\sigma}^{AM}$ in $C_{1/p}(L_2(\mathcal{M}), L_1(\mathcal{M}))$.

\tilde{D}_α^{AM} and \tilde{D}_α

For $1 < \alpha < \infty$,

$$\tilde{D}_\alpha(\rho\|\sigma) = \tilde{D}_\alpha^{AM}(\rho\|\sigma)$$

For $1/2 < \alpha < 1$: $h_\sigma^{\frac{1-\alpha}{2\alpha}} h_\rho^{1/2} \in L_{2\alpha}(\mathcal{M})$ and

$$\tilde{D}_\alpha(\rho\|\sigma) := \tilde{D}_\alpha^{AM}(\rho\|\sigma) = \frac{2\alpha}{\alpha-1} \log \|h_\sigma^{\frac{1-\alpha}{2\alpha}} h_\rho^{1/2}\|_{2\alpha}$$

Limit values:

- ▶ $\lim_{\alpha \nearrow 1} \tilde{D}_\alpha(\rho\|\sigma) = S(\rho\|\sigma)$
- ▶ $\lim_{\alpha \rightarrow 1/2} \tilde{D}_\alpha(\rho\|\sigma) = -\log F(\rho, \sigma)$ Uhlmann's fidelity

Data processing inequality for \tilde{D}_α , $\alpha \in (1/2, 1)$

We have to assume that $\Phi : L_1(\mathcal{M}) \rightarrow L_1(\mathcal{N})$ is trace preserving and **completely positive**, with Stinespring representation:

$$\Phi^* = T^* \pi(\cdot) T$$

π a $*$ -representation, T an isometry

- ▶ Let $p = 2\alpha$, $1/p + 1/q = 1$. Let $\rho = \omega_\eta$ and let $\omega := \omega_{\eta_0}$ be such that $\langle \eta, \eta_0 \rangle = \|\eta\|_{p,\sigma}^{AM} \|\eta_0\|_{q,\sigma}^{AM}$. Then

$$\langle \eta, \eta_0 \rangle = \langle T\eta, T\eta_0 \rangle \leq \|T\eta\|_{p,\Phi(\sigma)}^{AM} \|T\eta_0\|_{q,\Phi(\sigma)}^{AM}$$

- ▶ we obtain, with $\alpha^* = \alpha/(2\alpha - 1) > 1$:

$$\begin{aligned} \tilde{D}_\alpha(\rho\|\sigma) &\geq \tilde{D}_\alpha(\Phi(\rho)\|\Phi(\sigma)) + \tilde{D}_{\alpha^*}(\omega\|\sigma) - \tilde{D}_{\alpha^*}(\Phi(\omega)\|\Phi(\sigma)) \\ &\geq \tilde{D}_\alpha(\Phi(\rho)\|\Phi(\sigma)) \end{aligned}$$

Characterizations of sufficient channels by \tilde{D}_α

Theorem

The sandwiched Rényi relative entropies \tilde{D}_α characterize sufficiency for $\alpha \in (1/2, 1) \cup (1, \infty)$.

AJ, Ann. H. Poincaré, 2018; AJ, arXiv:1707.00047

That is:

$$\tilde{D}_\alpha(\Phi(\rho) \parallel \Phi(\sigma)) = \tilde{D}_\alpha(\rho \parallel \sigma) < \infty$$

implies that

$$\Psi \circ \Phi(\rho), \quad \Psi \circ \Phi(\sigma).$$

for some channel Ψ .

The proof in case $\alpha > 1$

Let $\alpha > 1$.

- ▶ the assumption

$$\tilde{D}_\alpha(\Phi(\rho) \parallel \Phi(\sigma)) = \tilde{D}_\alpha(\rho \parallel \sigma) < \infty$$

implies that $h_\rho \in L_\alpha(\mathcal{M}, \sigma)$ and Φ is a contraction preserving its norm.

- ▶ An easy proof for $\alpha = 2$:

$L_2(\mathcal{M}, \sigma)$ is a Hilbert space, Φ_σ is the adjoint of Φ

By properties of contractions on Hilbert spaces:

$$\Phi_\sigma \circ \Phi(h_\rho) = h_\rho$$

(note that **positivity** of Φ is enough for this)

The proof in case $\alpha > 1$

For general $\alpha > 1$, use [interpolation](#):

Let τ be a normal state, $s(\tau) \leq s(\sigma)$. Put

$$h_\tau(z) = h_\sigma^{(1-z)/2} h_\tau^z h_\sigma^{(1-z)/2} \in L_1(\mathcal{M}), \quad 0 \leq \operatorname{Re}(z) \leq 1,$$

continuous function, analytic in $0 < \operatorname{Re}(z) < 1$.

If the equality

$$\|\Phi(h_\tau(1/\alpha))\|_{\alpha, \Phi(\sigma)} = \|h_\tau(1/\alpha)\|_{\alpha, \sigma}$$

holds for some $\alpha > 1$, then it holds for all $\alpha > 1$.

The proof in case $\alpha > 1$

By assumptions $h_\rho = t h_\tau(1/\alpha)$ for some state τ , $t > 0$ - normalization, and we have

$$\|\Phi(h_\tau(1/\alpha))\|_{\alpha, \Phi(\sigma)} = \|h_\tau(1/\alpha)\|_{\alpha, \sigma}.$$

Then the equality holds also for $\alpha = 2$, so that

Φ is sufficient with respect to $\{\omega, \sigma\}$, where

$$h_\omega = s h_\tau(1/2) = s h_\sigma^{1/4} h_\tau^{1/2} h_\sigma^{1/4}, \quad s > 0.$$

Ok, but this is not what we wanted to prove!

The proof in case $\alpha > 1$

Assume $\mathcal{M} = B(\mathcal{H})$, replace h_ρ by the density operator ρ .

There is a decomposition $U\sigma U^* = \bigoplus_n \lambda_n \sigma_n^L \otimes \sigma_n^R$ such that Φ is sufficient with respect to $\{\rho, \sigma\}$ iff

$$U\rho U^* = \bigoplus_n \mu_n \rho_n^L \otimes \sigma_n^R$$

But we have such a decomposition for $\omega = s\sigma^{1/4}\tau^{1/2}\sigma^{1/4}$, hence also for $\rho = t\sigma^{1/2\beta}\tau^{1/\alpha}\sigma^{1/2\beta}$, which implies the result.

(In the general case, we use the characterization by conditional expectations.)

The case $\alpha \in (1/2, 1)$

Let $\alpha \in (1/2, 1)$. Then

$$h_{\sigma}^{\frac{1-\alpha}{2\alpha}} h_{\rho}^{1/2} = h_{\tau}^{1/2\alpha} u \in L_{2\alpha}(\mathcal{M})$$

for some $\tau \in \mathcal{M}_{*}^{+}$ and partial isometry $u \in \mathcal{M}$.

- We recall the inequality

$$\begin{aligned}\tilde{D}_{\alpha}(\rho\|\sigma) &\geq \tilde{D}_{\alpha}(\Phi(\rho)\|\Phi(\sigma)) + \tilde{D}_{\alpha^{*}}(\omega\|\sigma) - \tilde{D}_{\alpha^{*}}(\Phi(\omega)\|\Phi(\sigma)) \\ &\geq \tilde{D}_{\alpha}(\Phi(\rho)\|\Phi(\sigma))\end{aligned}$$

for $\alpha^{*} > 1$ and some state ω .

- Actually, we have $h_{\omega} = t h_{\tau}(1/\alpha^{*})$.

The case $\alpha \in (1/2, 1)$

- ▶ By assumptions, we obtain

$$\tilde{D}_{\alpha^*}(\omega \parallel \sigma) = \tilde{D}_{\alpha^*}(\Phi(\omega) \parallel \Phi(\sigma))$$

- ▶ since $\alpha^* > 1$, this implies that Φ is sufficient with respect to $\{\omega, \sigma\}$.
- ▶ use the decompositions (conditional expectations) as before.