

Notes for $\alpha - z$ Rényi divergence

Anna Jenčová

December 7, 2023

1 What is proved

1.1 DPI

This is proved for

$$\alpha \in (0, 1), \max\{\alpha, 1 - \alpha\} \leq z \quad \text{and} \quad \alpha > 1, \max\{\alpha/2, \alpha - 1\} \leq z \leq \alpha$$

See [5, Thm. 1(viii)], [4]. It is not possible to go beyond these bounds, [9].

1.2 Lower semicontinuity (LS)

Holds for $\alpha \in (0, 1)$ and for $\alpha > 1, z \geq \alpha/2$, [5].

1.3 Variational expressions

For $\alpha \in (0, 1), z \geq \max\{\alpha, 1 - \alpha\}$, we have [5, Theorem 1(vi)]

$$Q_{\alpha,z}(\psi\|\varphi) = \inf_{a \in \mathcal{M}_{++}} \left\{ \alpha \text{Tr} \left((a^{1/2} h_{\psi}^{\alpha/z} a^{1/2})^{z/\alpha} \right) + (1 - \alpha) \text{Tr} \left(a^{-1/2} h_{\varphi}^{(1-\alpha)/z} a^{-1/2} \right) \right\}.$$

For $\alpha > 1, z \geq \alpha/2$, we have [5, Theorem 2(vi)] and [6]

$$Q_{\alpha,z}(\psi\|\varphi) = \sup_{a \in \mathcal{M}_+} \left\{ \alpha \text{Tr} \left((a^{1/2} h_{\psi}^{\alpha/z} a^{1/2})^{z/\alpha} \right) - (\alpha - 1) \text{Tr} \left(a^{1/2} h_{\varphi}^{(\alpha-1)/z} a^{1/2} \right) \right\}.$$

The lower bound comes from the lower bound in LS. Proved for all $z > 0$ in type I case [8].

1.4 Martingale convergence

Holds in the bounds for DPI, [3]. A remark to this proof: I think that the proof of [3, Eq. (0.4)] can be simplified. The key ingredient here is the martingale convergence of the generalized conditional expectations, which gives [3, Eqs. (0.5)]

$$\psi_i \circ \mathcal{E}_{\mathcal{M}_i, \varphi} \rightarrow \psi \quad \text{in norm.}$$

(We denote $\varphi_i := \varphi|_{\mathcal{M}_i}$ and $\psi_i = \psi|_{\mathcal{M}_i}$.) In the bounds for DPI, we also have LS, so that

$$D_{\alpha,z}(\psi_i\|\varphi_i) \geq D_{\alpha,z}(\psi_i \circ \mathcal{E}_{\mathcal{M}_i, \varphi} \|\varphi_i \circ \mathcal{E}_{\mathcal{M}_i, \varphi})$$

and (using also [3, Eq.(0.5)])

$$\sup_i D_{\alpha,z}(\psi_i\|\varphi_i) \geq \liminf_i D_{\alpha,z}(\psi_i \circ \mathcal{E}_{\mathcal{M}_i, \varphi} \|\varphi_i \circ \mathcal{E}_{\mathcal{M}_i, \varphi}) \geq D_{\alpha,z}(\psi\|\varphi)$$

1.5 Monotonicity in z

For $\alpha > 1$, monotonicity is as

$$0 < z \leq z' \implies D_{\alpha,z}(\psi\|\varphi) \geq D_{\alpha,z'}(\psi\|\varphi).$$

This was proved in [2, Sec. 3] in type II_1 algebras, under the condition of lower semicontinuity of the map $\varphi \mapsto D_{\alpha,z'}(\psi\|\varphi)$, in particular, it holds for $\alpha/2 \leq z'$. Indeed, by the proof in [2, Sec. 3], we have for $z \leq z'$ and all $\varepsilon > 0$

$$Q_{\alpha,z}(\psi\|\varphi) \geq Q_{\alpha,z}(\psi\|\varphi + \varepsilon\tau) \geq Q_{\alpha,z'}(\psi\|\varphi + \varepsilon\tau),$$

so that the inequality follows by LS for z' .

Extension to the general case can be done using Haagerup reduction. For this, we so far need to assume that

$$\alpha - 1, \alpha/2 \leq z \leq z' \leq \alpha.$$

Let $\hat{\mathcal{M}}, \hat{\mathcal{M}}_n, \hat{\psi}, \hat{\varphi}$ be as in the Haagerup reduction and put $\psi_n = \hat{\psi}|_{\mathcal{M}_n}$, $\varphi_n = \hat{\varphi}|_{\mathcal{M}_n}$. Using [2, Sec. 3], we have for $z \leq z'$ and all n

$$D_{\alpha,z'}(\psi_n\|\varphi_n) \leq D_{\alpha,z}(\psi_n\|\varphi_n).$$

Using DPI and LS, we obtain

$$D_{\alpha,z'}(\psi\|\varphi) = D_{\alpha,z'}(\hat{\psi}\|\hat{\varphi}) \leq \liminf_n D_{\alpha,z'}(\psi_n\|\varphi_n) \leq \liminf_n D_{\alpha,z}(\psi_n\|\varphi_n) \leq D_{\alpha,z}(\hat{\psi}\|\hat{\varphi}) = D_{\alpha,z}(\psi\|\varphi).$$

It would be very useful if we could extend the use of Haagerup reduction beyond these bounds, see for example Section 2.2 below.

2 Some further results and remarks

2.1 Convergence of $D_{\alpha,z}$ as $\alpha \nearrow 1$

Using monotonicity in z , we can prove this for $z \leq 1$.

Lemma 1. *Assume that $0 \leq 1 - z < \alpha < 1$. Then for any normal state ψ ,*

$$D_{\beta,1}(\psi\|\varphi) \leq D_{\alpha,z}(\psi\|\varphi) \leq D_{\alpha,1}(\psi\|\varphi),$$

where $\beta = \frac{\alpha-1+z}{z}$.

Proof. The statement is trivial for $z = 1$, so we may assume $0 < 1 - z$. The second inequality follows by monotonicity of $z \mapsto D_{\alpha,z}(\psi\|\varphi)$ for $\alpha \in (0, 1)$. For the first inequality, note first that by the assumption, $\beta \in (0, 1)$ and by Hölder

$$Q_{\alpha,z}(\psi\|\varphi)^{\frac{1}{2z}} = \|h_{\psi}^{\frac{\alpha}{2z}} h_{\varphi}^{\frac{1-\alpha}{2z}}\|_{2z} = \|h_{\psi}^{\frac{1-z}{2z}} h_{\psi}^{\frac{\beta}{2}} h_{\varphi}^{\frac{1-\beta}{2}}\|_{2z} \leq \|h_{\psi}^{\frac{\beta}{2}} h_{\varphi}^{\frac{1-\beta}{2}}\|_2 = Q_{\beta,1}(\psi\|\varphi)^{\frac{1}{2}}.$$

This proves the statement. □

Using the lemma for $1 - \alpha$ small enough and properties of $D_{\alpha,1}$, we get for any $z \leq 1$:

$$\lim_{\alpha \nearrow 1} D_{\alpha,z}(\psi\|\varphi) = D(\psi\|\varphi).$$

2.2 Convergence of $D_{\alpha,z}$ as $\alpha \searrow 1$

The strategy of the previous section is limited by the bounds for which we currently have monotonicity in z for $\alpha > 1$. Namely, the inequality $D_{\alpha,z}(\psi\|\varphi) \leq D_{\alpha,1}(\psi\|\varphi)$ is currently only proved for

$$1 \leq z \leq \alpha \leq 2,$$

which is violated as $\alpha \searrow 1$ (unless $z = 1$ or $z = \alpha$). We have the following lower bound.

Lemma 2. *Let $1 < \alpha \leq z$. Then for any normal state ψ , we have*

$$D_{\beta,1}(\psi\|\varphi) \leq D_{\alpha,z}(\psi\|\varphi),$$

where $\beta = \frac{\alpha-1+z}{z}$.

Proof. Assume that $D_{\alpha,z}(\psi\|\varphi) < \infty$, otherwise there is nothing to prove. Then there is some $y \in L_{2z}(\mathcal{M})$ such that

$$h_{\psi}^{\frac{\alpha}{2z}} = y h_{\varphi}^{\frac{\alpha-1}{2z}}, \quad Q_{\alpha,z}(\psi\|\varphi) = \|y\|_{2z}^{2z}.$$

Since $\alpha \leq z$, we have $\frac{1}{2} - \frac{\alpha}{2z} \geq 0$ and

$$h_{\psi}^{1/2} = h_{\psi}^{\frac{1}{2} - \frac{\alpha}{2z}} y h_{\varphi}^{\frac{\alpha-1}{2z}}.$$

It follows that $h_{\psi}^{1/2} \in \mathcal{D}(\Delta_{\psi,\varphi}^{\frac{\alpha-1}{2z}})$ and

$$\Delta_{\psi,\varphi}^{\frac{\alpha-1+z}{2z}} h_{\varphi}^{1/2} = \Delta_{\psi,\varphi}^{\frac{\alpha-1}{2z}} \Delta_{\psi,\varphi}^{1/2} h_{\varphi}^{1/2} = \Delta_{\psi,\varphi}^{\frac{\alpha-1}{2z}} h_{\psi}^{1/2} = h_{\psi}^{\frac{z-1}{2z}} y.$$

It follows that

$$\|\Delta_{\psi,\varphi}^{\frac{\alpha-1+z}{2z}} h_{\varphi}^{1/2}\|_2 \leq \psi(1)^{\frac{z-1}{2z}} \|y\|_{2z}.$$

We therefore have for $\psi(1) = 1$,

$$D_{\alpha,z}(\psi\|\varphi) = \frac{1}{\alpha-1} \log \|y\|_{2z}^{2z} \geq \frac{z}{\alpha-1} \log \|\Delta_{\psi,\varphi}^{\frac{\alpha-1+z}{2z}} h_{\varphi}^{1/2}\|_2^2 = D_{\frac{\alpha-1+z}{z},1}(\psi\|\varphi).$$

□

2.3 Complex interpolation

This is to remark that for $\alpha > 1$ and $z \geq \alpha/2$, $Q_{\alpha,z}(\psi\|\phi)$ can be written using the Kosaki interpolation norms. Assume that both ψ and φ are faithful and $Q_{\alpha,z}(\psi\|\phi) < \infty$. We then have

$$h_{\psi} = h_{\psi}^{1-\frac{\alpha}{2z}} y h_{\varphi}^{\frac{\alpha-1}{2z}} = h_{\psi}^{\eta/q} y h_{\varphi}^{(1-\eta)/q}$$

for some $y \in L_{2z}(\mathcal{M})$, $q = \frac{2z}{2z-1}$ (the dual parameter to $2z$) and $\eta = \frac{2z-\alpha}{2z-1} \in [0,1]$. Hence h_{ψ} belongs to the space $L_{2z}^{\eta}(\mathcal{M}, \psi, \varphi)$, where

$$L_p^{\eta}(\mathcal{M}, \psi, \varphi) := C_{1/p}(h_{\psi}^{\frac{\eta}{2}} \mathcal{M} h_{\varphi}^{\frac{1-\eta}{2}}, \mathcal{M}_*) = C_{\eta}(L_p(\mathcal{M}, \varphi)_L, L_p(\mathcal{M}, \psi)_R),$$

[7, Thm. 11.1]. Let $\|\cdot\|_{2z,\psi,\varphi,\eta}$ denote the norm in this space, then it is easily seen that

$$Q_{\alpha,z}(\psi\|\phi) = \|y\|_{2z}^{2z} = \|h_{\psi}\|_{2z,\psi,\varphi,\eta}^{2z}.$$

We may be able to use the fact that $L_p^{\eta}(\mathcal{M}, \psi, \varphi)$ form an interpolating family with respect to both p and η to prove some results, e.g. monotonicity in z or in α . It may be possible to extend this for other values of α and z using [1].

References

- [1] J. Gu, Z. Yin, and H. Zhang. Interpolation of quasi noncommutative L_p -spaces. *arXiv:1905.08491*, 2019.
- [2] F. Hiai, Questions, note.
- [3] F. Hiai, Martingale convergence for $D_{\alpha,z}$, note.
- [4] A. Jenčová, DPI for $\alpha - z$ -Rényi divergence, note.
- [5] S. Kato. On α - z -Rényi divergence in the von Neumann algebra setting. *arXiv preprint arXiv:2311.01748*, 2023.
- [6] S. Kato, Variational expression for $\alpha > 1$, note.
- [7] H. Kosaki. Applications of the complex interpolation method to a von Neumann algebra: Non-commutative L_p -spaces. *J. Funct. Anal.*, 56:26–78, 1984.
- [8] M. Mosonyi. The strong converse exponent of discriminating infinite-dimensional quantum states. *Communications in Mathematical Physics*, 400(1):83–132, 2023.
- [9] H. Zhang. From Wigner-Yanase-Dyson conjecture to Carlen-Frank-Lieb conjecture. *Advances in Mathematics*, 365:107053, 2020.