

Randomization theorems for quantum channels



Anna Jenčová

Mathematical Institute, Slovak Academy of Sciences

Banff, July 2019


The problem

Given two channels



The problem


how precisely can we approximate



- α is a **superchannel**

The problem


how precisely can we approximate



- ▶ α is a **superchannel**
- ▶ approximation in **diamond norm**

The problem


how precisely can we approximate



- ▶ α is a **superchannel**
- ▶ approximation in **diamond norm**
- ▶ one shot

The problem

how precisely can we approximate



- ▶ α is a **superchannel**
- ▶ approximation in **diamond norm**
- ▶ one shot
- ▶ all superchannels or some restrictions on α

Equality case

- ▶ comparison of bipartite channels: (Gour, 2018)

Equality case

- ▶ comparison of bipartite channels: (Gour, 2018)
- ▶ comparison of channels: $d_{A_0} = d_{B_0} = 1$, restrictions on α :

Equality case

- ▶ comparison of bipartite channels: (Gour, 2018)
- ▶ comparison of channels: $d_{A_0} = d_{B_0} = 1$, restrictions on α :
 - ▶ post-processings (degradability)

$$A_1 = A_2, \quad \Phi_1 \stackrel{?}{=} \alpha \circ \Phi_2$$

(Chefles 2009, Buscemi 2012,2016,...)

Equality case

- ▶ comparison of bipartite channels: (Gour, 2018)
- ▶ comparison of channels: $d_{A_0} = d_{B_0} = 1$, restrictions on α :
 - ▶ post-processings (degradability)

$$A_1 = A_2, \quad \Phi_1 \stackrel{?}{=} \alpha \circ \Phi_2$$

(Chefles 2009, Buscemi 2012, 2016, ...)

- ▶ comparison of quantum experiments: c-q channels

$$\rho_1, \dots, \rho_k \in \mathcal{S}(B_1), \quad \sigma_1, \dots, \sigma_k \in \mathcal{S}(B_2), \quad \rho_i \stackrel{?}{=} \alpha(\sigma_i), \forall i$$

(Alberti & Uhlmann 1980, Buscemi 2012, Gour et al. 2018)

Equality case

- ▶ comparison of bipartite channels: (Gour, 2018)
- ▶ comparison of channels: $d_{A_0} = d_{B_0} = 1$, restrictions on α :
 - ▶ post-processings (degradability)

$$A_1 = A_2, \quad \Phi_1 \stackrel{?}{=} \alpha \circ \Phi_2$$

(Chefles 2009, Buscemi 2012, 2016, ...)

- ▶ comparison of quantum experiments: c-q channels

$$\rho_1, \dots, \rho_k \in \mathcal{S}(B_1), \quad \sigma_1, \dots, \sigma_k \in \mathcal{S}(B_2), \quad \rho_i \stackrel{?}{=} \alpha(\sigma_i), \forall i$$

(Alberti & Uhlmann 1980, Buscemi 2012, Gour et al. 2018)

- ▶ comparison of classical experiments: (Blackwell 1953, Torgersen 1970)

Equality case

- ▶ comparison of bipartite channels: (Gour, 2018)
- ▶ comparison of channels: $d_{A_0} = d_{B_0} = 1$, restrictions on α :
 - ▶ post-processings (degradability)

$$A_1 = A_2, \quad \Phi_1 \stackrel{?}{=} \alpha \circ \Phi_2$$

(Chefles 2009, Buscemi 2012, 2016, ...)

- ▶ comparison of quantum experiments: c-q channels

$$\rho_1, \dots, \rho_k \in \mathcal{S}(B_1), \quad \sigma_1, \dots, \sigma_k \in \mathcal{S}(B_2), \quad \rho_i \stackrel{?}{=} \alpha(\sigma_i), \forall i$$

(Alberti & Uhlmann 1980, Buscemi 2012, Gour et al. 2018)

- ▶ comparison of classical experiments: (Blackwell 1953, Torgersen 1970)
- ▶ more general comparison of classical channels: (Shannon 1958)

Equality case

Preorder: $\Phi_2 \succ \Phi_1 \iff \Phi_1 = \alpha(\Phi_2)$

Equality case

Preorder: $\Phi_2 \succ \Phi_1 \iff \Phi_1 = \alpha(\Phi_2)$

- ▶ formulated as an SDP problem

Equality case

Preorder: $\Phi_2 \succ \Phi_1 \iff \Phi_1 = \alpha(\Phi_2)$

- ▶ formulated as an SDP problem
- ▶ characterized by

Equality case

Preorder: $\Phi_2 \succ \Phi_1 \iff \Phi_1 = \alpha(\Phi_2)$

- ▶ formulated as an SDP problem
- ▶ characterized by
 - ▶ a set of inequalities in (some extension of) conditional min-entropy

Equality case

Preorder: $\Phi_2 \succ \Phi_1 \iff \Phi_1 = \alpha(\Phi_2)$

- ▶ formulated as an SDP problem
- ▶ characterized by
 - ▶ a set of inequalities in (some extension of) conditional min-entropy
 - ▶ risks of procedures in decision problems

Equality case

Preorder: $\Phi_2 \succ \Phi_1 \iff \Phi_1 = \alpha(\Phi_2)$

- ▶ formulated as an SDP problem
- ▶ characterized by
 - ▶ a set of inequalities in (some extension of) **conditional min-entropy**
 - ▶ risks of procedures in decision problems
 - ▶ success probabilities in **hypothesis testing** problems

Deficiency

Let us return to the general case:

- ▶ we define the **deficiency** as

$$\delta_{\mathcal{T}}(\Phi_1 \| \Phi_2) := \min_{\alpha \in \mathcal{T}} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond}$$

here \mathcal{T} is some subset of superchannels.

Deficiency

Let us return to the general case:

- ▶ we define the **deficiency** as

$$\delta_{\mathcal{T}}(\Phi_1 \| \Phi_2) := \min_{\alpha \in \mathcal{T}} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond}$$

here \mathcal{T} is some subset of superchannels.

- ▶ pseudo-distance:

$$\Delta_{\mathcal{T}}(\Phi_1, \Phi_2) := \max\{\delta_{\mathcal{T}}(\Phi_1 \| \Phi_2), \delta_{\mathcal{T}}(\Phi_2 \| \Phi_1)\}$$

Deficiency

Let us return to the general case:

- ▶ we define the **deficiency** as

$$\delta_{\mathcal{T}}(\Phi_1 \| \Phi_2) := \min_{\alpha \in \mathcal{T}} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond}$$

here \mathcal{T} is some subset of superchannels.

- ▶ pseudo-distance:

$$\Delta_{\mathcal{T}}(\Phi_1, \Phi_2) := \max\{\delta_{\mathcal{T}}(\Phi_1 \| \Phi_2), \delta_{\mathcal{T}}(\Phi_2 \| \Phi_1)\}$$

- ▶ equivalence relation:

$$\Phi_1 \sim_{\mathcal{T}} \Phi_2 \iff \Delta_{\mathcal{T}}(\Phi_1, \Phi_2) = 0$$

Deficiency

- ▶ These are extensions of **Le Cam deficiency/distance** for classical statistical experiments: $\mathcal{F}_1 = \{p_\theta, \theta \in \Theta\}$, $\mathcal{F}_2 = \{q_\theta, \theta \in \Theta\}$

$$\delta(\mathcal{F}_1 \| \mathcal{F}_2) = \min_{\alpha} \sup_{\theta} \|p_\theta - \alpha(q_\theta)\|_1$$

- ▶ **Randomization theorem** (Le Cam 1964): deficiency is characterized by comparing risks in decision problems: **informativity**

Randomization theorem for classical channels

Φ_1, Φ_2 - classical channels with equal input spaces: $A_1 = A_2 = A$,
 $\mathcal{T} = \text{post} :=$ set of post-processings

Theorem

Let $\epsilon \geq 0$, Then $\delta_{\text{post}}(\Phi_1 \| \Phi_2) \leq \epsilon$ if and only if:
for any ensemble $\mathcal{E} = \{\lambda_x, p_x\}$ of classical states $p_x \in \mathcal{S}(A)$,

$$P_{\text{succ}}(\Phi_1(\mathcal{E})) \leq P_{\text{succ}}(\Phi_2(\mathcal{E})) + \frac{\epsilon}{2} P_{\text{succ}}(\mathcal{E}),$$

here $\Phi_i(\mathcal{E}) = \{\lambda_x, \Phi_i(p_x)\}$.

Quantum randomization theorems

- ▶ for c-q channels (quantum experiments) (Matsumoto 2010): risks of quantum decision problems

Quantum randomization theorems

- ▶ for c-q channels (quantum experiments) (Matsumoto 2010): risks of quantum decision problems
- ▶ for quantum channels (J 2016): $\delta_{post}(\Phi_1 \parallel \Phi_2) \leq \epsilon \iff$

$$P_{succ}(\Phi_1 \otimes id_R(\mathcal{E})) \leq P_{succ}(\Phi_2 \otimes id_R(\mathcal{E})) + \frac{\epsilon}{2} P_{succ}(\mathcal{E})$$

for any ensemble on AR

Quantum randomization theorems

- ▶ for c-q channels (quantum experiments) (Matsumoto 2010): risks of quantum decision problems
- ▶ for quantum channels (J 2016): $\delta_{post}(\Phi_1 \parallel \Phi_2) \leq \epsilon \iff$

$$P_{succ}(\Phi_1 \otimes id_R(\mathcal{E})) \leq P_{succ}(\Phi_2 \otimes id_R(\mathcal{E})) + \frac{\epsilon}{2} P_{succ}(\mathcal{E})$$

for any ensemble on $AR \iff$

for any $\rho \in \mathcal{S}(AR)$:

$$2^{-H_{min}(R|B_1)_{\rho_1}} \leq 2^{-H_{min}(R|B_2)_{\rho_2}} + \frac{\epsilon}{2} 2^{-H_{min}(R|A)_{\rho}}$$

$\rho_i = (\Phi_i \otimes id_R)(\rho)$, H_{min} - conditional min-entropy

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Goals: characterize $\delta_{\text{sc}}(\Phi_1 \| \Phi_2)$ by

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Goals: characterize $\delta_{\text{sc}}(\Phi_1 \| \Phi_2)$ by

- ▶ inequalities in (extended) conditional min-entropies

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Goals: characterize $\delta_{\text{sc}}(\Phi_1 \| \Phi_2)$ by

- ▶ inequalities in (extended) conditional min-entropies
- ▶ success probabilities in some hypothesis testing schemes

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Goals: characterize $\delta_{\text{sc}}(\Phi_1 \| \Phi_2)$ by

- ▶ inequalities in (extended) conditional min-entropies
- ▶ success probabilities in some hypothesis testing schemes

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Goals: characterize $\delta_{\text{sc}}(\Phi_1 \| \Phi_2)$ by

- ▶ inequalities in (extended) conditional min-entropies
- ▶ success probabilities in some hypothesis testing schemes

Tools:

- ▶ the **duality** of diamond norm and conditional min-entropy (and extensions)

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Goals: characterize $\delta_{\text{sc}}(\Phi_1 \| \Phi_2)$ by

- ▶ inequalities in (extended) conditional min-entropies
- ▶ success probabilities in some hypothesis testing schemes

Tools:

- ▶ the **duality** of diamond norm and conditional min-entropy (and extensions)
 - ▶ comes from **dual SDP** (Chiribella & Ebler 2016, Gour 2018)

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Goals: characterize $\delta_{\text{sc}}(\Phi_1 \| \Phi_2)$ by

- ▶ inequalities in (extended) conditional min-entropies
- ▶ success probabilities in some hypothesis testing schemes

Tools:

- ▶ the **duality** of diamond norm and conditional min-entropy (and extensions)
 - ▶ comes from **dual SDP** (Chiribella & Ebler 2016, Gour 2018)
 - ▶ but also as a **dual norm** (Gutoski 2012, J 2014)

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Goals: characterize $\delta_{\text{sc}}(\Phi_1 \| \Phi_2)$ by

- ▶ inequalities in (extended) conditional min-entropies
- ▶ success probabilities in some hypothesis testing schemes

Tools:

- ▶ the **duality** of diamond norm and conditional min-entropy (and extensions)
 - ▶ comes from **dual SDP** (Chiribella & Ebler 2016, Gour 2018)
 - ▶ but also as a **dual norm** (Gutoski 2012, J 2014)
- ▶ **properties** (monotonicity) of conditional min-entropy

Outline

Φ_1, Φ_2 bipartite quantum channels, $\mathcal{T} = \text{sc} :=$ all superchannels

Goals: characterize $\delta_{\text{sc}}(\Phi_1 \| \Phi_2)$ by

- ▶ inequalities in (extended) conditional min-entropies
- ▶ success probabilities in some hypothesis testing schemes

Tools:


- ▶ the **duality** of diamond norm and conditional min-entropy (and extensions)
 - ▶ comes from **dual SDP** (Chiribella & Ebler 2016, Gour 2018)
 - ▶ but also as a **dual norm** (Gutoski 2012, J 2014)
- ▶ **properties** (monotonicity) of conditional min-entropy
- ▶ **minimax theorem** (Le Cam)

Choi isomorphism

Let $|I_A\rangle\rangle = \sum_i |i\rangle_A |i\rangle_{A'}, \quad A \simeq A'$



Choi isomorphism

Let $|I_A\rangle\rangle = \sum_i |i\rangle_A |i\rangle_{A'}$, $A \simeq A'$





Choi isomorphism

Let $|I_A\rangle\rangle = \sum_i |i\rangle_A |i\rangle_{A'}$, $A \simeq A'$




Choi isomorphism

Let $|I_A\rangle\rangle = \sum_i |i\rangle_A |i\rangle_{A'}$, $A \simeq A'$





Choi isomorphism:




Choi isomorphism

Let $|I_A\rangle\rangle = \sum_i |i\rangle_A |i\rangle_{A'}$, $A \simeq A'$



the inverse:



Ordered spaces of hermitian maps

- ▶ $\mathcal{L} = \mathcal{L}(A \rightarrow B) \equiv$ the space of hermitian-preserving maps

$$\phi : \mathcal{B}(A) \rightarrow \mathcal{B}(B), \quad \phi(Y^*) = \phi(Y)^*$$

Ordered spaces of hermitian maps

- ▶ $\mathcal{L} = \mathcal{L}(A \rightarrow B) \equiv$ the space of hermitian-preserving maps

$$\phi : \mathcal{B}(A) \rightarrow \mathcal{B}(B), \quad \phi(Y^*) = \phi(Y)^*$$

- ▶ $\mathcal{L}^+ = \mathcal{L}^+(A \rightarrow B) \equiv$ the cone of **completely positive maps**

Ordered spaces of hermitian maps

- ▶ $\mathcal{L} = \mathcal{L}(A \rightarrow B) \equiv$ the space of hermitian-preserving maps

$$\phi : \mathcal{B}(A) \rightarrow \mathcal{B}(B), \quad \phi(Y^*) = \phi(Y)^*$$

- ▶ $\mathcal{L}^+ = \mathcal{L}^+(A \rightarrow B) \equiv$ the cone of **completely positive maps**
- ▶ $(\mathcal{L}, \mathcal{L}^+)$ is an **ordered vector space**

Ordered spaces of hermitian maps

- ▶ $\mathcal{L} = \mathcal{L}(A \rightarrow B) \equiv$ the space of hermitian-preserving maps

$$\phi : \mathcal{B}(A) \rightarrow \mathcal{B}(B), \quad \phi(Y^*) = \phi(Y)^*$$

- ▶ $\mathcal{L}^+ = \mathcal{L}^+(A \rightarrow B) \equiv$ the cone of **completely positive maps**
- ▶ $(\mathcal{L}, \mathcal{L}^+)$ is an **ordered vector space**
- ▶ **Choi isomorphism:**

$$(\mathcal{L}, \mathcal{L}^+) \simeq (\mathcal{B}_h(BA'), \mathcal{B}(BA')^+)$$

The trace

For $\phi \in \mathcal{L}(A \rightarrow A)$, put $\tau(\phi) := \langle\langle I_A | C_\phi | I_A \rangle\rangle$

The trace

For $\phi \in \mathcal{L}(A \rightarrow A)$, put $\tau(\phi) := \langle\langle I_A | C_\phi | I_A \rangle\rangle$

$$\tau(\phi) = \text{Diagram 1} = \text{Diagram 2}$$

Diagram 1: A box labeled ϕ with two horizontal wires passing through it. The top wire is labeled A on both sides of the box. The bottom wire is labeled A' on both sides of the box. The wires are connected at their ends by semi-circular arcs, forming a closed loop.

Diagram 2: A vertical rectangle labeled C_ϕ . Two horizontal wires enter from the left and exit to the right. The top wire is labeled A on the right side. The bottom wire is labeled A' on the right side. The wires are connected at their right ends by a semi-circular arc, forming a closed loop.

The trace

For $\phi \in \mathcal{L}(A \rightarrow A)$, put $\tau(\phi) := \langle\langle I_A | C_\phi | I_A \rangle\rangle$

$$\tau(\phi) = \text{diagram} = \text{diagram}$$

The diagram shows the trace of a linear map ϕ as a linear map. On the left, a box labeled ϕ has two horizontal wires passing through it. The top wire is labeled A on both sides of the box, and the bottom wire is labeled A' on both sides. The wires are connected at their ends by semi-circular loops, forming a closed loop. This is followed by an equals sign and a second diagram. The second diagram shows a box labeled C_ϕ with two horizontal wires passing through it. The top wire is labeled A on both sides of the box, and the bottom wire is labeled A' on both sides. The wires are connected at their ends by semi-circular loops, forming a closed loop.


the trace of ϕ as a linear map

The trace

Partial trace: for $\phi \in \mathcal{L}(AA_0 \rightarrow BA_0)$


The trace

Partial trace: for $\phi \in \mathcal{L}(AA_0 \rightarrow BA_0)$

$$\tau_{A_0}(\phi) =$$


The trace


Partial trace: for $\phi \in \mathcal{L}(AA_0 \rightarrow BA_0)$

$$\tau_{A_0}(\phi) =$$


- ▶ if $\phi \in \mathcal{L}^+(AA_0 \rightarrow BA_0)$ then $\tau_{A_0}(\phi) \in \mathcal{L}^+(A \rightarrow B)$

The trace


Partial trace: for $\phi \in \mathcal{L}(AA_0 \rightarrow BA_0)$

$$\tau_{A_0}(\phi) =$$


- ▶ if $\phi \in \mathcal{L}^+(AA_0 \rightarrow BA_0)$ then $\tau_{A_0}(\phi) \in \mathcal{L}^+(A \rightarrow B)$
- ▶ does not map channels to channels

The trace

Partial trace: for $\phi \in \mathcal{L}(AA_0 \rightarrow BA_0)$

$$\tau_{A_0}(\phi) =$$


- ▶ if $\phi \in \mathcal{L}^+(AA_0 \rightarrow BA_0)$ then $\tau_{A_0}(\phi) \in \mathcal{L}^+(A \rightarrow B)$
- ▶ does not map channels to channels
- ▶ $\tau(\phi \circ (\alpha \otimes id)) = \tau(\tau_{A_0}(\phi) \circ \alpha)$

The ordered dual of $\mathcal{L}(A \rightarrow B)$

For $\mathcal{L} = \mathcal{L}(A \rightarrow B)$, we represent

$$\mathcal{L}^* \equiv \mathcal{L}(B \rightarrow A), \quad (\mathcal{L}^+)^* \equiv \mathcal{L}^+(B \rightarrow A)$$

The ordered dual of $\mathcal{L}(A \rightarrow B)$

For $\mathcal{L} = \mathcal{L}(A \rightarrow B)$, we represent

$$\mathcal{L}^* \equiv \mathcal{L}(B \rightarrow A), \quad (\mathcal{L}^+)^* \equiv \mathcal{L}^+(B \rightarrow A)$$

with duality

$$\langle \psi, \phi \rangle = \tau(\psi \circ \phi) = \tau(\phi \circ \psi) = \text{Tr } C_\phi C_{\psi^*} :$$


The ordered dual of $\mathcal{L}(A \rightarrow B)$

For $\mathcal{L} = \mathcal{L}(A \rightarrow B)$, we represent

$$\mathcal{L}^* \equiv \mathcal{L}(B \rightarrow A), \quad (\mathcal{L}^+)^* \equiv \mathcal{L}^+(B \rightarrow A)$$

with duality

$$\langle \psi, \phi \rangle = \tau(\psi \circ \phi) = \tau(\phi \circ \psi) = \text{Tr } C_\phi C_{\psi^*} :$$



Diamond norm and conditional min-entropy

For $\phi \in \mathcal{L} = \mathcal{L}(A \rightarrow B)$, the diamond norm is defined as

$$\|\phi\|_{\diamond} = \sup_{\rho \in \mathcal{S}(AA')} \|\phi \otimes id_{A'}(\rho)\|_1$$

Diamond norm and conditional min-entropy

For $\phi \in \mathcal{L} = \mathcal{L}(A \rightarrow B)$, the diamond norm is defined as

$$\|\phi\|_{\diamond} = \sup_{\rho \in \mathcal{S}(AA')} \|\phi \otimes id_{A'}(\rho)\|_1$$

- ▶ distinguishability norm for **channels** $\mathcal{B}(A) \rightarrow \mathcal{B}(B)$

Diamond norm and conditional min-entropy

For $\phi \in \mathcal{L} = \mathcal{L}(A \rightarrow B)$, the diamond norm is defined as

$$\|\phi\|_{\diamond} = \sup_{\rho \in \mathcal{S}(AA')} \|\phi \otimes id_{A'}(\rho)\|_1$$

- ▶ distinguishability norm for **channels** $\mathcal{B}(A) \rightarrow \mathcal{B}(B)$
- ▶ constructed from the convex structure of the set of channels

Diamond norm and conditional min-entropy

Let

- ▶ $\mathcal{C} = \mathcal{C}(A \rightarrow B) \subset \mathcal{L}^+$ the set of channels

Diamond norm and conditional min-entropy

Let

- ▶ $\mathcal{C} = \mathcal{C}(A \rightarrow B) \subset \mathcal{L}^+$ the set of channels
- ▶ \mathcal{C} is an affine section of \mathcal{L}^+

Diamond norm and conditional min-entropy

Let

- ▶ $\mathcal{C} = \mathcal{C}(A \rightarrow B) \subset \mathcal{L}^+$ the set of channels
- ▶ \mathcal{C} is an affine section of \mathcal{L}^+
- ▶ dual section:

$$\tilde{\mathcal{C}} := \{\gamma \in (\mathcal{L}^+)^*, \langle \gamma, \alpha \rangle = 1, \forall \alpha \in \mathcal{C}\}$$

Diamond norm and conditional min-entropy

Let

- ▶ $\mathcal{C} = \mathcal{C}(A \rightarrow B) \subset \mathcal{L}^+$ the set of channels
- ▶ \mathcal{C} is an affine section of \mathcal{L}^+
- ▶ dual section:

$$\tilde{\mathcal{C}} := \{\gamma \in (\mathcal{L}^+)^*, \langle \gamma, \alpha \rangle = 1, \forall \alpha \in \mathcal{C}\}$$

Diamond norm and conditional min-entropy

Let

- ▶ $\mathcal{C} = \mathcal{C}(A \rightarrow B) \subset \mathcal{L}^+$ the set of channels
- ▶ \mathcal{C} is an affine section of \mathcal{L}^+
- ▶ dual section:

$$\tilde{\mathcal{C}} := \{\gamma \in (\mathcal{L}^+)^*, \langle \gamma, \alpha \rangle = 1, \forall \alpha \in \mathcal{C}\}$$

Dual expressions:

$$\begin{aligned}\|\phi\|_{\diamond} &= \min_{\alpha \in \mathcal{C}} \min\{\lambda > 0, -\lambda\alpha \leq \phi \leq \lambda\alpha\} \\ &= \max_{\gamma \in \tilde{\mathcal{C}}} \max_{-\gamma \leq \eta \leq \gamma} \langle \eta, \phi \rangle\end{aligned}$$

Diamond norm and conditional min-entropy

The dual norm: for $\psi \in \mathcal{L}^*$,

$$\begin{aligned}\|\psi\|^\diamond &= \min_{\gamma \in \tilde{\mathcal{C}}} \min\{\lambda > 0, -\lambda\gamma \leq \psi \leq \lambda\gamma\} \\ &= \max_{\alpha \in \mathcal{C}} \max_{-\alpha \leq \xi \leq \alpha} \langle \psi, \xi \rangle\end{aligned}$$

Diamond norm and conditional min-entropy

The dual norm: for $\psi \in \mathcal{L}^*$,


$$\begin{aligned}\|\psi\|^\diamond &= \min_{\gamma \in \tilde{\mathcal{C}}} \min\{\lambda > 0, -\lambda\gamma \leq \psi \leq \lambda\gamma\} \\ &= \max_{\alpha \in \mathcal{C}} \max_{-\alpha \leq \xi \leq \alpha} \langle \psi, \xi \rangle\end{aligned}$$

For cp maps: if $\psi \in (\mathcal{L}^+)^*$:

$$\begin{aligned}\|\psi\|^\diamond &= \min_{\gamma \in \tilde{\mathcal{C}}} \min\{\lambda > 0, \psi \leq \lambda\gamma\} \\ &= \max_{\alpha \in \mathcal{C}} \langle \psi, \alpha \rangle\end{aligned}$$

Diamond norm and conditional min-entropy


- maps in $\tilde{\mathcal{C}}$:



$$\sigma \in \mathcal{S}(A)$$

Diamond norm and conditional min-entropy

- maps in $\tilde{\mathcal{C}}$:




- in Choi representation:

$$\tilde{\mathcal{C}} = \{\sigma_A \otimes I_B, \sigma_A \in \mathcal{S}(A)\}$$

Diamond norm and conditional min-entropy

- maps in $\tilde{\mathcal{C}}$:



- in Choi representation:


$$\tilde{\mathcal{C}} = \{\sigma_A \otimes I_B, \sigma_A \in \mathcal{S}(A)\}$$

- for $\psi \in (\mathcal{L}^+)^*$, $\rho = C_\psi$:

$$\|\psi\|^\diamond = \min_{\sigma_A \in \mathcal{S}(A)} \min\{\lambda > 0, \rho \leq \lambda \sigma_A \otimes I_B\} = 2^{-H_{\min}(B|A)_\rho}$$

Diamond norm and conditional min-entropy

- maps in $\tilde{\mathcal{C}}$:



- in Choi representation:

$$\tilde{\mathcal{C}} = \{\sigma_A \otimes I_B, \sigma_A \in \mathcal{S}(A)\}$$

- for $\psi \in (\mathcal{L}^+)^*$, $\rho = C_\psi$:

$$\|\psi\|^\diamond = \min_{\sigma_A \in \mathcal{S}(A)} \min\{\lambda > 0, \rho \leq \lambda \sigma_A \otimes I_B\} = 2^{-H_{\min}(B|A)_\rho}$$

- conditional min-entropy: $H_{\min}(B|A)_\rho$

Diamond norm and conditional min-entropy


Dual expression:

$$\|\psi\|^\diamond = \max_{\alpha \in \mathcal{C}} \langle \psi, \alpha \rangle = \max_{\alpha \in \mathcal{C}} \langle \langle I_B | (\alpha \otimes id)(\rho) | I_B \rangle \rangle$$

Diamond norm and conditional min-entropy

Dual expression:


$$\|\psi\|^\diamond = \max_{\alpha \in \mathcal{C}} \langle \psi, \alpha \rangle = \max_{\alpha \in \mathcal{C}} \langle\langle I_B | (\alpha \otimes id)(\rho) | I_B \rangle\rangle$$



Diamond norm and conditional min-entropy

Dual expression:

$$\|\psi\|^\diamond = \max_{\alpha \in \mathcal{C}} \langle \psi, \alpha \rangle = \max_{\alpha \in \mathcal{C}} \langle\langle I_B | (\alpha \otimes id)(\rho) | I_B \rangle\rangle$$



operational interpretation of $H_{\min}(B|A)_\rho$ (König et al., 2009):

(up to d_B) the largest fidelity with maximally entangled state, that can be obtained by applying a channel on A

The 2-diamond norm and conditional 2-min-entropy

Let now

$$\blacktriangleright \mathcal{L} = \mathcal{L}(A_1 A_2 \rightarrow B_1 B_2)$$

The 2-diamond norm and conditional 2-min-entropy


Let now

- ▶ $\mathcal{L} = \mathcal{L}(A_1 A_2 \rightarrow B_1 B_2)$
- ▶ $\mathcal{C}_2 = \mathcal{C}_2(A_1 \rightarrow B_1, A_2 \rightarrow B_2) \equiv$ set of **superchannels**:

The 2-diamond norm and conditional 2-min-entropy

Let now


- ▶ $\mathcal{L} = \mathcal{L}(A_1 A_2 \rightarrow B_1 B_2)$
- ▶ $\mathcal{C}_2 = \mathcal{C}_2(A_1 \rightarrow B_1, A_2 \rightarrow B_2) \equiv$ set of **superchannels**:



The 2-diamond norm and conditional 2-min-entropy

Let now

- ▶ $\mathcal{L} = \mathcal{L}(A_1 A_2 \rightarrow B_1 B_2)$
- ▶ $\mathcal{C}_2 = \mathcal{C}_2(A_1 \rightarrow B_1, A_2 \rightarrow B_2) \equiv$ set of **superchannels**:




- ▶ \mathcal{C}_2 is an affine section of \mathcal{L}^+

The 2-diamond norm and conditional 2-min-entropy

The dual section $\tilde{\mathcal{C}}_2$: set of superchannels


The 2-diamond norm and conditional 2-min-entropy

The dual section $\tilde{\mathcal{C}}_2$: set of superchannels



The 2-diamond norm and conditional 2-min-entropy

The dual section $\tilde{\mathcal{C}}_2$: set of superchannels



σ is a state, γ a channel

The 2-diamond norm and conditional 2-min-entropy

- ▶ we can define a pair of dual norms in $\mathcal{L}, \mathcal{L}^*$ as before:

$$\begin{aligned}\|\phi\|_{2\Diamond} &= \min_{\alpha \in \mathcal{C}_2} \min\{\lambda > 0, -\lambda\alpha \leq \phi \leq \lambda\alpha\} \\ &= \max_{\gamma \in \tilde{\mathcal{C}}_2} \max_{-\gamma \leq \eta \leq \gamma} \langle \eta, \phi \rangle \\ \|\psi\|^{2\Diamond} &= \min_{\gamma \in \tilde{\mathcal{C}}_2} \min\{\lambda > 0, -\lambda\gamma \leq \psi \leq \lambda\gamma\} \\ &= \max_{\alpha \in \mathcal{C}_2} \max_{-\alpha \leq \xi \leq \alpha} \langle \psi, \xi \rangle\end{aligned}$$

The 2-diamond norm and conditional 2-min-entropy

- ▶ we can define a pair of dual norms in \mathcal{L} , \mathcal{L}^* as before:

$$\begin{aligned}\|\phi\|_{2\Diamond} &= \min_{\alpha \in \mathcal{C}_2} \min\{\lambda > 0, -\lambda\alpha \leq \phi \leq \lambda\alpha\} \\ &= \max_{\gamma \in \tilde{\mathcal{C}}_2} \max_{-\gamma \leq \eta \leq \gamma} \langle \eta, \phi \rangle \\ \|\psi\|^{2\Diamond} &= \min_{\gamma \in \tilde{\mathcal{C}}_2} \min\{\lambda > 0, -\lambda\gamma \leq \psi \leq \lambda\gamma\} \\ &= \max_{\alpha \in \mathcal{C}_2} \max_{-\alpha \leq \xi \leq \alpha} \langle \psi, \xi \rangle\end{aligned}$$

- ▶ $\|\cdot\|_{2\Diamond}$ is a distinguishability norm for superchannels

The 2-diamond norm and conditional 2-min-entropy

- ▶ we can define a pair of dual norms in \mathcal{L} , \mathcal{L}^* as before:

$$\begin{aligned}\|\phi\|_{2\Diamond} &= \min_{\alpha \in \mathcal{C}_2} \min\{\lambda > 0, -\lambda\alpha \leq \phi \leq \lambda\alpha\} \\ &= \max_{\gamma \in \tilde{\mathcal{C}}_2} \max_{-\gamma \leq \eta \leq \gamma} \langle \eta, \phi \rangle \\ \|\psi\|^{2\Diamond} &= \min_{\gamma \in \tilde{\mathcal{C}}_2} \min\{\lambda > 0, -\lambda\gamma \leq \psi \leq \lambda\gamma\} \\ &= \max_{\alpha \in \mathcal{C}_2} \max_{-\alpha \leq \xi \leq \alpha} \langle \psi, \xi \rangle\end{aligned}$$

- ▶ $\|\cdot\|_{2\Diamond}$ is a distinguishability norm for superchannels
- ▶ from $\|\cdot\|^{2\Diamond}$, we obtain the **conditional 2-min-entropy**

The 2-diamond norm and conditional 2-min-entropy


For $\psi \in (\mathcal{L}^+)^*$, $\rho = C_\psi$:

$$\begin{aligned}\|\psi\|^{2\Diamond} &= \min_{\sigma, \gamma} \min\{\lambda > 0, \psi \leq \lambda \sigma * \gamma \otimes \text{Tr}_{B_2}\} = 2^{-H_{\min}^{(2)}(B|A)_\rho} \\ &= \max_{\alpha \in \mathcal{C}_2} \langle \alpha, \psi \rangle = \max_{\alpha \in \mathcal{C}_2} \langle \langle I_{B_1 B_2} | (\alpha \otimes id)(\rho) | I_{B_1 B_2} \rangle \rangle\end{aligned}$$

The 2-diamond norm and conditional 2-min-entropy

For $\psi \in (\mathcal{L}^+)^*$, $\rho = C_\psi$:

$$\begin{aligned} \|\psi\|^{2_\diamond} &= \min_{\sigma, \gamma} \min\{\lambda > 0, \psi \leq \lambda \sigma * \gamma \otimes \text{Tr}_{B_2}\} = 2^{-H_{\min}^{(2)}(B|A)_\rho} \\ &= \max_{\alpha \in \mathcal{C}_2} \langle \alpha, \psi \rangle = \max_{\alpha \in \mathcal{C}_2} \langle\langle I_{B_1 B_2} | (\alpha \otimes id)(\rho) | I_{B_1 B_2} \rangle\rangle \end{aligned}$$



The 2-diamond norm and conditional 2-min-entropy

For $\psi \in (\mathcal{L}^+)^*$, $\rho = C_\psi$:

$$\begin{aligned}\|\psi\|^{2\Diamond} &= \min_{\sigma, \gamma} \min\{\lambda > 0, \psi \leq \lambda \sigma * \gamma \otimes \text{Tr}_{B_2}\} = 2^{-H_{\min}^{(2)}(B|A)_\rho} \\ &= \max_{\alpha \in \mathcal{C}_2} \langle \alpha, \psi \rangle = \max_{\alpha \in \mathcal{C}_2} \langle\langle I_{B_1 B_2} | (\alpha \otimes id)(\rho) | I_{B_1 B_2} \rangle\rangle\end{aligned}$$

Useful properties of $H_{\min}^{(2)}$: (Gour, 2018)

- ▶ monotonicity for any superchannel $\Theta \in \mathcal{C}_2(B_3 \rightarrow B_1, A_1 \rightarrow A_3)$:

$$\|\phi\|^{2\Diamond} \geq \|\Theta(\phi)\|^{2\Diamond}$$

- ▶ additivity

$$\|\phi \otimes \psi\|^{2\Diamond} = \|\phi\|^{2\Diamond} \|\psi\|^{2\Diamond}$$

Comparison of bipartite channels

We compute the deficiency $\delta_{sc}(\Phi_1 \parallel \Phi_2)$:

$$\min_{\alpha \in \mathcal{C}_2} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond} =$$

Comparison of bipartite channels

We compute the deficiency $\delta_{sc}(\Phi_1 \parallel \Phi_2)$:

$$\min_{\alpha \in \mathcal{C}_2} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond} = \min_{\alpha \in \mathcal{C}_2} \max_{\|\gamma\|_{\diamond} \leq 1} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle$$

Comparison of bipartite channels

We compute the deficiency $\delta_{sc}(\Phi_1 \parallel \Phi_2)$:

$$\min_{\alpha \in \mathcal{C}_2} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond} = 2 \min_{\alpha \in \mathcal{C}_2} \max_{\substack{\|\gamma\|_{\diamond} \leq 1, \\ \gamma \geq 0}} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle$$

Comparison of bipartite channels

We compute the deficiency $\delta_{sc}(\Phi_1 \parallel \Phi_2)$:

$$\min_{\alpha \in \mathcal{C}_2} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond} = 2 \min_{\alpha \in \mathcal{C}_2} \max_{\substack{\|\gamma\|_{\diamond} \leq 1, \\ \gamma \geq 0}} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle$$

(minimax thm.) $= 2 \max_{\gamma} \min_{\alpha} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle$

Comparison of bipartite channels

We compute the deficiency $\delta_{sc}(\Phi_1 \parallel \Phi_2)$:

$$\begin{aligned}\min_{\alpha \in \mathcal{C}_2} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond} &= 2 \min_{\alpha \in \mathcal{C}_2} \max_{\substack{\|\gamma\|_{\diamond} \leq 1, \\ \gamma \geq 0}} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle \\ (\text{minimax thm.}) &= 2 \max_{\gamma} \min_{\alpha} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle \\ &\leq 2 \max_{\gamma} \left(\langle \gamma, \Phi_1 \rangle - \max_{\alpha \in \mathcal{C}_2} \langle \gamma, \alpha(\Phi_2) \rangle \right)\end{aligned}$$

Comparison of bipartite channels

We compute the deficiency $\delta_{sc}(\Phi_1 \parallel \Phi_2)$:

$$\begin{aligned}\min_{\alpha \in \mathcal{C}_2} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond} &= 2 \min_{\alpha \in \mathcal{C}_2} \max_{\substack{\|\gamma\|_{\diamond} \leq 1, \\ \gamma \geq 0}} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle \\ (\text{minimax thm.}) &= 2 \max_{\gamma} \min_{\alpha} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle \\ &\leq 2 \max_{\gamma} \left(\max_{\beta \in \mathcal{C}_2} \langle \gamma, \beta(\Phi_1) \rangle - \max_{\alpha \in \mathcal{C}_2} \langle \gamma, \alpha(\Phi_2) \rangle \right)\end{aligned}$$

Comparison of bipartite channels

We compute the deficiency $\delta_{sc}(\Phi_1 \parallel \Phi_2)$:


$$\begin{aligned}\min_{\alpha \in \mathcal{C}_2} \|\Phi_1 - \alpha(\Phi_2)\|_{\diamond} &= 2 \min_{\alpha \in \mathcal{C}_2} \max_{\substack{\|\gamma\|_{\diamond} \leq 1, \\ \gamma \geq 0}} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle \\ (\text{minimax thm.}) &= 2 \max_{\gamma} \min_{\alpha} \langle \gamma, \Phi_1 - \alpha(\Phi_2) \rangle \\ &\leq 2 \max_{\gamma} \left(\max_{\beta \in \mathcal{C}_2} \langle \gamma, \beta(\Phi_1) \rangle - \max_{\alpha \in \mathcal{C}_2} \langle \gamma, \alpha(\Phi_2) \rangle \right) \\ &\leq \epsilon ?\end{aligned}$$

Comparison of bipartite channels

$$\max_{\alpha \in \mathcal{C}_2} \langle \gamma, \alpha(\Phi_i) \rangle = ?$$


Comparison of bipartite channels

$$\max_{\alpha \in \mathcal{C}_2} \langle \gamma, \alpha(\Phi_i) \rangle = ?$$




Comparison of bipartite channels

$$\max_{\alpha \in \mathcal{C}_2} \langle \gamma, \alpha(\Phi_i) \rangle = ?$$



Comparison of bipartite channels


$$\max_{\alpha \in \mathcal{C}_2} \langle \gamma, \alpha(\Phi_i) \rangle = ?$$



$$= \tau_{B_0}(\gamma * \Phi_i)$$


Comparison of bipartite channels

$$\max_{\alpha \in \mathcal{C}_2} \langle \gamma, \alpha(\Phi_i) \rangle = ?$$



Comparison of bipartite channels

$$\max_{\alpha \in \mathcal{C}_2} \langle \gamma, \alpha(\Phi_i) \rangle = \max_{\alpha \in \mathcal{C}_2} \langle \tau_{B_0}(\gamma * \Phi_i), \alpha \rangle = \|\tau_{B_0}(\gamma * \Phi_i)\|^{\diamond}$$



Comparison of bipartite channels

Theorem

Let $\epsilon \geq 0$. Then $\delta_{sc}(\Phi_1 \| \Phi_2) \leq \epsilon$ if and only if for all systems A_3, B_3 and all $\gamma \in \mathcal{L}^+(B_3 B_0 \rightarrow A_3 A_0)$ we have

$$\|\tau_{B_0}(\gamma * \Phi_1)\|^{2\diamond} \leq \|\tau_{B_0}(\gamma * \Phi_2)\|^{2\diamond} + \frac{\epsilon}{2} \|\gamma\|^\diamond.$$


Comparison of bipartite channels

Theorem

Let $\epsilon \geq 0$. Then $\delta_{sc}(\Phi_1 \| \Phi_2) \leq \epsilon$ if and only if for all systems A_3, B_3 and all $\gamma \in \mathcal{L}^+(B_3 B_0 \rightarrow A_3 A_0)$ we have

$$\|\tau_{B_0}(\gamma * \Phi_1)\|^{2\diamond} \leq \|\tau_{B_0}(\gamma * \Phi_2)\|^{2\diamond} + \frac{\epsilon}{2} \|\gamma\|^\diamond.$$

where




Comparison of bipartite channels

Theorem

Let $\epsilon \geq 0$. Then $\delta_{sc}(\Phi_1 \| \Phi_2) \leq \epsilon$ if and only if for all systems A_3, B_3 and all $\gamma \in \mathcal{L}^+(B_3 B_0 \rightarrow A_3 A_0)$ we have

$$\|\tau_{B_0}(\gamma * \Phi_1)\|^{2\diamond} \leq \|\tau_{B_0}(\gamma * \Phi_2)\|^{2\diamond} + \frac{\epsilon}{2} \|\gamma\|^\diamond.$$

where



We can restrict to $A_3 \simeq A_1$ and $B_3 \simeq B_1$.

Conditional min-entropy and guessing probabilities

For an ensemble $\mathcal{E} = \{\lambda_x, \rho_x\}$, $\rho_x \in \mathcal{S}(A)$, let

$$\phi_{\mathcal{E}} \in \mathcal{L}^+(B \rightarrow A), \quad C_{\phi_{\mathcal{E}}} = \rho_{\mathcal{E}} := \sum_x |x\rangle\langle x| \otimes \lambda_x \rho_x$$

$$\phi_{\mathcal{E}} = \begin{array}{c} x \\ \hline \boxed{\lambda_x \rho_x} \\ \hline A \end{array}$$

-classical-to-quantum map

Conditional min-entropy and guessing probabilities

For an ensemble $\mathcal{E} = \{\lambda_x, \rho_x\}$, $\rho_x \in \mathcal{S}(A)$, let

$$\phi_{\mathcal{E}} \in \mathcal{L}^+(B \rightarrow A), \quad C_{\phi_{\mathcal{E}}} = \rho_{\mathcal{E}} := \sum_x |x\rangle\langle x| \otimes \lambda_x \rho_x$$

$$\phi_{\mathcal{E}} = \begin{array}{c} x \\ \hline \boxed{\lambda_x \rho_x} \\ \hline A \end{array}$$

-classical-to-quantum map

Optimal success probability (König et al. 2009)

$$P_{succ}(\mathcal{E}) = \max_M \sum_x \lambda_x \text{Tr}[\rho_x M_x] = \|\phi_{\mathcal{E}}\|^\diamond = 2^{-H_{min}(X|A)_{\rho_{\mathcal{E}}}}$$

Conditional min-entropy and guessing probabilities

Let $\gamma \in \mathcal{L}^+(R \rightarrow A)$, $\rho = C_\gamma \in \mathcal{S}(AR)$. We produce an ensemble

$$\mathcal{E}_\rho = \left\{ \frac{1}{d_R^2}, \rho_x \right\}, \quad \rho_x = (id_A \otimes \mathcal{U}_x^R)(\rho) \in \mathcal{S}(AR),$$

Conditional min-entropy and guessing probabilities

Let $\gamma \in \mathcal{L}^+(R \rightarrow A)$, $\rho = C_\gamma \in \mathcal{S}(AR)$. We produce an ensemble

$$\mathcal{E}_\rho = \left\{ \frac{1}{d_R^2}, \rho_x \right\}, \quad \rho_x = (id_A \otimes \mathcal{U}_x^R)(\rho) \in \mathcal{S}(AR),$$

where

$$\mathcal{U}_x^R \equiv \text{generalized Pauli unitaries on } \mathcal{H}_R.$$

Conditional min-entropy and guessing probabilities

Let $\gamma \in \mathcal{L}^+(R \rightarrow A)$, $\rho = C_\gamma \in \mathcal{S}(AR)$. We produce an ensemble

$$\mathcal{E}_\rho = \left\{ \frac{1}{d_R^2}, \rho_x \right\}, \quad \rho_x = (id_A \otimes \mathcal{U}_x^R)(\rho) \in \mathcal{S}(AR),$$

where

$$\mathcal{U}_x^R \equiv \text{generalized Pauli unitaries on } \mathcal{H}_R.$$


Then we have

$$P_{\text{succ}}(\mathcal{E}_\rho) = \frac{1}{d_R} \|\gamma\|^\diamond = \frac{1}{d_R} 2^{-H_{\min}(R|A)_\rho}$$

Conditional min-entropy and guessing probabilities

Channel discrimination problem:

- ▶ an ensemble of channels $\mathcal{E}_R = \{\frac{1}{d_R^2}, \mathcal{U}_x^R\}$
- ▶ testers (PPOVMs) with input state ρ :



- ▶ success probability:

$$P_{succ}(\mathcal{E}_R, \rho) := \max_M \sum_x \text{Tr} [(id \otimes \mathcal{U}_x^R)(\rho) M_x] = \frac{1}{d_R} 2^{-H_{min}(R|A)_\rho}$$

Conditional min-entropy and guessing probabilities

For any ensemble $\mathcal{E} = \{\frac{1}{d_R}, \Psi_x\}_{x=1}^{d_R^2}$ of **unital** channels:


$$P_{succ}(\mathcal{E}, \rho) \leq \frac{1}{d_R} 2^{-H_{min}(R|A)_\rho}$$

Conditional min-entropy and guessing probabilities

For any ensemble $\mathcal{E} = \{\frac{1}{d_R^2}, \Psi_x\}_{x=1}^{d_R^2}$ of **unital** channels:

$$P_{succ}(\mathcal{E}, \rho) \leq \frac{1}{d_R} 2^{-H_{min}(R|A)_\rho}$$

For a pair of quantum channels $\Phi_i : A \rightarrow B_i$, $\delta_{post}(\Phi_1 \| \Phi_2)$ can be characterized by comparing testers of the form




for this type of tasks, for any $\rho \in \mathcal{S}(AR)$.

Comparison of bipartite channels by guessing probabilities

A more complicated situation - we maximize over α_1, α_2 :


Comparison of bipartite channels by guessing probabilities

A more complicated situation - we maximize over α_1, α_2 :




Comparison of bipartite channels by guessing probabilities

A more complicated situation - we maximize over α_1, α_2 :




Comparison of bipartite channels by guessing probabilities

A more complicated situation - we maximize over α_1, α_2 :




Comparison of bipartite channels by guessing probabilities

A more complicated situation - we maximize over α_1, α_2 :




Comparison of bipartite channels by guessing probabilities

A more complicated situation - we maximize over α_1, α_2 :



Comparison of bipartite channels by guessing probabilities

A more complicated situation - we maximize over α_1, α_2 :




$$= \frac{1}{d_{B_3} d_{B_0}} \sum_{x,y} \text{Tr} [\rho_{xy} (\alpha_1 * \Phi_i)^* (M_x \otimes B_y)]$$

Comparison of bipartite channels by guessing probabilities

Let $\mathcal{E} = \{\lambda_{xy}, \rho_{xy}\}$ be an ensemble on $B_3 A_3 A_0 B_0$ and $\{N_y\}$ a POVM on $B_0 B'_0$. Consider the following scheme:

Comparison of bipartite channels by guessing probabilities


Let $\mathcal{E} = \{\lambda_{xy}, \rho_{xy}\}$ be an ensemble on $B_3 A_3 A_0 B_0$ and $\{N_y\}$ a POVM on $B_0 B'_0$. Consider the following scheme:



Comparison of bipartite channels by guessing probabilities

Let $\mathcal{E} = \{\lambda_{xy}, \rho_{xy}\}$ be an ensemble on $B_3 A_3 A_0 B_0$ and $\{N_y\}$ a POVM on $B_0 B'_0$. Consider the following scheme:


1. **pre-processing**: pick and apply any $\alpha \in \mathcal{C}(A_3 \rightarrow R A_i)$;



Comparison of bipartite channels by guessing probabilities

Let $\mathcal{E} = \{\lambda_{xy}, \rho_{xy}\}$ be an ensemble on $B_3 A_3 A_0 B_0$ and $\{N_y\}$ a POVM on $B_0 B'_0$. Consider the following scheme:


1. pre-processing: pick and apply any $\alpha \in \mathcal{C}(A_3 \rightarrow R A_i)$;
2. apply Φ_i ;



Comparison of bipartite channels by guessing probabilities

Let $\mathcal{E} = \{\lambda_{xy}, \rho_{xy}\}$ be an ensemble on $B_3 A_3 A_0 B_0$ and $\{N_y\}$ a POVM on $B_0 B'_0$. Consider the following scheme:

1. **pre-processing**: pick and apply any $\alpha \in \mathcal{C}(A_3 \rightarrow R A_i)$;
2. apply Φ_i ;
3. **measurement**: pick any POVM $\{M_x\}$ and measure $M \otimes N$.



Comparison of bipartite channels by guessing probabilities

Let $\mathcal{E} = \{\lambda_{xy}, \rho_{xy}\}$ be an ensemble on $B_3 A_3 A_0 B_0$ and $\{N_y\}$ a POVM on $B_0 B'_0$. Consider the following scheme:

1. **pre-processing**: pick and apply any $\alpha \in \mathcal{C}(A_3 \rightarrow R A_i)$;
2. apply Φ_i ;
3. **measurement**: pick any POVM $\{M_x\}$ and measure $M \otimes N$.

The optimal success probability is

$$P_{succ}(\mathcal{E}, \Phi_i, N) := \max_{\alpha, M} P_{succ}(\mathcal{E}, (\Phi_i * \alpha)^*(M \otimes N))$$

Comparison of bipartite channels: guessing probabilities

Theorem

$\delta_{sc}(\Phi_1 \| \Phi_2) \leq \epsilon$ if and only if for any A_3, B_3 , any POVM N on $B_0 B'_0$ and any ensemble $\mathcal{E} = \{\lambda_{xy}, \rho_{xy}\}$, on $B_3 A_3 A_0 B'_0$, we have

$$P_{succ}(\mathcal{E}, \Phi_1, N) \leq P_{succ}(\mathcal{E}, \Phi_2, N) + \frac{\epsilon}{2} P_{succ}(\mathcal{E})$$

We may restrict to $A_3 \simeq A_1, B_3 \simeq B_1$ and $N = \mathcal{B}$ the Bell measurement.

Deficiency for $\mathcal{T} \subset \mathcal{C}_2$

Let $\mathcal{T} \subseteq \mathcal{C}_2$ be

- ▶ convex
- ▶ closed
- ▶ $\mathcal{T} \circ \mathcal{T} \subseteq \mathcal{T}$

Deficiency for $\mathcal{T} \subseteq \mathcal{C}_2$

Let $\mathcal{T} \subseteq \mathcal{C}_2$ be

- ▶ convex
- ▶ closed
- ▶ $\mathcal{T} \circ \mathcal{T} \subseteq \mathcal{T}$

Then we obtain similar results with modified conditional 2-min-entropy:

$$H_{min, \mathcal{T}}^{(2)} := -\log\left(\max_{\alpha \in \mathcal{T}} \langle \gamma, \alpha \rangle\right)$$

Deficiency for $\mathcal{T} \subseteq \mathcal{C}_2$

Let $\mathcal{T} \subseteq \mathcal{C}_2$ be

- ▶ convex
- ▶ closed
- ▶ $\mathcal{T} \circ \mathcal{T} \subseteq \mathcal{T}$

Then we obtain similar results with modified conditional 2-min-entropy:

$$H_{min, \mathcal{T}}^{(2)} := -\log\left(\max_{\alpha \in \mathcal{T}} \langle \gamma, \alpha \rangle\right)$$

For characterization by guessing probabilities: restrictions on allowed pairs (α, M) of pre-processing and measurement.