

JPhysA-113511 - [View Abstract](#)

Effective methods for constructing extreme quantum observables



***Do you want to get recognition for this review on [Publons](#)?**

Don't let your reviewing work go unnoticed! Researchers the world over use Publons to effortlessly track their valuable peer review contributions for any journal. If you opt in, your Publons profile will automatically be updated to show a verified record of this review in full compliance with the journal's review policy. If you don't have a Publons profile, you will be prompted to create a free account.[Learn more about Publons](#)

☐ Yes

☒ No

GUIDE TO REVIEWER

Papers published in [Journal of Physics A: Mathematical and Theoretical](#) should contain novel results that substantially advance their relevant field. Articles that repeat or add only incrementally to already published work will not be acceptable.

QUALITY RATING

Please rate the manuscript for the originality, significance and clarity of the content. Guidance is available [here](#)

	Very Low Not publishable	Low Publishable only in an archival journal	Medium Would require major improvement to be publishable in this journal	High Likely to be publishable in this journal	Very High (Top 10%) Definitely publishable in this journal
* Originality	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>
* Significance	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>
* Clarity	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>

QUESTIONS

Please answer these questions, which are for the benefit of the Editors (the answers will not normally be transmitted to the authors). If you have any explanatory comments, please add them to the "Comments to the Editor(s)" box below.

*** Please rate the [scientific rigour](#) of the article**

☒ High ☐ Medium ☐ Low

Is this article fully or partially in your area of expertise?

☒ Fully in my area ☐ Partially in my area

Is this article better suited to the scope of another journal?

(If yes, please indicate in the "Comments to the Editor(s)" box below which journal(s))([list of all IOP journals for reference](#))

☐ Yes ☒ No

Has the author made reference to the most recent and most appropriate work?

(If no, please provide any missing references in your Comments to the authors.)

☒ Yes ☐ No

Is this work likely to be of wider interest beyond the article's immediate field?

☒ Yes ☐ No

*** Recommendation**

- ☒ Accept
- ☐ Revisions required before acceptance (only changes to the text are required)
- ☐ Major revisions required before acceptance
- ☐ Reject but may resubmit (may be publishable in this journal following these substantial revisions)
- ☐ Reject (adds little new knowledge to the field; may be publishable in an archival journal)
- ☐ Reject (contains basic errors and faulty judgements; not publishable in any journal)

COMMENTS TO THE EDITOR(S)

ΩSpecial Characters

Please enter any comments you wish to make to the Editor(s) that are not part of your report for the authors. Please bear in mind that authors can request to see these comments under the Data Protection Act 2018 if they so wish.

*COMMENTS TO THE AUTHOR(S)

ΩSpecial Characters

Please support your answers with [written comments](#) for the authors. If you choose to attach your full report as a file, please state this in the box below.

The paper contributes to the important problem of understanding the convex structure of (finite outcome) POVMs (in finite dimensions). A description of the extreme POVMs is valuable also because in many situations such POVMs correspond to optimal measurements. The results bring an insight into the possible ranks of operators in an extreme POVM and propose a method of construction of such POVMs in some cases. Although the obtained descriptions do not cover all possibilities (as the authors admit) I find the results significant enough for publication. I have only a few minor comments: - p.2, second paragraph: since the authors work only with finite outcome measurements, I think it is enough just to say that the set of POVMs is the convex hull of extreme ones -p.2, l. 14: "ranks rank" -p.4, l.2: "show" -> shows -p.5 l.4: "\eta_s" -> \eta_1 - p.6, Thm. 2: it is not clear from the introduced notation that we can have $t^r=0$ -p.6, the proof of Thm. 2: "We" -> we -many places: "compliment" -> complement -p.9 (also Fig. 1) "braking" -> breaking