maestro? voala

March 24, 2026

For Emmy Noether, in the spirit that symmetry is not decoration but structure.

Abstract

This note does not claim any classical identity of calculus, linear algebra, or geometry to be new. Its purpose is to isolate a unified *readout-frame* viewpoint: what appears as paradox, collapse, or magic in one representation may become clean structure in another.

The central technical device is the operator

$$\mathrm{CORE}_u := M_{u'} \circ U_u,$$

which packages substitution together with inner-derivative transport into a single operator object. We then illustrate the same structural slogan on two classical examples: Euler's formula, viewed as the complex readout of a real rotation, and the projection paradox of a higher-dimensional rigid object.

1 Guiding principle

The viewpoint of this note can be summarized informally as follows:

$$\textit{wrong frame} \Rightarrow \textit{fake magic}, \qquad \textit{right frame} \Rightarrow \textit{clean structure}.$$

The main claim is not that classical formulas change, but that their *visible meaning* can change dramatically when one changes the readout frame. In particular:

- a real rotation may look as if it destroys direction,
- yet the same action in the complex readout preserves direction up to phase;
- a higher-dimensional rigid object may look collapsed or contradictory,
- yet the contradiction may belong only to the projection, not to the source.

The role of the CORE notation is to compress such transport into a single operator step.

2 The CORE compression principle

We do not claim the classical chain rule itself to be new. The contribution of this note is only the introduction of a compressed operator packaging of that mechanism.

Definition 1 (Substitution and multiplication channels)**.** *Let $u = u(x)$ be a differentiable function. Define the substitution operator U_u and multiplication operator $M_{u'}$ by*

$$U_u[F](x) := F(u(x)), \qquad M_{u'}[g](x) := u'(x)\, g(x).$$

We then define the compressed operator

$$\mathrm{CORE}_u := M_{u'} \circ U_u.$$

Proposition 1 (Compressed chain rule)**.** *With the notation above, the classical chain rule can be written as*

$$D_x \circ U_u = \mathrm{CORE}_u \circ D_u.$$

Proof. Let F be differentiable. Then

$$(D_x \circ U_u)[F](x) = \frac{\mathrm{d}}{\mathrm{d}x} F(u(x)) = u'(x)F'(u(x)).$$

On the other hand,

$$(\mathrm{CORE}_u \circ D_u)[F](x) = (M_{u'} \circ U_u)[F'](x) = u'(x)F'(u(x)).$$

Thus both sides agree on every differentiable F, which proves the identity. □

So the novelty claimed here is not the underlying calculus identity, but its operator-level compression into a single readout step generated by CORE_u.

Antiderivative form

The same packaging yields the usual substitution rule for integrals. If $F'(u) = f(u)$, then

$$\int f(u(x))u'(x)\,\mathrm{d}x = F(u(x)) + C.$$

In the present notation, one may regard this informally as the statement that the integrand is the transport of f through the same compressed channel CORE_u.

Remark 1. *Thus substitution is not presented here as a two-step manipulation, but as a one-step transport mechanism generated by* CORE_u*.*

3 Euler's formula as readout of a real rotation

We now illustrate the same philosophy on a classical identity. We do not claim Euler's formula to be new. The point is only that it can be seen as the readout of a real geometric rotation in the correct frame.

3.1 The real rotation generator

Consider the matrix

$$J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

which generates rotation in $\mathbb{R}^2$.

Define the complex readout map

$$\Pi : \mathbb{R}^2 \to \mathbb{C}, \qquad \Pi(x, y) := x + \mathrm{i}y.$$

Proposition 2 (Intertwining relation)**.** *The readout map* Π *satisfies*

$$\Pi \circ J = \mathrm{i}\,\Pi.$$

Proof. For $(x, y) \in \mathbb{R}^2$,

$$J(x, y) = (-y, x),$$

hence

$$(\Pi \circ J)(x, y) = \Pi(-y, x) = -y + \mathrm{i}x = \mathrm{i}(x + \mathrm{i}y) = \mathrm{i}\,\Pi(x, y).$$

□

This is the structural core: the real rotation generator J becomes multiplication by i after the correct readout.

3.2 Exponentiation and the master step

By functional calculus,

$$\Pi \circ e^{\theta J} = e^{\mathrm{i}\theta}\Pi.$$

Now apply both sides to the basis vector

$$e_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Then

$$\Pi(e^{\theta J} e_1) = e^{\mathrm{i}\theta}\Pi(e_1) = e^{\mathrm{i}\theta}.$$

But geometrically, $e^{\theta J}$ is the real rotation by angle θ, so

$$e^{\theta J} e_1 = \begin{pmatrix} \cos\theta \\ \sin\theta \end{pmatrix}.$$

Applying Π gives

$$\Pi(e^{\theta J} e_1) = \cos\theta + \mathrm{i}\sin\theta.$$

Therefore

$$e^{\mathrm{i}\theta} = \cos\theta + \mathrm{i}\sin\theta.$$

Remark 2. *In this frame, Euler's formula is not "magic": it is the complex readout of an ordinary real rotation. The phase is geometry remembered by the operator.*

4 Direction survives as phase

The same example can be phrased in eigenlanguage.

Over $\mathbb{R}^2$, the rotation matrix J has no nonzero real eigenvectors. So in the real frame, direction appears to be destroyed.

Over $\mathbb{C}^2$, however, one has eigenvectors

$$\begin{pmatrix} 1 \\ -\mathrm{i} \end{pmatrix}, \qquad \begin{pmatrix} 1 \\ \mathrm{i} \end{pmatrix},$$

with eigenvalues

$$-\mathrm{i}, \qquad \mathrm{i}.$$

Thus the action does not destroy direction absolutely; rather, it preserves direction in the complex frame up to a phase factor.

So the slogan becomes:

in the real frame, direction is lost; in the complex frame, direction survives as phase.

5 Projection paradox and the collapsing tesseract

We now turn to a geometric analogue of the same principle.

Let $X \in \mathbb{R}^4$ be a rigid higher-dimensional object. Suppose one does not see X directly, but only a projection

$$Y = P_\theta(X),$$

where the visible image depends on a parameter θ encoding angle, frame, or orientation.

Then it may happen that

$$P_{\theta_1}(X) \neq P_{\theta_2}(X) \qquad \text{even though} \qquad X = X.$$

This is the basic projection paradox: the visible geometry may vary wildly while the source object remains unchanged.

Remark 3. *Thus projection can create contradiction without any contradiction in the source. What looks like collapse may be only readout loss.*

In informal words: a single rigid object may cast many incompatible shadows. So one should not confuse *shape* with any single *appearance*.

A useful slogan is:

$$\text{shape} \neq \text{single appearance}, \qquad \text{shape} = \text{equivalence class of projections.}$$

This is directly parallel to the phase story above: the underlying object remains coherent, while its low-level readout changes with frame.

6 Unified interpretation

We may now state the common pattern in all examples of this note.

1. A source object or action is structurally coherent.
2. A chosen readout frame makes only part of that structure visible.
3. In a poor frame, one observes loss, contradiction, or apparent magic.
4. In the correct frame, the same event becomes lawful transport.

For the chain rule:

$$\text{substitution} + \text{inner derivative} \quad \rightsquigarrow \quad \mathrm{CORE}_u.$$

For Euler's formula:

$$\text{real rotation} \quad \rightsquigarrow \quad \text{complex phase transport.}$$

For higher-dimensional geometry:

$$\text{rigid source} \quad \rightsquigarrow \quad \text{family of incompatible projections.}$$

So the unifying claim is modest but precise:

many apparent paradoxes are not contradictions in the object, but artifacts of the readout frame.

7 Optional operator channel viewpoint

The same packaging philosophy can also be applied to other operator channels. For example, if one works with an oscillatory kernel

$$f_{\mathrm{osc}}(k) := \frac{(-1)^{k+1}}{k}$$

and with an auxiliary transform $\mathcal{Z}$, one may introduce the notation

$$\mathrm{CORE}_{\mathrm{odd}}[f] := \mathcal{Z}[f_{\mathrm{osc}} f].$$

No claim is made here that the individual ingredients are new. The only point is that this notation continues the same style: a multi-step oscillatory readout may be compressed into a single operator channel.

8 Conclusion

This note does not claim any new classical theorem. Its claim is only that several standard mechanisms admit a common compressed description.

The central idea is that one should distinguish carefully between

$$\text{object}, \qquad \text{action}, \qquad \text{readout}.$$

Once this distinction is made, several apparently mysterious phenomena become structurally transparent.

In short:

$$\textit{wrong frame} \Rightarrow \textit{fake magic}, \qquad \textit{right frame} \Rightarrow \textit{clean structure}.$$

Appendix: CORE decomposition of the $\frac{di}{dt} = \ddot{i}$ meme

This appendix is intentionally humorous. It does not claim a new theorem of calculus. Its only point is to show that even a typographic joke can be read through a layered CORE-style frame.

1. The hidden source object

The visible symbol i is not taken as the primitive source object. Instead, we regard the dotless letter

$$\imath$$

as the underlying source.

The ordinary printed symbol i is then interpreted as the result of a first readout map

$$R_{\text{typ}}(\imath) := i,$$

where the dot is part of the glyph-level presentation.

2. The dynamic dot

Now introduce a second operation, namely the time-derivative transport

$$\dot{M}[g] := \dot{g}.$$

This adds a *dynamic* dot, not a typographic one.

3. The meme CORE operator

Define the humorous CORE operator

$$\text{CORE}_{\text{meme}} := \dot{M} \circ R_{\text{typ}}.$$

Applying it to the hidden source $\imath$, one obtains

$$\text{CORE}_{\text{meme}}(\imath) = \dot{M}\big(R_{\text{typ}}(\imath)\big) = \dot{M}(i) = \ddot{i}.$$

Thus the joke may be read as a two-layer transport:

$$\imath \xrightarrow{R_{\text{typ}}} i \xrightarrow{\dot{M}} \ddot{i}.$$

4. Readout interpretation

In the wrong frame, the expression

$$\frac{di}{dt}$$

looks like an ordinary derivative of the symbol i.

In the meme frame, however, the two dots are assigned different roles:

- the first dot belongs to the glyph-level readout,
- the second dot belongs to temporal dynamics.

So the punchline is not really calculus, but readout layering:

$$\text{glyph dot} + \text{time dot} = \ddot{i}.$$

5. Final slogan

In compressed form, the entire joke says

$$\text{wrong frame}: \ \frac{di}{dt}, \qquad \text{right frame}: \ \text{glyph dot} + \text{time dot} = \ddot{i}.$$

Or, in one line:

The first dot is typography, the second dot is dynamics.

[Hidden lemma: contradiction may belong to the readout] Let V and W be vector spaces, and let $\{P_\theta : V \to W\}_{\theta \in \Theta}$ be a family of readout maps depending on a parameter θ. Assume that the family is nontrivial, i.e. there exist $\theta_1 \neq \theta_2$ such that

$$P_{\theta_1} \neq P_{\theta_2}.$$

Then there exists an object $X \in V$ such that

$$P_{\theta_1}(X) \neq P_{\theta_2}(X), \qquad \text{while of course} \qquad X = X.$$

In particular, incompatible visible outputs do not imply any contradiction in the source object itself.

Proof. Since $P_{\theta_1} \neq P_{\theta_2}$ as maps, their difference

$$P_{\theta_1} - P_{\theta_2} : V \to W$$

is not the zero map. Hence there exists some $X \in V$ such that

$$(P_{\theta_1} - P_{\theta_2})(X) \neq 0.$$

Equivalently,

$$P_{\theta_1}(X) \neq P_{\theta_2}(X).$$

But this is a statement only about two different readouts of the same object X; it does not alter the identity of the source. Therefore the apparent contradiction belongs to the readout layer, not to the source layer. □

[Readout-frame paradox] A single rigid source may admit mutually incompatible visible appearances under different readout frames:

$$P_{\theta_1}(X) \neq P_{\theta_2}(X) \qquad \text{although} \qquad X = X.$$

Thus projection can create contradiction in appearance without contradiction in the object.

9 Gaussian normalization as the correct readout frame

We add one further classical example of the same slogan:

$$\text{wrong normalization} \Rightarrow \text{constant clutter}, \qquad \text{right normalization} \Rightarrow \text{self-duality}.$$

Nothing claimed here is new in analysis. The point is only structural: the same Gaussian family may look cumbersome in one scale and canonically closed in another.

9.1 The Gaussian family

For $a > 0$, consider the Gaussian

$$g_a(x) := e^{-ax^2}.$$

Its integral is the classical identity

$$\int_{-\infty}^{\infty} e^{-ax^2}\, dx = \sqrt{\frac{\pi}{a}}.$$

In particular,

$$\int_{-\infty}^{\infty} e^{-x^2}\, dx = \sqrt{\pi}, \qquad \int_{-\infty}^{\infty} e^{-\pi x^2}\, dx = 1.$$

So the factor $\sqrt{\pi}$ has not disappeared by magic; it has been absorbed into the choice $a = \pi$.

9.2 Fourier readout

Now choose the Fourier transform convention

$$\widehat{f}(\xi) := \int_{-\infty}^{\infty} f(x) e^{-2\pi i x\xi}\, dx.$$

Then the Gaussian

$$f(x) := e^{-\pi x^2}$$

has the exact self-duality property

$$\widehat{f}(\xi) = e^{-\pi \xi^2}.$$

Thus $e^{-\pi x^2}$ is the canonical Gaussian for this Fourier frame: it keeps the same visible form under the transform, with no extra constants.

Proposition 3 (Gaussian self-duality in the correct frame)**.** *With the Fourier convention above, one has*

$$\widehat{e^{-\pi x^2}}(\xi) = e^{-\pi \xi^2}.$$

Proof. Let

$$F(\xi) := \int_{-\infty}^{\infty} e^{-\pi x^2} e^{-2\pi i x\xi}\, dx.$$

Differentiate under the integral:

$$F'(\xi) = \int_{-\infty}^{\infty} e^{-\pi x^2} (-2\pi i x) e^{-2\pi i x\xi}\, dx.$$

Since

$$\frac{d}{dx} e^{-\pi x^2} = -2\pi x e^{-\pi x^2},$$

we may rewrite this as

$$F'(\xi) = i \int_{-\infty}^{\infty} \frac{d}{dx}(e^{-\pi x^2}) e^{-2\pi i x\xi}\, dx.$$

Integrating by parts, the boundary term vanishes, so

$$F'(\xi) = -i \int_{-\infty}^{\infty} e^{-\pi x^2} \frac{d}{dx}(e^{-2\pi i x\xi})\, dx = -2\pi\xi\, F(\xi).$$

Hence F satisfies

$$F'(\xi) = -2\pi\xi F(\xi),$$

so

$$F(\xi) = Ce^{-\pi\xi^2}.$$

Evaluating at $\xi = 0$,

$$F(0) = \int_{-\infty}^{\infty} e^{-\pi x^2}\, dx = 1,$$

therefore $C = 1$, and

$$\widehat{e^{-\pi x^2}}(\xi) = e^{-\pi\xi^2}.$$

□

9.3 Interpretation

The point is not merely computational. In the scale e^{-x^2}, one sees the Gaussian together with the constant $\sqrt{\pi}$. In the scale $e^{-\pi x^2}$, one sees the same family in the frame where Fourier transport closes on itself.

So the moral is exactly the same as before:

$$\text{wrong frame} \Rightarrow \text{constant clutter}, \qquad \text{right frame} \Rightarrow \text{clean invariant structure}.$$

Equivalently: the Gaussian is not changed by the transform as an object-class, but only becomes visibly self-dual in the properly normalized readout.

Remark 4. *There is no contradiction between*

$$\int_{-\infty}^{\infty} e^{-x^2}\, dx = \sqrt{\pi}$$

and

$$\widehat{e^{-\pi x^2}}(\xi) = e^{-\pi\xi^2}.$$

These are two readouts of the same Gaussian family e^{-ax^2} under different normalizations. The factor $\sqrt{\pi}$ is not lost; it is absorbed into the canonical choice $a = \pi$.

$$\boxed{e^{-x^2} \text{ gives } \sqrt{\pi}, \qquad e^{-\pi x^2} \text{ gives self-duality.}}$$