

Toward a more informative multimodal data analysis of the pediatric AML transcriptome

Jenea I. Adams


2nd year Ph.D. Student (GCB)

Xing Lab Roundtable

Monday, March 12th, 2021

Acute myeloid leukemia (AML) is the most fatal of childhood cancers with no good treatments


- Caused by accumulation of immature myeloid cells in bone marrow
- Affects 25% of children with leukemia
- Chemotherapy for acute lymphoblastic leukemia (ALL) is generally successful
 - Kills healthy and aberrant B-cells → stable prognosis



Current treatment strategies for pediatric AML are based adult genomes/transcriptomes


- Mean age of AML onset is 45 years
- "Pediatric" considered 0 years to ~30 years old
- Age-specific genomic signatures have been linked to prognosis (Bullinger *et al.*, 2017)
- Hematopoietic expansion has age-specific regulation (Bullinger *et al.*, 2017)

What about splicing?




Alternative splicing and correlated molecular pathways could help to better distinguish pediatric vs adult AML


PEGASAS correlates gene ontology and alternative splicing events



A



B



Hsu *et al.*, 2016

Hypothesis: Alternative splicing and correlated molecular pathways could play important roles in distinguishing pediatric vs adult AML

Aim 1


Improve analysis of splicing in large heterogeneous RNA-seq datasets

Aim 2



Discover age-specific, pathway-dependent alternative splicing patterns in pediatric AML RNA-seq data

Aim 1: Improve analysis of splicing in large heterogeneous RNA-seq datasets

- PEGASAS lacks an approach to make batch effect-informed correlations
- Capture and use expression heterogeneity to mitigate batch effects → Surrogate variable analysis (SVA)
- Using SVA could improve reproducibility and downstream accuracy
- Approach
 - Compare pathway-relevant exons detected with/without SVA



Aim 2: Discover age-specific, pathway-dependent alternative splicing patterns in pediatric AML RNA-seq data



Age and genetics: how do prognostic factors at diagnosis explain disparities in acute myeloid leukemia?

Manali I Patel ¹, Yifei Ma, Beverly S Mitchell, Kim F Rhoads

Affiliations + expand

PMID: 23608826 DOI: [10.1097/COC.0b013e31828d7536](https://doi.org/10.1097/COC.0b013e31828d7536)