11B Analysis with S455 Setup

Tobias Jenegger

R3B Collaboration Meeting
30. Nov. 2020

Setup and Detectors

Particle Identification

12C(p,2p)11B reaction

Summary & Outlook

Supported by BMBF 05P15WOFNA and 05P19WOFN1.

The results presented here are based on the experiment s444/s473, which was performed at the beam line/infrastructure Cave C at the GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt (Germany) in the frame of FAIR Phase-0.

TUM Members:
Roman Gernhäuser,Lukas Ponnath,Philipp Klenze,Tobias Jenegger

The S455 Setup (February 2020)



Beam energy: from 400 to 800 AMeV
Intensities: 1kHz - 1MHz
Beamtype: 12C,
Target: C, CH2

Particle Identification

$$B\rho = \frac{\beta \gamma M}{q}$$



Time of Flight Measurement: Start to TOFW

Flight-path Reconstruction: Tracking Detectors (MWPC1, 2 and 3)

Charge Measurement : TWIM Music

Flightpath Reconstruction



Radius Reconstruction:

$$R = \frac{L_{eff}}{2\sin\left(\frac{\theta_{in} + \theta_{out}}{2}\right)}$$

Known:

- position and inflight angle (θ_{in}) before GLAD
- position after GLAD (MWPC3)

compute θ_{out} iteratively:



Charge versus A/q



12C(p,2p)11B reaction



$$p_A + p_T = p_1 + p_2 + p_{A-1} + p_\gamma$$



Gamma Spectrum of 11B



Doppler Correction

$$E_{\gamma} = \gamma E_{lab}\left(1 - \beta\cos\left(\theta\right)\right)$$

Polar Angular Distribution of protons for 12C(p,2p)11B







Arzimuthal Distribution of protons for 12C(p,2p)11B



CALIFA left half (90°) iPhos

CALIFA right half (45°) iPhos

Momentum Distribution 2p & 11B



expected: barrier line

Explanation for smearing needed:

→ use simulation

→ boosting to 12C frame:

$$p_{12C} = p_i + p_{11B}$$



$$\cos(\gamma) \approx 1$$



Summary & Outlook

→ Particle Identification works out

→ Gamma spectrum and angular distribution plots look reasonable

→ Further investigations for momenta distributions of the outgoing particles needed

→ Expand analysis towards 10B isotope

Thank you!

CALIFA @ Technical University of Munich (TUM)

Roman Gernhäuser, Lukas Ponnath, Philipp Klenze, Tobias Jenegger

GEFÖRDERT VOM
Bundesministerium für Bildung und Forschung

Backup