

$^{12}\text{C}(\text{p},2\text{p})^{11}\text{B}$ Quasi Free Scattering in Inverse Kinematics at R³B

Tobias Jenegger

R3B Collaboration Meeting 2022

Setup Experiment S444

$^{12}\text{C}(\text{p},2\text{p})^{11}\text{B}$ reaction

Analysis


Summary & Outlook

TUM Members:
Roman Gernhäuser, Lukas Ponnath, Philipp Klenze, Tobias Jenegger



Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2094 – 390783311, BMBF 05P19WOFN1, 05P21WOFN1 and the FAIR Phase-0 program




Quasi Free Scattering Analysis with Experiment S444/467 (2020)



SETUP:


Beam energy: 400 AMeV
 Beamtype: ^{12}C
 Beamtime: 3 hours
 Intensities: 1kHz - 1MHz
 Target: CH_2 (12.29 mm)

$^{12}\text{C}(\text{p},2\text{p})^{11}\text{B}$ reaction




Fragment Particle Identification

- Time Measurement (START & TOFW)
- Charge Measurement (TWIM Music)
- Flightpath Reconstruction:









$$B * \rho = \frac{\beta * \gamma * M}{q}$$

Tobias Jenegger



Identification of the two correlated Protons






Event Selection Criteria:


- ^{11}B fragment identification
- Two hits (protons) with $E_{\text{hit}} > 30 \text{ MeV}$
- $\theta_1 + \theta_2 < 90^\circ$
- $\Delta\varphi = 180^\circ \pm 40^\circ$

Reconstruction of Inner Momenta

Before Scattering:



After Scattering:







(Four-)Momentum conservation relation:

$$p_{12C} + p_{tg} = p_1 + p_2 + p_{11B}$$

assuming QE scattering in mean field potential:


$$p_{12C} = p_i + p_{11B}$$

$$p_i \approx p_{missing} = p_1 + p_2 - p_{tg} \text{ (no ISI / FSI)}$$





Momentum reconstruction of ^{11}B




Before Scattering:






After Scattering:



Tobias Jenegger




Correlations between Fragment and Proton Pair




Summary

➤ Identification of the QFS-process:


^{11}B Fragment





Polar Angles of two protons



Gamma Spectrum of ^{11}B




➤ Inner Momenta and according correlation plots:




- Analyze data with other CH₂ target lengths (24.53/24 mm)
- Background subtraction with carbon target (5.4/10.86/21.98 mm)
→ Get cross section for QFS-process
- Further investigations of nuclear properties, eg. proton separation energy

$$S_p = T_{tg} - T_{p1} - T_{p2} - T_{11B} = (1 - \gamma) m_p - \gamma (T_1 + T_2) + \beta \gamma (p_{1\parallel} + p_{2\parallel}) - \frac{k^2}{2m_{11B}}$$



- S444 Commissioning Experiment can also be used for 12C – 12C inelastic scattering analysis





Thank you!

Special Thanks to:

Panin Valerii

SOFIA Group (A. Chatillon, J. Taieb, et. al.)



Tobias Jenegger

