

Hormonal-Social Adaptation Population Model: A Computational Framework for Exploring Behavioral Stabilization Under Ecological Release

Anonymous
Independent Researcher

2026

Abstract

Ecological conditions that reduce mortality pressure can alter the selective environment experienced by populations, yet most population regulation models focus on density-dependent resource competition and predation without incorporating endocrine-mediated behavioral feedbacks. We present HSAP (Hormonal-Social Adaptation Population model), an agent-based model testing whether coupled behavioral and endocrine mechanisms can influence population stability under reduced external threat. In HSAP, individuals regulate testosterone, cortisol, and estrogen in response to perceived predation risk and population density. These hormonal states drive behavioral outputs—aggression, fertility, mating drive—that feed back into population dynamics through a density-dependent behavioral sink mechanism with hysteresis. We evaluated 61 scenarios (4 core conditions, 3 sink variants, 54-factorial parameter sweep) across 1,970 validated simulation runs. Under reduced predation and disease pressure, populations stabilized at higher densities with reduced male testosterone and aggression (Cohen $d = 3.29$). Under high space constraint with abundant resources, populations declined to 31% of the resource-matched uncrowded condition, demonstrating crowding pathology through fertility suppression. Under behavioral sink conditions, populations either recovered or produced partial collapse with 14% extinction risk. A null model suite of 11 alternatives could not distinguish HSAP from logistic growth, predator-prey, or density-fertility models on population trajectories alone; HSAP became identifiable only when behavioral and endocrine observables were included. These results are simulation-consistent: under the model’s assumptions, endocrine-linked behavioral feedbacks can produce qualitatively distinct population dynamics under identical resource conditions, but population data alone are insufficient to identify the mechanistic pathways involved. The model does not establish that these mechanisms operate in natural populations; empirical validation requires paired endocrine and behavioral field measurements.

1 Introduction

1.1 Population regulation through ecological pathways

Population regulation is conventionally modeled through four classes of mechanism: resource competition, predation, disease, and density-dependent effects on fecundity and mortality [Sinclair, 1989, Hixon et al., 2002]. These frameworks—logistic growth, Lotka-Volterra predator-prey dynamics, SIR disease models, and their extensions—share a common assumption: population size is determined by external conditions acting on individuals through resource limitation and mortality risk. While these mechanisms are well-established and empirically supported, they do not fully account for a recurring pattern in field observations: populations sometimes stabilize at densities

well below resource-imposed carrying capacities, particularly following predator removal or during post-irruption phases [Krebs et al., 1995, Krebs, 2001].

1.2 Behavioral regulation as an alternative pathway

One class of explanations invokes behavioral plasticity. Individuals adjust aggression, reproductive effort, and social organization in response to perceived environmental conditions, creating feedback loops between individual physiology and population dynamics. This behavioral regulation pathway is distinct from resource-based regulation: it operates through individual decision rules rather than external mortality or fecundity limitation.

Endocrine hormones—particularly testosterone, cortisol, and estrogen—are plausible mediators of such adjustments. Testosterone influences aggression and reproductive investment across vertebrate taxa [Wingfield et al., 1990, Sapolsky, 2005]. Cortisol mediates stress responses and can suppress reproductive function [Creel et al., 2013]. Estrogen regulates female reproductive effort and social behavior [Ellison, 2003]. The causal chain from environmental perception through hormonal state to behavioral output is well-documented in individual studies, but has rarely been incorporated into population-level models.

1.3 Behavioral sink theory and crowding effects

Calhoun [1962] identified “behavioral sink” phenomena in overcrowded populations, where density-dependent behavioral deterioration—including reproductive suppression, aggression, and social withdrawal—could trap populations in unsustainable states even after density was reduced. The behavioral sink concept suggests that populations can enter self-reinforcing states of reproductive failure that persist beyond the ecological conditions that triggered them.

Subsequent work has extended this framework. Christian and Davis [1964] proposed endocrine-mediated mechanisms linking population density to reproductive suppression through stress hormone pathways. These studies converge on a common hypothesis: that behavioral and endocrine feedbacks can generate population-level regulatory effects independent of resource competition.

1.4 Existing models and the integration gap

Several modeling frameworks have explored aspects of endocrine-mediated population regulation. Endocrine-based models of stress and dominance have been applied to primate social dynamics [Sapolsky, 2005], but typically at the group level rather than as population-level regulators. Agent-based models of behavioral ecology have incorporated hormonal state variables [Higham et al., 2013], but usually without formal null model comparisons to test whether endocrine mechanisms add explanatory power beyond simpler alternatives.

The gap in the literature is a model that integrates: (1) individual-level endocrine feedback from environment to behavior; (2) sex-specific behavioral dynamics (male aggression, female competition); (3) a density-driven behavioral sink with recovery; and (4) rigorous null model comparison to test whether population trajectories alone can identify the endocrine mechanism.

1.5 HSAP contribution

We present HSAP (Hormonal-Social Adaptation Population model) as a falsifiable agent-based model that addresses this gap. HSAP proposes a specific causal chain: low external threat $\rightarrow$ male testosterone downshift $\rightarrow$ reduced male aggression, elevated female aggression $\rightarrow$ reduced fertility

→ population stabilization. The model is designed to be tested against its own predictions, not merely to demonstrate that endocrine feedbacks can produce interesting dynamics.

Our contributions are: (1) a computational framework linking individual endocrine state to population dynamics through behavioral feedbacks; (2) a demonstration that identical resource conditions produce qualitatively different population outcomes depending on behavioral and endocrine parameters; (3) an honest null model comparison showing that population trajectories alone are insufficient for mechanism identification; and (4) identification of the observable classes required for empirical testing.

Scope and boundaries. HSAP is a mammalian population model. It does not model human sexual orientation; non-reproductive sexual or social behavior is treated only as species-specific ethological behavior requiring explicit supporting data. The model is simulation-consistent, not proven; empirical validation requires paired endocrine and behavioral field measurements.

The paper is organized as follows: Section 2 describes the model. Section 3 reports results. Section 4 discusses implications and limitations. Section 5 concludes.

2 Methods

We follow the ODD (Overview, Design concepts, Details) protocol for describing individual-based models [Grimm et al., 2006].

2.1 Purpose

HSAP tests whether endocrine-linked behavioral feedbacks can generate population stabilization under reduced external threat. The model proposes that low predator pressure triggers a male testosterone downshift, reducing male aggression and reproductive effort, while female aggression increases under reduced male competition. These behavioral changes, combined with density-dependent reproductive restraint, can stabilize populations below carrying capacity through endocrine-mediated pathways rather than resource limitation alone.

2.2 Agent Architecture

Each agent is characterized by sex (assigned 50/50 at birth), age (in model time steps; 10 steps = 1 model-year), and a suite of physiological and behavioral state variables. Agents are initialized with age drawn uniformly from [5, 40) and energy, health, and rank set to 1.0, 1.0, and 0.5 respectively.

Hormonal state. Each agent carries three hormonal axes: testosterone, estrogen, and cortisol. Male testosterone is initialized near 1.0 (`base_male_testosterone`) with Gaussian noise (SD=0.15); female testosterone near 0.3 (SD=0.05). Estrogen is initialized near 1.0 for females and 0.3 for males. Cortisol is initialized near 0.5 for both sexes (SD=0.1). All hormonal values are clamped to [0, 1] at each time step.

Behavioral state. Derived from hormones and environment at each step: `aggression_tendency`, `maternal_defense_tendency` (females), `mating_drive`, `fertility`, and `offspring_survival_probability`. These are recomputed every time step; agents do not carry fixed behavioral types.

Social state. Each agent has a dominance rank (clamped [0, 1]), territory quality, `cooperation_bonus` (accumulated from cooperative actions, capped at 0.3), and a pregnancy flag with gestation count-down (`gestation_time` = 5 steps).

2.3 Environment

The environment provides five static parameters and several dynamic state variables.

Static parameters (set per scenario):

- **resource_abundance** (0 to 1): baseline resource availability
- **predator_pressure** (0 to 1): external threat level
- **disease_pressure** (0 to 1): disease prevalence
- **space_constraint** (0 to 1): controls how quickly territory availability declines with density
- **carrying_capacity** (integer): K for density computation

Dynamic state (updated each step):

- **resource_abundance**: regenerates toward maximum, consumed by population; modulated by seasonality
- **territory_availability**: $\max(0, 1 - \rho \cdot S)$, where $\rho = N/K$
- **predation_risk**: $P \cdot (1 + 0.5 \cdot \rho)$
- **disease_risk**: $D \cdot (1 + 0.5 \cdot \rho)$
- **resource_scarcity**: $\max(0, 1 - R)$, used in cortisol, aggression, and mortality equations

2.4 Endocrine Update

Hormones are recomputed for every agent at each time step, creating the causal feedback loop that defines HSAP.

Male testosterone is driven by five factors:

$$T_{m,t+1} = \min(T_{\max}, T_{\text{base}} + 0.3 \cdot \text{rank} + 0.4 \cdot M_d - 0.3 \cdot \rho - 0.4 \cdot (1 - P) - 0.2 \cdot (1 - \text{health})) \quad (1)$$

The term $-0.4 \cdot (1 - P)$ is the critical endocrine feedback: when predator pressure is high, the term is near zero (testosterone is not reduced); when predator pressure is low, the term subtracts up to 0.4 (testosterone is downshifted). This is the mechanism through which threat reduction propagates to behavioral change.

Female testosterone responds to resource scarcity and density:

$$T_{f,t+1} = \min(T_{\max}, 0.3 + 0.15 \cdot \text{rank} + 0.4 \cdot (1 - R) + 0.3 \cdot \rho) \quad (2)$$

Cortisol (both sexes) integrates density, rank, health, predation, and resource scarcity:

$$C = 0.5 + 0.1 \cdot \rho - 0.3 \cdot \text{rank} + 0.5 \cdot (1 - \text{health}) + 0.2 \cdot P + 0.3 \cdot (1 - R) \quad (3)$$

114 **Estrogen** (females) is suppressed by cortisol:

$$E = 1.0 - 0.3 \cdot C \quad (4)$$

115 2.5 Behavioral Output

116 From hormones, agents derive:

- 117 • **Aggression**: base = $T \cdot 0.5 + C \cdot 0.2$, with sex-specific additions (females: offspring defense
118 + resource competition; males: dominance bonus)
- 119 • **Fertility**: base = 0.5, modified by hormones, age, density-dependent reproductive restraint
120 (activates above density 0.7), and territory availability
- 121 • **Mating drive**: males = $T \cdot 0.5 + 0.3$; females = $E \cdot 0.4 + 0.2$
- 122 • **Offspring survival probability**: affected by testosterone, cortisol, density, and resource
123 scarcity

124 **Behavioral decision process.** At each time step, agents select one action from a fixed reper-
125 toire: forage, dominance_event, disperse, withdraw, cooperate, guard_offspring. Selection follows
126 a priority-based stochastic process:

- 127 1. **Forage** (deterministic): if energy < 0.3
- 128 2. **Guard offspring** (deterministic): if female with dependent offspring and cortisol > 0.6
- 129 3. **Dominance event**: probability = aggression_tendency $\times$ 0.3
- 130 4. **Disperse**: if density > 0.6, age > 15, probability = 0.1
- 131 5. **Withdraw**: if cortisol > 0.7, probability = $0.3 + \eta \cdot \beta_{\text{withdraw}}$
- 132 6. **Cooperate**: if energy > 0.7, probability = 0.1
- 133 7. **Forage** (default)

Table 1: Action effects on agent state

Action	Energy	Rank
Forage (success)	+0.2 to +0.4	—
Forage (fail)	$-0.05(1 + \rho)$	—
Dominance (win)	-0.15	+0.05
Dominance (lose)	-0.225	-0.05
Disperse	-0.3	-0.1
Withdraw	+0.05	-0.02
Cooperate	+0.05	—
Guard offspring	-0.1	—

134 All action effects are clamped to $[0, 1]$.

2.6 Reproduction

Mating occurs when a randomly selected male (probability proportional to `mating_drive`) encounters a fertile female above a fertility gate (default 0.3). Both must have energy above a threshold (0.4) and be in reproductive age (10–60 model-years). On successful mating, the female becomes pregnant with gestation time of 5 steps.

At birth, litter size is drawn from a normal distribution (mean=4.0, SD=1.0, floored to 1). Each pup survives with probability determined by the mother’s `offspring_survival_probability`. The mother pays an energy cost of 0.3. Newborns face an additional mortality check.

Infanticide triggers when density exceeds 0.8, with probability proportional to agent aggression and density above threshold.

Offspring neglect triggers when density exceeds 0.7 or cortisol exceeds 0.7, removing one offspring.

2.7 Mortality

Death probability per time step is computed as:

$$P_{\text{death}} = \min(1, (P_{\text{base}} + P_{\text{pred}} + P_{\text{disease}} + P_{\text{starvation}} + P_{\text{injury}} + P_{\text{old}} + P_{\text{crowd}}) \times \tau \times M) \quad (5)$$

where $\tau = 0.1$ (age increment per step) and M is the mortality multiplier (1.0 normally, 0.4 during post-sink recovery).

Table 2: Mortality components

Component	Formula
Base	0.02 (constant)
Predation	$0.05 \cdot P_{\text{risk}} \cdot (1 - 0.02 \cdot (1 - R)) \cdot \text{modifiers}$
Disease	$0.03 \cdot D_{\text{risk}} \cdot (1 + 0.5 \cdot (1 - \text{health})) \cdot (1 + 0.5 \cdot \rho)$
Starvation	If energy < 0.2: $0.3 \cdot (1 - R) \cdot (1 - \text{energy}/0.2)$
Injury	0.1 if injured
Old age	$0.01 \cdot (\text{age} - 50)$ if age > 50
Crowding	$0.05 \cdot \max(0, \rho - 0.8) + \eta \cdot M_{\text{sink}}$

Here η is the behavioral sink factor (Section 2.8) and M_{sink} is a binary mortality modifier active during sink engagement.

2.8 Behavioral Sink and Recovery

The behavioral sink is a density-driven phase transition that suppresses reproductive output even after density falls below carrying capacity.

Activation. When density exceeds `behavioral_sink_on_threshold` (default 0.75), the sink engages. A `sink_factor` is computed as a continuous measure of sink severity:

$$\eta = \min\left(1, \frac{\rho - 0.75}{1 - 0.75}\right) \quad (6)$$

Effects while active. Fertility is multiplicatively suppressed: $F \times (1 - \eta \cdot 0.85)$. Mating drive is suppressed: $M_d \times (1 - \eta \cdot 0.4)$. Withdrawal behavior and offspring neglect probabilities are increased.

Deactivation (hysteresis). The sink disengages only when density falls below `behavioral_sink_off_threshold` (default 0.50), creating a hysteresis band (on at 0.75, off at 0.50). A minimum duration of 30 steps must elapse before deactivation is permitted. An optional auto-recovery timer (`behavioral_sink_auto_recovery_c`) forces disengagement after a fixed number of steps.

Post-sink recovery. After sink disengagement, a recovery phase lasting 100 steps begins. During recovery: fertility and mating drive are multiplicatively boosted by 1.3; mortality is reduced to 40% of normal. If population drops below 30 during recovery, refugees (up to 10 per step) are injected to supplement the depleted population.

Sink Factor in the HSAP Composite Indicator. The composite indicator incorporates sink state: when the sink is active, the indicator is dampened, reflecting the disruption of normal social-endocrine regulation.

2.9 Metrics

Per time step, HSAP records: population size, density, sex ratio, resource abundance, predator pressure, disease pressure, territory availability, age structure (juvenile/adult/senior counts), mean hormones (testosterone, estrogen, cortisol, `male_T`, `female_T`), behavioral outputs (male/female aggression, female defense), reproductive metrics (fertility, births, deaths, matings, pregnancies, infanticide, neglect, refugees), sink state (active, post-recovery, `sink_factor`), external threat index, and the HSAP composite indicator.

The HSAP composite indicator is an analytical summary metric, not a directly measurable biological quantity. It is a weighted composite of five signals: low external threat (25%), low male aggression (20%), high female aggression (20%), low fertility (20%), and population stability (15%), with a viability dampener for populations below 50. The weights were chosen to reflect the model's hypothesized causal chain; they are not fit to data.

Summary metrics (computed over the full run): final population, peak population, minimum population, crash ratio (final/peak), time to stability, mean male/female testosterone, mean male/female aggression, mean fertility, mean cortisol, extinction status, and run number.

2.10 Null Models

We compared HSAP against 11 null models that lack endocrine feedbacks or behavioral mechanisms. Each null model was fit to HSAP's population trajectory and evaluated by mean squared error (MSE). Full equations are provided in the supplementary materials.

We also tested six HSAP ablation variants, each removing one component: male testosterone downshift, female aggression channel, cortisol, endocrine responsiveness, reproductive restraint, and sink recovery.

2.11 Software and Reproducibility

The model is implemented in Python 3.10+ using NumPy, pandas, Matplotlib, and tqdm. Sensitivity analysis uses SALib. Genetic algorithm search uses DEAP. The model code is available in

Table 3: Null model suite

Model	Mechanism	Key equation
N0_logistic	Logistic growth	$dN/dt = rN(1 - N/K)$
N1_predator_prey	Lotka-Volterra	$dN/dt = \alpha N - \beta NP$
N2_density_fertility	Density-dependent fertility	$F = F_0 \cdot (1 - N/K)$
N3_disease_pressure	Disease mortality	$M_d = D \cdot (1 + 0.5 \cdot N/K)$
N4_dominance_hierarchy	Rank-based allocation	No endocrine feedback
N5_random_hormone	Random hormones	Hormones $\sim U(0, 1)$
N6_resource_only	Resource-limited growth	Growth limited by R only
N7_density_recovery	Density + recovery	Density dependence + Allee term
N8_random_behavior	Random actions	Uniform action selection
N9_social_behavior	Social only	No endocrine coupling
N10_endocrine_no_behavior	Endocrine only	Hormones do not affect behavior

`src/hsap/` with full test coverage in `tests/`.

All scenario definitions are frozen in `src/hsap/scenarios.py`. Simulation outputs are frozen in `results/freezes/`. The pipeline is reproducible via:

```
python scripts/run_resumable_pipeline.py validate
python scripts/run_resumable_pipeline.py summarize
python scripts/make_paper_figures.py --dpi 300
```

2.12 Parameterization and Model Calibration

HSAP parameters were not estimated through optimization against empirical population trajectories. Instead, parameters define a computational experiment space used to examine whether coupled ecological and behavioral feedback mechanisms can generate emergent population dynamics. Biological literature informed the direction and plausibility of several mechanisms (e.g., testosterone suppresses aggression when threat is low; cortisol suppresses fertility), while numerical values function as model assumptions and scenario controls. The model contains 71 free parameters across six parameter classes (endocrine, behavioral, reproduction, mortality, environment, and GA). A full parameter table with provenance is provided in the supplementary materials.

This parameterization strategy is appropriate for the model’s purpose: HSAP is a computational exploration, not a predictive statistical model. The question is not whether specific parameter values match empirical data, but whether the class of feedback structures described by HSAP can produce qualitatively distinct population dynamics under defined assumptions.

2.13 Spatial Representation

The model uses a well-mixed population approximation. Agents are not assigned explicit spatial coordinates; interactions represent aggregate social dynamics rather than localized territorial processes. Territory availability is computed as a scalar function of density ($\tau = \max(0, 1 - \rho \cdot S)$), not as a spatially explicit resource landscape. This abstraction is appropriate for the model’s scope: HSAP tests endocrine-behavioral feedback mechanisms, not spatial ecology.

2.14 Interaction Model

Social interactions are represented as pairwise anonymous encounters rather than persistent network relationships. At each step, agents are processed in random order. Mating pairs are formed by selecting a male with probability proportional to mating drive and pairing with a randomly selected fertile female. Dominance contests occur when an agent selects the dominance event action; the opponent is not explicitly modeled (outcomes depend on the acting agent’s rank and testosterone). There is no kin recognition, no reciprocal altruism, and no reputation system. Cooperation accumulates a bounded bonus (≤ 0.3) but does not create social bonds.

2.15 Resource Scarcity

Resource scarcity is defined as:

$$\text{resource_scarcity} = \max(0, 1 - R_t) \quad (7)$$

where R_t is the current resource abundance. This scalar appears in the cortisol update, female testosterone update, aggression computation, offspring survival probability, and mortality equations. It represents the gap between current resource availability and maximum resource capacity, not a direct measure of starvation risk (which is modeled separately through the energy-based starvation mortality component).

3 Results

All results are from the validated pipeline: 1,970 validated simulation runs across 61 scenarios (4 core $\times$ 50, 3 sink $\times$ 50, 54 factorial $\times$ 30), with null/ablation suite completed for 7 core scenarios. Scenario definitions are frozen in `src/hsap/scenarios.py`. Full traceability to source data is provided in `arxiv/review/RESULTS_TRACEABILITY.md`.

3.1 Experimental Scope

The computational experiment comprised:

- **4 core scenarios** (A–D): 50 runs each, testing baseline, abundance, crowding, and high-predation conditions
- **3 sink scenarios** (C_stable, E, F): 50 runs each, testing behavioral sink activation, recovery, and partial collapse
- **54-factorial sweep**: 30 runs each, varying predator pressure (0.1/0.4/0.8), disease pressure (0.05/0.3/0.7), resource abundance (0.4/1.0/1.5), and space constraint (0.2/0.8)
- **11 null models**: fit to each core scenario’s population trajectory, evaluated by MSE
- **6 ablation variants**: removing one model component at a time
- **GA exploration**: fitness landscape mapping across parameter space

Table 4: Core scenario outcomes

Scenario	Population	Male T	Male Agg	Fertility	Extinction
A_normal_baseline	94 (89–100)	0.994	0.895	0.554	0%
B_hsap_abundance	141 (136–147)	0.986	0.847	0.568	0%
C_crowded_abundance	44 (41–46)	0.982	0.848	0.436	0%
D_high_predation	77 (72–81)	0.997	0.933	0.557	0%

3.2 Core Scenario Outcomes

All four core scenarios produced distinct population outcomes (Table 4, Figure ??). Summary statistics are reported as mean (95% CI) across 50 runs.

A_normal_baseline (predator_pressure=0.25, disease_pressure=0.20, resource_abundance=0.60, space_constraint=0.40, K=500) served as the reference condition.

B_hsap_abundance (predator_pressure=0.10, disease_pressure=0.05, resource_abundance=1.50, space_constraint=0.20, K=500) produced 50% higher population than A. Male testosterone shifted from 0.994 to 0.986, directionally consistent with relaxed competitive endocrine signaling. The 95% confidence intervals for male testosterone do not overlap between A and B (Cohen $d = 3.29$). Male aggression decreased from 0.895 to 0.847. The effect was statistically robust but modest in absolute magnitude (0.8 percentage points on a [0, 1] scale), indicating that the model’s population-level separation is not driven by a large endocrine swing alone. Notably, B’s mean fertility (0.568) is only marginally higher than A’s (0.554), suggesting that the population increase in B is not primarily driven by increased reproductive output.

C_crowded_abundance (predator_pressure=0.10, disease_pressure=0.05, resource_abundance=1.50, space_constraint=0.90, K=200) stabilized at 44 agents, 47% below baseline despite identical resource availability to B. Fertility was reduced to 0.436 compared to 0.568 in B, indicating crowding pathology through fertility suppression rather than increased mortality.

D_high_predation_survival (predator_pressure=0.45, disease_pressure=0.15, resource_abundance=0.75, space_constraint=0.40, K=500) stabilized at 77 agents, 18% below baseline. Male testosterone (0.997) and aggression (0.933) were elevated above baseline, consistent with predator-mediated endocrine state.

3.3 Population Trajectories

The four core scenarios demonstrated that identical resource conditions (B and C share resource_abundance=1.50) can produce qualitatively different population outcomes depending on space constraint and predator/disease pressure. Population trajectories showed:

- **Stabilization** (A, B, D): populations reached steady state within 200–300 model-years
- **Partial collapse** (C): population declined to 31% of the resource-matched uncrowded condition (B)
- **No extinction** in core scenarios across all 200 runs

The comparison between B and C shows that, under model assumptions, space constraint alone can suppress population to 31% of the resource-matched uncrowded condition, with the crowding pathology manifesting through reduced fertility rather than increased mortality.

3.4 Sink, Recovery, and Collapse

The behavioral sink produced qualitatively distinct dynamics depending on scenario conditions (Table ??, Figure ??).

C_crowded_stable (control; same environment as C_crowded_abundance but with behavioral sink trigger raised to 0.92) produced outcomes identical to C_crowded_abundance (pop=44, male_T=0.982, 0% sink onset). This confirms that the behavioral sink requires its density-triggered mechanism; removing the trigger eliminates sink dynamics.

E_behavioral_sink_recovery (predator_pressure=0.20, disease_pressure=0.10, resource_abundance=1.0, space_constraint=0.80, K=200, sink_on_threshold=0.25) produced the characteristic sink-recovery pattern. The sink engaged in 62% of timesteps and disengaged in 36%. Population dipped from an initial transient to a minimum of ~ 75 agents, then recovered to ~ 98 agents. The behavioral sink is reversible under these conditions: the hysteresis mechanism allowed density to fall below the deactivation threshold, triggering recovery with fertility and mating drive boosts and mortality relief.

F_behavioral_sink_partial_collapse (predator_pressure=0.10, disease_pressure=0.05, resource_abundance=0.40, space_constraint=0.80, K=200, sink_on_threshold=0.15) produced severe partial collapse. Population was reduced to ~ 4 agents (CI=[3, 5]). Sink engagement rate was 55%, but recovery rate was 0% — the behavioral sink never disengaged. Extinction occurred in 14% of runs (95% CI=[6%, 24%]). This is partial collapse with nonzero extinction risk.

3.5 Null Model Non-Identifiability

The null model comparison reveals that population trajectories alone are partly non-identifiable across the null model suite (Table ??, Figure ??).

Of 11 null models tested, 7 (logistic growth, predator-prey, density-fertility, dominance hierarchy, random hormone, density with recovery, random behavior) could not distinguish A from B using population trajectories alone (A/B MSE ratio < 1.5). The $1.5\times$ threshold is a practical classification boundary: below this ratio, the null model fits both scenarios comparably well, indicating that population data alone cannot discriminate the endocrine mechanism from simpler alternatives.

Two models showed partial separation: disease pressure (N3: A MSE=25,637, B MSE=74,023) and endocrine-no-behavior (N10: A MSE=56,787, B MSE=86,016). One model (resource-only, N6) distinguished A from B but with poor overall fit (A MSE=23,850).

This non-identifiability is one of the paper’s main results. It implies that a field study measuring population size alone cannot reject HSAP in favor of logistic growth, predator-prey dynamics, or density-dependent fertility. HSAP becomes testable only when population outcomes are paired with behavioral and endocrine observables — specifically, male testosterone and aggression measurements that null models cannot reproduce.

3.6 Ablation Analysis

The ablation analysis quantifies the contribution of each model component (Table 5, Figure ??).

Removing the male testosterone downshift channel increased B’s population MSE from 73 (full model) to 565 — a 7.8-fold increase. Removing the female aggression channel increased it to 397 (5.4-fold). These two components are the minimal endocrine mechanisms required for HSAP’s distinguishing power.

Removing endocrine responsiveness (the coupling between hormones and behavior) produced the largest MSE increase on B (2,458) compared to A (913), indicating that the endocrine-behavioral coupling is more consequential under reduced threat.

Table 5: Ablation results (B scenario MSE)

Ablation	B MSE	Fold increase	Interpretation
Full model	73	1.0×	Reference
No male T downshift	565	7.8×	Core mechanism
No female aggression	397	5.4×	Core mechanism
No cortisol	898	12.3×	Broad effect
No endocrine responsiveness	2,458	33.7×	Coupling critical
No reproductive restraint	2,293	31.4×	Density feedback
No sink recovery	0	—	Sink not activated

Removing sink recovery produced MSE=0 across all scenarios, because without the sink mechanism, the behavioral sink never engages. This is consistent with the internal consistency of the model.

3.7 Factorial Sweep

The 54-factorial analysis (predator_pressure: 0.1/0.4/0.8; disease_pressure: 0.05/0.3/0.7; resource_abundance: 0.4/1.0/1.5; space_constraint: 0.2/0.8; K=500) identifies space constraint as the dominant population regulator (Figure ??).

Under low space constraint (0.20), populations ranged from 99–143 agents regardless of predation, disease, or resource levels. Under high space constraint (0.80), populations ranged from 4–13 agents under high predation and disease, showing that crowding pathology can override resource abundance.

The factorial analysis also reveals interaction effects: high predator pressure combined with high disease pressure produced population collapse only under high space constraint, suggesting that multiple stressors are necessary for collapse when space is abundant. Extinction was rare (only 2 of 54 conditions showed nonzero extinction, both at 3.3%).

3.8 Sensitivity Analysis

The sensitivity analysis confirms that space constraint is the dominant parameter controlling population outcomes across the factorial space. Predator pressure and disease pressure have secondary effects, primarily manifesting under high space constraint. Resource abundance has the weakest independent effect on population size, consistent with the crowding pathology finding that space, not resources, limits population under crowded conditions.

3.9 Genetic Algorithm Exploration

The GA exploration mapped the fitness landscape across endocrine and behavioral parameter space. Results are presented as exploratory search, not optimization proof. The GA identified parameter regions where population stabilization is maximized, but these results should be interpreted as demonstrating the existence of favorable parameter regions, not as evidence that biological populations operate in those regions.

3.10 Negative Results and Boundaries

These results demonstrate behavior of the HSAP computational model under specified parameterizations. They do not establish:

1. That the modeled mechanisms operate in natural populations
2. That the parameter values are empirically calibrated
3. That the observed trajectories are unique to endocrine-linked behavioral regulation
4. That HSAP is the correct or only model capable of producing these dynamics
5. That the HSAP composite indicator weights are biologically meaningful

The results support the narrower claim that this class of coupled ecological-behavioral feedback model can generate population-level dynamics that differ from simpler null formulations under controlled simulation conditions. The non-identifiability result (Section 3.5) shows that population data alone are insufficient to distinguish HSAP from simpler alternatives — a finding that strengthens the case for empirical testing while simultaneously limiting the conclusions that can be drawn from simulation alone.

4 Discussion

4.1 Summary of Findings

Under the model’s assumptions, HSAP demonstrates that coupled endocrine-behavioral feedbacks can generate population-level dynamics that differ from simpler null formulations. Four core scenarios produced distinct population outcomes under identical resource conditions: the same resource abundance (1.50) produced populations of 141 (**B_hsap_abundance**) and 44 (**C_crowded_abundance**), depending on space constraint and threat level. The behavioral sink mechanism produced two distinct outcomes — recovery (E) or partial collapse (F) — depending on environmental context, not mechanism design.

These results are simulation-consistent, not proven. They demonstrate that the class of feedback structures described by HSAP is sufficient to produce qualitatively distinct population dynamics under defined assumptions. They do not demonstrate that these mechanisms operate in natural populations.

4.2 What This Paper Is Not

This paper is not:

1. **An empirical claim.** HSAP is a computational model. It does not establish that endocrine feedbacks drive population dynamics in any species. The parameter values are illustrative, not empirically calibrated.
2. **A predictive model.** HSAP does not predict population sizes, trajectories, or outcomes for specific ecosystems. It explores a class of feedback structures, not a specific parameterization.
3. **A universal explanation.** HSAP does not claim that endocrine-behavioral feedbacks are the primary mechanism of population regulation. It claims they are a candidate mechanism worth testing.

393 4. **A human behavioral model.** HSAP is a mammalian population model. It does not model
394 human sexual orientation, human social dynamics, or human population regulation. Non-
395 reproductive behavior is treated only as species-specific ethological behavior.

396 5. **An optimization result.** The genetic algorithm exploration maps fitness landscapes; it does
397 not identify optimal parameter values for biological populations.

398 The paper’s contribution is narrower: it identifies a family of coupled ecological-behavioral
399 feedback mechanisms that can produce population-level dynamics distinguishable from simpler al-
400 ternatives — but only when behavioral and endocrine observables are included alongside population
401 data.

402 4.3 The Non-Identifiability Result

403 The null model comparison is the paper’s most important result for empirical science. Seven of 11
404 null models could not distinguish HSAP from simpler alternatives (logistic growth, predator-prey,
405 density-fertility) on population trajectories alone. Population data alone are insufficient to identify
406 the mechanistic pathways driving stabilization or collapse.

407 This non-identifiability has two implications:

408 **For modelers:** Population trajectories alone cannot discriminate between endocrine-mediated
409 regulation and simpler ecological mechanisms. Any model producing reasonable population dynam-
410 ics should be tested against null alternatives before claiming mechanistic specificity.

411 **For empiricists:** Field studies measuring population size alone cannot reject HSAP in favor
412 of logistic growth, or vice versa. HSAP becomes testable only when population outcomes are
413 paired with behavioral and endocrine observables — specifically, male testosterone and aggression
414 measurements that null models cannot reproduce.

415 This is a call for richer data in conservation biology, not a dismissal of population monitoring.
416 Population data remain essential, but they are insufficient for mechanism identification.

417 4.4 Testable Predictions

418 HSAP generates three predictions for post-predator-removal populations, each falsifiable with paired
419 endocrine-behavioral data:

420 1. **Endocrine signature.** Male testosterone should decrease following predator removal, with
421 magnitude proportional to the reduction in perceived threat. This is the most direct test of
422 the endocrine feedback loop.

423 2. **Crowding pathology.** Populations in space-constrained environments should show higher
424 cortisol and lower fertility than resource-matched populations in unconstrained environments,
425 even when resource availability is identical.

426 3. **Behavioral sink persistence.** Populations under high combined stressors (predation +
427 disease + crowding) should show persistent reproductive suppression despite below-capacity
428 density, consistent with the behavioral sink mechanism.

429 These predictions require paired endocrine and behavioral field measurements. They cannot be
430 tested using population data alone — a direct consequence of the non-identifiability result.

4.5 Model Limitations

Several limitations constrain the interpretation of HSAP results:

No empirical calibration. All 71 parameters are model assumptions. The endocrine update equations, behavioral output functions, and sink mechanism are hypothesized, not empirically derived. Species-specific calibration is future work.

No spatial geography. The model uses a well-mixed population approximation. Agents are not assigned spatial coordinates. This excludes spatial processes such as habitat fragmentation, metapopulation dynamics, and localized resource depletion.

Anonymous social interactions. Social interactions are pairwise and anonymous. There is no kin recognition, no persistent social network, no reciprocal altruism, and no reputation system. This excludes network-mediated effects such as kin selection and social learning.

No seasonality. Seasonal variation was not modeled. Simulations represent normalized ecological conditions.

B_hsap_abundance fertility limitation. In scenario B (abundance), mean fertility (0.568) is only marginally higher than in scenario A (baseline, 0.554). The HSAP mechanism reduces male testosterone and aggression but does not substantially reduce female fertility under the tested parameterization. This is a limitation, not a feature.

Cohen d inflation. The reported Cohen $d = 3.29$ for male testosterone between A and B is inflated by the low variance across simulation runs. Simulation precision is not biological variance. Effect sizes should be interpreted with caution.

Mixed-depth factorial. Core scenarios use 50 runs; factorial scenarios use 30 runs. This reflects a deliberate trade-off between statistical power and parameter-space exploration.

4.6 Implications for Conservation Biology

The behavioral sink concept has practical implications for conservation. If populations can enter self-reinforcing states of reproductive failure that persist beyond the ecological conditions that triggered them, then:

1. **Population recovery may require active intervention** beyond predator control or habitat restoration. Behavioral and endocrine recovery may lag ecological recovery.
2. **Population monitoring should include behavioral and endocrine metrics.** Population size alone cannot distinguish between resource-limited regulation and endocrine-mediated behavioral sink dynamics.
3. **Multiple stressors may create irreversible states.** Under model assumptions, the partial collapse scenario (F) shows that combinations of predation, disease, and crowding can produce persistent reproductive suppression that does not resolve when individual stressors are reduced.

These implications are simulation-consistent, not proven. They suggest that conservation biology should consider endocrine and behavioral monitoring alongside population monitoring — a recommendation that requires empirical validation.

4.7 Future Work

HSAP provides a foundation for several extensions:

1. **Species-specific calibration.** Parameterize HSAP for a specific mammalian species with empirical endocrine and behavioral data.
2. **Spatial extension.** Add explicit spatial structure to test whether habitat fragmentation interacts with behavioral sink dynamics.
3. **Network extension.** Replace anonymous interactions with social network structure to test whether network topology affects endocrine-mediated regulation.
4. **Empirical validation.** Design field studies that pair population monitoring with endocrine sampling to test HSAP’s predictions in post-predator-removal systems.
5. **Mechanism decomposition.** Identify which components of the endocrine-behavioral feedback loop are necessary and sufficient for population-level effects under field-relevant conditions.

5 Conclusion

HSAP is a falsifiable agent-based model showing that endocrine-linked behavioral feedbacks can generate population stabilization, crowding pathology, recovery, or partial collapse under reduced external threat. The model’s most important finding is negative: population trajectories alone are insufficient to identify the mechanistic pathways driving these dynamics. Seven of 11 null models produced indistinguishable population outcomes. HSAP becomes testable only when population outcomes are paired with behavioral and endocrine observables.

This result has practical implications for conservation biology. Post-predator-removal monitoring programs that measure only population size may miss endocrine and behavioral signatures of regulatory change. HSAP argues for integrating endocrine sampling into population monitoring, particularly in systems where predator removal has produced unexpected population dynamics.

The model is simulation-consistent, not proven. It shows that endocrine-linked behavioral feedbacks are sufficient to produce qualitatively distinct population dynamics under identical resource conditions. Whether these feedbacks are necessary — or even present — in real populations remains an empirical question.

References

- John B. Calhoun. A behavioral sink. *Symposia of the Zoological Society of London*, 8:295–315, 1962.
- John J. Christian and David E. Davis. Endocrines, behavior, and population. *Science*, 146(3651):1550–1560, 1964. doi: 10.1126/science.146.3651.1550.
- Scott Creel, Benjamin Dantzer, Wolfgang Goymann, Dustin I Rubenstein, Mikael Sandell, and Andrew J Young. The ecology of stress: effects of the social environment. *Functional Ecology*, 27:81–88, 2013.
- Peter T. Ellison. *On Fertile Ground: A Natural History of Human Reproduction*. Harvard University Press, Cambridge, MA, 2003.
- Volker Grimm, Uta Berger, Finn Bastiansen, Sigrunn Eliassen, Vincent Ginot, Jarl Giske, John Goss-Custard, Toni Grand, Simone K. Heinz, Geir Huse, Andreas Huth, Jane U. Jepsen, Christian Jorgensen, Wolf M. Mooij, Birger Mueller, Guy Pe’er, Cyril Piou, Steven F. Railsback,

- Andrew M. Robbins, Monica M. Robbins, Eva Rossmanith, Nadja Rueger, Espen Strand, Sami Souissi, Richard A. Stillman, Rune Vabo, Ute Visser, and Donald L. DeAngelis. A standard protocol for describing individual-based and agent-based models. *Ecological Modelling*, 198(1-2): 115–126, 2006. doi: 10.1016/j.ecolmodel.2006.04.023.
- James P. Higham, Michael Heistermann, and Dario Maestripieri. The endocrinology of male rhesus macaque social and reproductive status: A test of the challenge and social stress hypotheses. *Behavioral Ecology and Sociobiology*, 67(1):19–30, 2013. doi: 10.1007/s00265-012-1420-6.
- Mark A. Hixon, Stephen W. Pacala, and Stuart A. Sandin. Population regulation: Historical context and contemporary challenges of open vs. closed systems. *Ecology*, 83(6):1490–1508, 2002. doi: 10.1890/0012-9658(2002)083[1490:PRHCAC]2.0.CO;2.
- Charles J Krebs. *Ecology: the experimental analysis of distribution and abundance*. Addison Wesley, 2001.
- Charles J. Krebs, Stan Boutin, Rudy Boonstra, A. R. E. Sinclair, J. N. M. Smith, Mark R. T. Dale, Karl Martin, and R. Turkington. Impact of food and predation on the snowshoe hare cycle. *Science*, 269(5227):1112–1115, 1995. doi: 10.1126/science.269.5227.1112.
- Robert M. Sapolsky. The influence of social hierarchy on primate health. *Science*, 308(5722): 648–652, 2005. doi: 10.1126/science.1106477.
- A.R.E. Sinclair. Population regulation in animals. In J.M. Cherrett, editor, *Ecological Concepts: the Contribution of Ecology to the Understanding of the Natural World*, pages 197–241. Blackwell Scientific Publications, Oxford, 1989.
- John C. Wingfield, Robert E. Hegner, Alfred M. Dufty, and Gregory F. Ball. The “challenge hypothesis”: theoretical implications for patterns of testosterone secretion, mating systems, and breeding strategies. *The American Naturalist*, 136(6):829–846, 1990. doi: 10.1086/285134.